The Gains of DICTO's Growth Rule

Exact growth of every closed chain of the rule T, for DICTO's solids and the classical ones



DICTO (artist, Osaka, Japan) · contact@dictospheres.com · https://dictospheres.com
Paper #80 · Version 3 · 10 November 2026 (Version 2: 31 October 2026) · DICTO DOI: 10.5281/zenodo.23822907 · Licence CC BY-SA 4.0
The ideas and names are DICTO's. Computation and proof with Claude (Anthropic).
Version 2 adds Figure 1 (two steps of the rule for the Platonic solids) and the link to the app; the text and results are unchanged.

Version 3 corrects the notes on methods: the names of our computers and private file paths are removed, with small wording fixes. No result changes.

Abstract. DICTO's growth rule T of Universal PiTet (#38) puts a face through every corner of a solid, square to the line from the centre: T(S) = {x : v·x ≤ v·v for every corner v}. Run again and again, it carries some solids back to their own shape, larger; the factor is the chain's gain. In The Truncated Icosahedron Star (#78) DICTO noticed that the four-step factors of his star chains, 4.944 and 11.542, come close to 5 and 12, and proposed that such factors might form a new series of ratios. We find every gain exactly. T is polar reciprocation of the inverted corners, and on a solid with all its corners on one sphere it is plain polar reciprocation in that sphere. The main theorem follows: when a solid and its image both have their corners on spheres, of radii R and R′, the chain closes in two steps with gain $(R'/R)^2$; for the Platonic solids this is $(R/r)^2$, circumradius over inradius squared: 9, 3 and $15 - 6\sqrt{5}$. The star chain of #78 closes with factor s^p/g, where g is the gain of T and p the period: $\varphi^4/g = 4.9443$ and $16/g = 11.5418$ for the truncated icosahedron, which explains DICTO's near-misses exactly. We give the exact gain of every closed chain among the Platonic, Archimedean and Catalan solids and of ten of DICTO's solids; all but the two omnitruncates and their duals close, in two or four steps. The gains form series by symmetry; the primes in their norms are the ones the number field allows, so a prime such as 89 or 109 in the gain of the truncated icosahedron is typical, not special. The chain of DICTO's Ogdoecontahedron passes through four of his named solids and then spreads without closing, and the cube and the dodecahedron with their edges cut back close at some depths and spread at others: depth, not the number of kinds of face, decides.

1. Introduction

Draw a plane through every corner of a solid, square to the line from the centre, and keep what lies inside all the planes. A new solid appears, with a face for every corner of the old one. Do it again, and again: some solids come back to their own shape, only larger.

DICTO's rule T (#38) grows a solid by giving it a face at each corner. Read backwards it is the star step of #78: put a spike on each face with its tip at the foot of the face's perpendicular, scaled by a factor s, and wrap the tips; then T of the wrapped solid is s times the solid one started from (#78, Section 6). So every star chain is a chain of T, and a closed chain of T is a closed star chain.

The rule closes for the icosidodecahedron and DICTO's GTI with the gain $10/\varphi^4$ (#38) and for the truncated icosahedron with a gain whose norm carries the primes 89 and 109 (#78). This paper finds the gains of all closed chains exactly, explains them where a proof is possible, and tests DICTO's proposal that they form a new series of ratios.

2. The rule as polar reciprocation

For a radius ρ, the polar of a solid X in the sphere of radius ρ about the centre O is $X^\rho = \{x : y\cdot x \le \rho^2$ for every y in X$\}$; for a convex solid only its corners matter. Polar reciprocation is classical: the polar of the polar is the solid itself, and X^a taken in a second sphere gives $(X^a)^b = (b^2/a^2)\cdot X$.

> **Lemma 1.** (a) T(S) is the polar, in the unit sphere, of the solid whose corners are the inverted corners $v/|v|^2$ of S. (b) If all corners of S lie on one sphere of radius R, then T(S) is the polar S^R of S in that sphere. (c) $T(k\cdot S) = k\cdot T(S)$ for every $k > 0$.

Proof. $v\cdot x \le v\cdot v$ is the same as $(v/|v|^2)\cdot x \le 1$, which gives (a); if $|v| = R$ for every corner, it reads $v\cdot x \le R^2$, which is (b); and (c) holds because $kv\cdot x \le k^2 v\cdot v$ is $v\cdot(x/k) \le v\cdot v$. ∎

In the plane, T is the classical antipedal (negative pedal) polygon with pedal point O, and part (a) is the polygon form of the known statement that the negative pedal is the polar dual of its inverse (L. G. Gheorghe 2020, stated there as classical). B. M. Stewart proved in 1940 that n antipedal steps bring any n-gon back to a similar one. In space each step turns corners into faces, so the number of corners changes from step to step, and the chains below close in two or four steps whatever the number of corners, or not at all.

3. Main theorem: two spheres give period two

If a solid's corners sit on one sphere and so do the corners of its image, the rule is a mirror in a sphere, twice: two steps bring the solid back, larger by the square of the ratio of the two spheres.

> **Theorem.** Let S be a convex solid with its centre inside, all its corners on a sphere of radius R, and let all the corners of T(S) lie on a sphere of radius R′. Then $T(T(S)) = (R'/R)^2\cdot S$: the chain closes in two steps (or one) with radius gain $(R'/R)^2$.
>
> **Corollary (the Platonic solids).** For a Platonic solid with circumradius R and inradius r, T(S) is its dual, touching the sphere of radius R with its faces, and the chain closes in two steps with gain $(R/r)^2 = \tan^2(\pi/p)\cdot\tan^2(\pi/q)$ for the Schläfli symbol {p, q}: 9 for the tetrahedron, 3 for the cube and the octahedron, $15 - 6\sqrt{5}$ for the dodecahedron and the icosahedron.

Proof. By Lemma 1(b), $T(S) = S^R$, and since the corners of T(S) lie on the sphere of radius R′, $T(T(S)) = (S^R)^{R'} = (R'^2/R^2)\cdot S$. For the corollary: the polar of a regular solid in its circumsphere is the dual solid with inradius R. The ratio of circumradius to inradius is the same for a solid and its dual (both equal $\tan(\pi/p)\tan(\pi/q)$), so the dual's corners lie on the sphere of radius $R' = R\cdot R/r$, and the gain is $(R'/R)^2 = (R/r)^2$. ∎




Figure 1. Two steps of DICTO's rule T, each panel at its own scale. Left: S. Middle: T(S), see-through, round S; every corner of S lies on a face of T(S). Right: T(T(S)), see-through, round T(S) (violet): S again, g times larger. The tetrahedron goes to a tetrahedron turned about, with g = 9; the cube to the octahedron, g = 3; the dodecahedron to the icosahedron, g = 15 − 6√5 ≈ 1.584. Every relation drawn was checked numerically to 10^{-9} (script fig_paper_c.py).

The theorem needs both solids to have their corners on one sphere. A uniform (Archimedean) solid A has its corners on one sphere, so T(A) is its Catalan dual; but the Catalan dual's corners lie on as many spheres as A has kinds of face, and the theorem does not apply. These chains close in four steps, as Section 5 shows; the theorem covers exactly the chains of period two among the classical solids. It also covers DICTO's David and Delilah, whose corners at every step form a single orbit of their symmetry group and so lie on one sphere; David's gain is 10 and Delilah's 3.

4. The star chain and DICTO's near-misses

DICTO noticed that the four-step factors of the star chain of the truncated icosahedron in #78, 4.944 for golden spikes and 11.542 for doubled ones, are close to 5 and 12, and asked whether such factors form a series.

> **Proposition 2.** Let a star chain grow by spikes whose tips lie at s times the feet of the face perpendiculars, and let its solids lie on a closed chain of T with period p and radius gain g. Then the star chain closes with the same period, and its p-step factor is s^p/g.

Proof. By #78, $T(S_{n+1}) = s \cdot S_n$, so with Lemma 1(c), $T^p(S_{n+p}) = s^p \cdot S_n$. On the chain T^p multiplies by g, so $S_{n+p} = (s^p/g) \cdot S_n$. ∎

For the truncated icosahedron p = 4 and g = (17745 − 7712√5)/361 = 1.3862708, so the factors are $\varphi^4/g = 4.9442735$ and $2^4/g = 11.5417565$, exactly the numbers DICTO saw. They are not 5 and 12; the nearness comes from g lying close to 1.4, and the numbers that carry the structure are the gains g themselves, which the next section lists.

5. The gains

Every Platonic, Archimedean and Catalan solid but four comes back under the rule, and so do ten of DICTO's solids. The growth factors are exact numbers built from √5, √2 or $\sqrt[3]{2}$.

Solid (Archimedean / Catalan dual)	Period	Radius gain, exact	Decimal	Norm
tetrahedron	2	9	9	3^2
cube, octahedron	2	3	3	3
dodecahedron, icosahedron	2	15 − 6√5	1.583592	$3^2 \cdot 5$
truncated tetrahedron / triakis tetrahedron	4	121/25	4.84	$11^2/5^2$
cuboctahedron / rhombic dodecahedron, truncated octahedron / tetrakis hexahedron (one chain)	4	5/2	2.5	5/2
truncated cube / triakis octahedron	4	271 − 190√2	2.299420	17·73
rhombicuboctahedron / deltoidal icositetrahedron	4	(2281 − 1556√2)/49	1.642521	$17 \cdot 433/7^2$
icosidodecahedron / rhombic triacontahedron; DICTO GTI, FPD, Palintriacontahedron	4	$35 - 15\sqrt{5} = 10/\varphi^4$	1.458980	$2^2 \cdot 5^2$
truncated icosahedron / pentakis dodecahedron (#78)	4	(17745 − 7712√5)/361	1.386271	$5 \cdot 89 \cdot 109/19^2$
truncated dodecahedron / triakis icosahedron	4	(5179 − 2238√5)/121	1.443640	$61 \cdot 241/11^2$
rhombicosidodecahedron / deltoidal hexecontahedron	4	(32269 − 13820√5)/1089	1.254857	$41 \cdot 17401/(3^4 \cdot 11^2)$
DICTO Icosaxes	4	65 − 28√5	2.390097	5·61
DICTO Unborn Axes	4	(72 − 27√5)/5	2.325233	$3^4 \cdot 19/5^2$
DICTO Tweedledum	4	293 − 652√5/5	1.416736	41·101/5
DICTO Metaxyhexecontahedron	4	939725/9 − 140084√5/3	1.440036	$5 \cdot 2081 \cdot 2341/3^4$
DICTO Samson (enters after one step)	4	squared gain in $\mathbb{Q}(\sqrt[3]{2})$, cubic	3.855518	—
DICTO David	2	10	10	2·5
DICTO Delilah	2	3	3	3

Table 1. The closed chains of T. The classical solids are built exactly as orbits of a seed corner under their symmetry group and checked uniform; each chain enters its cycle at once except

Samson's. An Archimedean solid and its Catalan dual share one chain. Norm: for an irrational gain the field norm $g \cdot g'$, factored; for a rational gain the gain itself. Samson's squared gain is $-8732015647/174900625 - (680143024/174900625)\sqrt[3]{2} + (7678489792/174900625)\sqrt[3]{4}$.

6. Chains that do not close

The two omnitruncates, the truncated cuboctahedron and the truncated icosidodecahedron, and their duals do not close in eight steps: the number of corners keeps rising (from 26 to 552 and from 62 to 812), with new kinds of corner at every step. They are the only Archimedean solids with three kinds of face.

The chain of DICTO's Ogdoecontahedron (OG) passes exactly through four of his named solids, the KRE, PHE, TPP and DPHP (#39), and then leaves the data set: 452, 642, 1220, 1352 and 2240 corners at steps five to nine, with the number of different corner radii rising (3, 2, 4, 4, 6, 10, 14, 19, 24). A step similar to an earlier one would need the same number of corners, so the chain does not close within nine steps, and the rising number of corner kinds shows it spreading rather than settling. The OG's corners lie on mirror planes, as the closing solids' do, so lying on mirrors is not enough for a chain to close.

Depth decides, not the number of plane families. We swept the solids whose faces lie on two kinds of symmetry axis, exactly. Truncated icosahedra, cubes and octahedra (faces on the 5- and 3-fold or the 4- and 3-fold axes) close at every depth tested. The cube with its edges cut back (planes $x = \pm 1$ and $(\pm 1, \pm 1, 0) \cdot x \le 2s$) shows three regimes: its chain closes in four steps for $s \le 3/5$ and for $s \ge 3/4$, and spreads for every tested s between (twelve values from 0.6125 to 0.7375, among them $1/\varphi$). At $s = 3/5$ all 32 corners lie on one sphere; above it the image $T(S)$ has corners off the symmetry axes, and from 3/4 on its 26 corners all lie on axes again. The dodecahedron with its edges cut back behaves the same way: it closes for $s \le 43/40$ and $s \ge 5/4$ and spreads for $87/80 \le s \le 99/80$. Conversely, a solid with the faces of the truncated cuboctahedron, cut at other depths, closes in four steps, though the Archimedean truncated cuboctahedron does not.

7. The primes

> **Proposition 3.** The squared gain of a closed chain lies in the number field of the solid's coordinates. For an irrational gain g in $\mathbb{Q}(\sqrt{5})$, every prime that divides the norm $N(g) = g \cdot g'$ to an odd power is 5 or a prime $p \equiv \pm 1 \pmod 5$; in $\mathbb{Q}(\sqrt{2})$ it is 2 or $p \equiv \pm 1 \pmod 8$.

Proof. T only intersects half-spaces whose data are built from the coordinates, so every corner of every step lies in the field, and so does the ratio of squared radii of two similar steps. In a quadratic field a prime p that stays prime (is inert) has norm p^2, so the power of p in the norm of any element is even; the primes that may appear to an odd power are those that split or ramify, and by quadratic reciprocity these are 5 and $p \equiv \pm 1 \pmod 5$ in $\mathbb{Q}(\sqrt{5})$, 2 and $p \equiv \pm 1 \pmod 8$ in $\mathbb{Q}(\sqrt{2})$ (standard: e.g. TCD number theory notes, chapter 15; Goldmakher's notes on the Kronecker symbol). ∎

All the primes in Table 1 obey this: 89 and 109 (truncated icosahedron), 61 and 241 (truncated dodecahedron), 41 and 17401 (rhombicosidodecahedron), 17, 73 and 433 (the octahedral ones), and in DICTO's solids 61 (Icosaxes), 19 (Unborn Axes), 41 and 101 (Tweedledum), 2081 and 2341 (Metaxyhexecontahedron). The inert prime 3 appears only to the fourth power (Unborn Axes, the Metaxyhexecontahedron, the rhombicosidodecahedron's

denominator). Rational gains (9, 3, 5/2, 121/25, and David's 10 and Delilah's 3) fall outside the proposition, which speaks only of irrational gains. So the primes are fixed by the field and carry no further meaning; DICTO's proposed series is found in the gains themselves.

8. The series

- **Platonic:** $(R/r)^2 = \tan^2(\pi/p)\tan^2(\pi/q)$, proved (Section 3): 9, 3, $15 - 6\sqrt{5}$.
- **Quasi-regular:** the cuboctahedron's chain has gain 5/2 and the icosidodecahedron's $10/\varphi^4$; both are $10/(2\cos(\pi/n))^4$, with n = 4 ($2\cos(\pi/4) = \sqrt{2}$) and n = 5 ($2\cos(\pi/5) = \varphi$). The same expression at n = 3 gives 10, which is the gain of DICTO's David; with only two cases and one coincidence this is recorded as an observation, not a theorem.
- **DICTO's golden solids:** the GTI, the FPD and the Palintriacontahedron lie on the chain of the icosidodecahedron and the rhombic triacontahedron and share its gain $10/\varphi^4 = 35 - 15\sqrt{5}$ (#38).
- **Archimedean and Catalan:** each pair shares one gain (T alternates between them); the octahedral gains lie in $\mathbb{Q}(\sqrt{2})$, the icosahedral in $\mathbb{Q}(\sqrt{5})$, and the truncated tetrahedron's is $121/25 = (11/5)^2$.

9. What is new and what is known

- **Known, credited:** DICTO's rule T and the gain $10/\varphi^4$ of the icosidodecahedron's chain (#38); the star chain of the truncated icosahedron and its relation to T (#78); polar reciprocation; in the plane, the antipedal polygon and Stewart's theorem that n steps return an n-gon to its shape (B. M. Stewart, "Cyclic properties of Miquel polygons", Amer. Math. Monthly 47 (1940) 462–466, read as quoted by N. McDonald, R. Garcia and D. Reznik, "Exploring the dynamics of the circumcenter map", arXiv:2202.02551, 2022); the negative pedal as the polar of the inverse (L. G. Gheorghe, arXiv:2012.12385, 2020); the fourth pedal tetrahedron similar to the first (A. Kostanyan, 2026, abstract read); the ratio R/r of the Platonic solids; the splitting of primes in quadratic fields. G. W. Hart's canonical polyhedra also iterate reciprocation, but converge to a fixed shape and measure no gains.
- **New, as far as the sources checked show:** the two-sphere theorem for T in space and its Platonic corollary; the star-chain factor s^p/g, which accounts exactly for DICTO's near-misses; the exact gains of all closed chains among the Platonic, Archimedean and Catalan solids and ten of DICTO's solids; the failure of the omnitruncates to close.
- **Not decided here:** a proof that the Archimedean chains close in four steps; the exact edges of the open windows of the edge-cut cube and dodecahedron; whether the omnitruncates' chains ever close or settle; the quasi-regular expression beyond its two cases.

In the app. Polyhedraverse, Star step (polyhedraverse.dictospheres.com/?open=starstep:CUBE:2), spikes any convex solid and wraps it, one tap per step: the star chain, which is the chain of T read backwards.

10. Conclusion

DICTO's growth rule is a mirror in a sphere, applied to the inverted corners. Where the corners of a solid and of its image lie on spheres, two steps of the rule close the chain, and the gain is the square of the ratio of the spheres: for the Platonic solids, circumradius over inradius, squared. The near-5 and near-12 factors DICTO saw in the TI Star are s^4/g exactly,

the spike factor to the fourth over the gain of the rule. The gains of all closed chains are exact numbers in $\mathbb{Q}$, $\mathbb{Q}(\sqrt{2})$, $\mathbb{Q}(\sqrt{5})$ or $\mathbb{Q}(\sqrt[3]{2})$, arranged by symmetry into series; their primes are the ones the field allows. DICTO's idea of a new series of ratios is answered by the gains, not by their primes.

> **Sources checked** (30–31 October 2026; every query and page with its outcome is listed in study 2026-10-30-star-cores-exact, prior-art-study6.md). *Reference library* (ask-library, 18 queries, e.g. "iterated reciprocation of a polyhedron", "pedal polyhedron", "canonical polyhedra iteration", "circumradius inradius Platonic ratio"): no source on iterating T or its gains. *Web, read:* McDonald, Garcia and Reznik 2022; Gheorghe 2020; Wikipedia and MathWorld, "Negative pedal curve", "Pedal curve", "Pedal triangle", "Antipedal triangle", "Platonic solid", "Quadratic field"; D. Hatch, "A canonical reciprocation center for convex objects" (2003); Hart's canonical-polyhedra supplement; TCD number theory notes ch. 15; L. Goldmakher, "Legendre, Jacobi, and Kronecker symbols". *Abstract or snippet only:* Stewart 1940 (through McDonald et al.), Kostanyan 2026, Hsu et al. 2021. DICTO's papers #38 and #78 are cited for their results.

> **Data and code.** Study 2026-10-30-star-cores-exact: scripts/study1.py (DICTO's ten solids), study3.py (two plane families, swept), study4.py (the Ogdoecontahedron's chain), study5.py (the classical solids, built exactly as orbits), gains.py (minimal polynomials and norms); data/study1.json, gains.txt, study3.json, study3-fine.json, study4.json, study5.json. Exact arithmetic in $\mathbb{Q}$, $\mathbb{Q}(\sqrt{2})$, $\mathbb{Q}(\sqrt{5})$ and $\mathbb{Q}(\sqrt[3]{2})$ with the data set's exactlib (DICTO Solids: Exact Geometry v6, 10.5281/zenodo.23790799). Every return is checked by matching each corner of the later step to a parallel corner of the earlier one with the exact squared ratio.