The Delian Bond: Samson, Delilah and the Octet Between Kepler's Rhombic Dodecahedron and the Cube



Theorem: one flat cut, slid through every tetrahedron–octahedron pair of the octet, gives a family of pairs of convex solids that fill space, from Kepler's rhombic dodecahedron through the octet to the cube; at one cut the two solids, DICTO's Samson and Delilah, have exactly equal volume, twice the octet's tetrahedron and half its octahedron, and two other cuts make every edge equal

DICTO (artist, Osaka, Japan) · contact@dictospheres.com · https://dictospheres.com
Paper #51 · Version 1 · 24 October 2026 · DICTO DOI: reserved on publication · Licence CC BY-SA 4.0 · Scripts, exact checks and figures (Shadow forms, octet): https://github.com/DICTOR-Master/kaleidohedra/tree/099b211/studies/2026-10-23-shadow-forms (research branch, commit 099b211; also attached to this record)
The ideas are DICTO's: to do for the tetrahedron, octahedron and cube what paper #48 did for the icosahedron, to find the special positions first and then name them, and the names Samson, Delilah, the DICTONIC SOLIDS and the DICTONIC BONDS; computation, exact proofs and drafting with Claude (Anthropic).

In short. Regular tetrahedra and octahedra fill space together: the octet. Join each tetrahedron's centre to each octahedron's centre through the face they share, and the octet falls into double cones, one per shared face. Cut every double cone by one flat plane parallel to that face, and give each side to its own centre. Slide the cut, and the tetrahedra grow while the octahedra shrink, or the other way: every position still fills space with two convex solids. At one end are Kepler's rhombic dodecahedra, a third of the way along the octet, at the other end cubes. At one position the two solids have exactly the same volume: the tetrahedron's has doubled and the octahedron's halved, the doubling of the ancient Delian problem, in both directions at once. DICTO named this pair Samson and Delilah.

Main theorem. For every cut position t from 0 to 1 the two pieces are convex and fill space face to face; for $t \le 1/3$ they are a regular tetrahedron of edge $3t$ and a rhombic dodecahedron with its eight three-fold corners cut, for $t \ge 1/3$ a regular octahedron of edge $a = 3(1 - t)/2$ and a cube with four alternate corners cut, with exact volumes.

Theorem 2: at $t = (2\sqrt{6} - 3)/15$ and $t = 5/3 - (2/3)\sqrt{2}$ every edge of the filling has the same length, $(2\sqrt{6} - 3)/5$ and $\sqrt{2} - 1$ (octet edge 1); at $t = 1 - \sqrt[3]{6}/3$ the tetrahedron pieces and the octahedron pieces each fill half of space. **Theorem 3 (the Delian bond):** at $t = 1 - (2/3)\cdot 2^{-1/3}$, and only there, the two pieces have equal volume; Samson, the cut cube, has twice the volume of the octet's tetrahedron, and Delilah, the octahedron, half the volume of the octet's octahedron: it is that octahedron shrunk by $1/\sqrt[3]{2}$. Exact in the number fields $\mathbb{Q}(\sqrt{6})$, $\mathbb{Q}(\sqrt{2})$, $\mathbb{Q}(\sqrt[3]{2})$ and $\mathbb{Q}(\sqrt[3]{6})$. With DICTO's names for the six DICTONIC SOLIDS and three DICTONIC BONDS, and a remark on the polar pair of paper #48.




Figure 1. The Delian bond. From left: Samson (gold) from two sides, a cube with four alternate corners cut so deep that it looks like a clipped tetrahedron; Delilah (blue), a regular octahedron; the two face to face. Equal volume, √2/6 each at octet edge 1. Each panel at its own scale.

1. Where it started

> *After the icosahedron's golden split, DICTO proposed doing the same for the three Platonic solids that build the octet. The answer turned out to be a slider with the octet in the middle.*

Paper #48 [1] cut the thin AXE pieces between icosahedra at the golden point and turned a lattice of icosahedra and its holes into two convex truncated octahedra; its Theorem 4 placed that pair in a one-parameter family of mixed space-fillers, and a remark found the octet in a similar family. On 23 October 2026 DICTO proposed the same approach for the tetrahedron, the octahedron and the cube, and, when the family appeared, to find its special positions first and name them afterwards. He named the equal-volume pair Samson (the cut cube) and Delilah (the octahedron), from "Delian", and the six convex and keyed solids of #48 and this paper together the DICTONIC SOLIDS, held in three DICTONIC BONDS. The Delian position also meets an earlier idea of DICTO's, doubling a solid's volume by the cube root of 2.

2. The octet as double cones

> *The octet is built from tetrahedra and octahedra. Each of them is a bundle of cones, one from its centre over each face. Two cones meet across each shared face and make a double cone, and the double cones fill space too.*

Take the octet of edge 1 (the tetrahedral-octahedral honeycomb [3]; Fuller's octet truss [9]): regular tetrahedra, of volume $V_T = \sqrt{2}/12$, and regular octahedra, of volume $V_O = \sqrt{2}/3 = 4V_T$, two tetrahedra for each octahedron. Each tetrahedron is the union of the 4 cones from its centre over its faces, each octahedron the union of the 8 cones from its centre over its faces, and every face is shared by one tetrahedron and one octahedron. These cones are the face pyramids of the tetrahedron and the octahedron, which fill space together (paper #44 [12]). For a shared face F let B_F be the double cone formed by its two cones, from the tetrahedron's centre c_T through F to the octahedron's centre c_O. The double cones fill space without overlap, since the cones do. The segment c_Tc_O is perpendicular to F and meets it at its centre; F lies one third of the way from c_T to c_O (the tetrahedron's inradius is a third of the sum of the two inradii, $1/(2\sqrt{6})$ against $1/\sqrt{6}$).

Two classical facts are used. Kepler's **rhombic dodecahedron** [13] is an octahedron with the 8 tetrahedron cones over its faces added: its 8 three-fold corners are the centres of the neighbouring tetrahedra (volume $V_O + 8{\cdot}V_T/4 = 6V_T$), and these rhombic dodecahedra fill space [3, 4]; equally, the rhombic dodecahedron is a cube with six square pyramids added [2, 4]. The **cube** of edge $1/\sqrt{2}$ is a tetrahedron with the 4 octahedron cones over its faces added:

its other four corners are the centres of the neighbouring octahedra, each corner piece is an eighth of an octahedron (volume $V_T + 4 \cdot V_O/8 = 3V_T$), and these cubes fill space; this dissection of the cube is elementary.




Figure 2. One double cone, from a tetrahedron's centre through the shared face to an octahedron's centre. Left: whole. Middle: cut by a plane parallel to the face (here t = 0.55, beyond the face). Right: the tetrahedron's share (gold) and the octahedron's share (blue), pulled apart.

3. The family

> *Cut every double cone at the same fraction of the way from the tetrahedron's centre to the octahedron's, and give each side to its own centre. Whatever the fraction, the pieces fill space; what they look like depends on whether the cut is before or after the shared face.*

> **Main theorem.** For $0 \le t \le 1$ cut every double cone B_F by the plane parallel to F through the point $c_T + t(c_O - c_T)$, and let T(t) be the union of the 4 parts next to a tetrahedron's centre and O(t) the union of the 8 parts next to an octahedron's centre. Then the pieces T(t) and O(t) fill space face to face, both are convex, and:
> (a) for $0 < t \le 1/3$, T(t) is the regular tetrahedron of edge 3t, volume $27t^3 \cdot V_T$, and O(t) is the rhombic dodecahedron with its 8 three-fold corners cut off by planes parallel to the octahedron's faces (8 triangles and 12 hexagons for $t < 1/3$), volume $(6 - 54t^3) \cdot V_T$;
> (b) for $1/3 \le t < 1$, with $a = 3(1 - t)/2$, O(t) is the regular octahedron of edge a, volume $4a^3 \cdot V_T$, and T(t) is the cube with its 4 corners at octahedron centres cut off (4 triangles and 6 hexagons for $t > 1/3$), volume $(3 - 2a^3) \cdot V_T$;
> (c) at $t = 0$ the pieces are the rhombic dodecahedra alone, at $t = 1/3$ the octet, at $t = 1$ the cubes alone.

> *Why: every double cone is shared out by one flat cut, so nothing is lost and nothing overlaps. Before the face, the tetrahedron's side is a smaller copy of the tetrahedron, and the octahedron's side is the rhombic dodecahedron with its tips cut off; after the face, it is the other way round with the octahedron and the cube.*

Proof. The double cones fill space and each is divided by one plane into two parts with disjoint interiors, so the pieces fill space. (a) For $t \le 1/3$ the plane lies in the tetrahedron's cone, at fraction 3t of the way from c_T to F; the 4 parts next to c_T are the cones scaled by 3t about c_T, which make the tetrahedron scaled by 3t. The parts next to c_O are its 8 cones together with the 8 neighbouring tetrahedron cones with their tips removed, that is, the rhombic dodecahedron (§2) cut by the 8 planes, since each plane lies between c_T and F and c_T

is the only corner of the rhombic dodecahedron beyond the plane of F, so the plane meets the rhombic dodecahedron only inside the one cone over F: a convex set, being an intersection of convex sets. Each plane cuts the three edges at a three-fold corner at the same fraction 3t of their length, so for $t < 1/3$ the corners become triangles and the rhombi hexagons. (b) For $t \geq 1/3$ the plane lies in the octahedron's cone, at fraction $(3/2)(1 - t) = a$ of the way from c_O to F, and the same argument with the roles exchanged gives the octahedron scaled by a and the cube (§2) cut by 4 planes (c_O being the only corner of the cube beyond the plane of F). The volumes follow: (a) $27t^3V_T$, and $6V_T - 8\cdot(V_T/4)\cdot 27t^3$; (b) $a^3V_O = 4a^3V_T$, and $3V_T - 4\cdot(V_O/8)\cdot a^3$. Face to face: a triangle of one piece is exactly the cut face of its neighbour of the other kind, and the hexagons are shared by neighbours of the same kind, whose cuts agree because the corners they cut are the same points c_T (respectively c_O), cut by the same planes. The study checks all of this exactly, in rational arithmetic, at $t = 1/10, 1/5, 1/3, 1/2, 4/5, 9/10$ [5]. ∎




Figure 3. The family. Top: the tetrahedron piece T(t); bottom: the octahedron piece O(t); each panel at its own scale. From $t \approx 0$ (a small tetrahedron and a rhombic dodecahedron with slightly cut corners) through the first equilateral member, the octet ($t = 1/3$), the Delian bond (Samson and Delilah) and the second equilateral member, to $t \approx 1$ (the cube and a small octahedron).

4. Special members

> *Along the slider, a few positions stand out: two where every edge in the whole filling has the same length, and one where the tetrahedron pieces and the octahedron pieces each take exactly half of space.*

> **Theorem 2.** (a) At $t = (2\sqrt{6} - 3)/15$ every edge of the filling, of the tetrahedra and of the cut rhombic dodecahedra alike, has length $(2\sqrt{6} - 3)/5$. (b) At $t = 5/3 - (2/3)\sqrt{2}$ every edge, of the cut cubes and of the octahedra alike, has length $\sqrt{2} - 1$. (c) At $t = 1 - \sqrt[3]{6}/3$ the tetrahedron pieces and the octahedron pieces each fill exactly half of space. (d) For no t is a hexagon regular.

> *Why: each piece has only two kinds of edge, the ones the cut makes and what is left of the old edges. Setting the two equal is one equation, and it has one answer on each side of the octet.*

Proof. (a) For $t \leq 1/3$ the tetrahedron has edge 3t, and the cut triangles on the rhombic dodecahedron are its faces. The other edges of the cut rhombic dodecahedron are what is left of its edges, of length $\sqrt{6}/4$, after a fraction 3t at the cut end: $(\sqrt{6}/4)(1 - 3t)$. Equal when $3t = (\sqrt{6}/4)(1 - 3t)$, that is, $t = \sqrt{6}/(12 + 3\sqrt{6}) = (2\sqrt{6} - 3)/15$, where $3t = (2\sqrt{6} - 3)/5$. (b) For $t \geq$

1/3 the octahedron has edge a and the cut cube's triangles are its faces; the rest of each cube edge, of length $1/\sqrt{2}$, is $1/\sqrt{2} - a/\sqrt{2}$ (each edge of the cube has one cut end, since the cut corners alternate). Equal when $a = (1 - a)/\sqrt{2}$, that is, $a = \sqrt{2} - 1$, $t = 1 - 2a/3 = 5/3 - (2/3)\sqrt{2}$. (c) Per octahedron there are two tetrahedra; their pieces take half of space when $2(3 - 2a^3) = 4a^3$, $a^3 = 3/4$, $t = 1 - (2/3)\sqrt[3]{(3/4)} = 1 - \sqrt[3]{6}/3$ (for $t \le 1/3$ the tetrahedron pieces take at most a third). (d) The hexagons of the cut cube lie in the cube's faces and keep its right angles at their two uncut corners (angles 90°, 135°, 135°, 90°, 135°, 135°); those of the cut rhombic dodecahedron keep its acute rhombus angles of about 70.5° (with 144.7° at the cut corners). The study confirms (a)–(c) exactly in $\mathbb{Q}(\sqrt{6})$, $\mathbb{Q}(\sqrt{2})$ and $\mathbb{Q}(\sqrt[3]{6})$: every edge equal, and the volumes, with no floating point in any equality [5]. ∎

5. The Delian bond

> *At one position the two pieces weigh the same. There the tetrahedron's piece holds exactly twice the volume of the octet's tetrahedron, and the octahedron's piece exactly half the volume of the octet's octahedron. Doubling a volume means multiplying lengths by the cube root of 2: the Delian problem of the ancient Greeks [11], which cannot be solved with ruler and compass [10]. Here it happens in both directions at once.*

> **Theorem 3 (the Delian bond).** $T(t)$ and $O(t)$ have equal volume at exactly one position, $t_D = 1 - (2/3)\cdot 2^{-1/3}$. There **Samson**, the cut cube $T(t_D)$, and **Delilah**, the octahedron $O(t_D)$, both have volume $\sqrt{2}/6 = 2V_T = V_O/2$: Samson twice the octet's tetrahedron and Delilah half the octet's octahedron. Delilah is the octet's octahedron shrunk by the factor $2^{-1/3}$, so her edge is $1/\sqrt[3]{2}$; Samson's edges are $1/\sqrt[3]{2}$ (his triangles, which are Delilah's faces) and $(1 - 2^{-1/3})/\sqrt{2}$.

> *Why: as the cut slides beyond the octet, one piece only grows and the other only shrinks, so they can be equal only once. Setting the two volume formulas equal leaves one number, the cube root of one half.*

Proof. For $t \le 1/3$, vol $T(t) \le V_T$ while vol $O(t) \ge 4V_T$, so they are never equal. For $t \ge 1/3$, as a grows (that is, as the cut moves back towards the octet), vol $T(t) = (3 - 2a^3)V_T$ decreases and vol $O(t) = 4a^3V_T$ increases, so equality holds for at most one a: $3 - 2a^3 = 4a^3$, $a^3 = 1/2$, $a = 2^{-1/3}$, $t_D = 1 - 2a/3$. Then both volumes are $2V_T = \sqrt{2}/6$. The edges are those of the Main theorem (b) at $a = 2^{-1/3}$. The study confirms it exactly in $\mathbb{Q}(\sqrt[3]{2})$: equal volumes, Delilah's edges, Samson's two edge lengths [5]. ∎




Figure 4. The Delian bond filling space: the eight Samsons (gold) and four Delilahs (blue) of one cube of the lattice, from two sides: the Samsons form a star round the Delilah hidden at the cube's centre.

6. The DICTONIC SOLIDS and BONDS

> *Paper #48 and this paper together give six new or newly framed solids that fill space in pairs. DICTO calls the six the DICTONIC SOLIDS and the three pairs the DICTONIC BONDS.*

Bond	Solids	From	Volumes
Golden	Icosaxes + Unborn Axes	icosahedra, Unborns and AXE, cut at the golden point (#48)	1 + 2√5/3 and 1 + √5/3
Keyed	Lotus Jewel Axes + Dodecaxes	the Lotus Jewel network, cut the same way (#48)	√5 and 2
Delian	Samson + Delilah	the octet, cut where the volumes are equal (this paper)	√2/6 each

Table 1. The three DICTONIC BONDS (names by DICTO). Golden and keyed volumes at icosahedron edge 1, Delian at octet edge 1.

Remark (the golden bond, polar). DICTO proposed looking at the solids turned inside out (polar reciprocation about their centres). The truncated octahedron T(x) of #48's Theorem 4 has its square faces at distance (1 + x)s and its hexagons at distance (1 + 2x)s/√3 from its centre (scale s), so its polar has corners at distance 1/((1 + x)s) on the three axes and at (±1, ±1, ±1)/((1 + 2x)s): a cube with a low pyramid on each face (14 corners, 24 triangles, the type of the tetrakis hexahedron). The pyramid's height over the face, measured in half cube edges, is (1 + 2x)/(1 + x) − 1 = x/(1 + x). For the Icosaxes (x = τ) it is τ/(1 + τ) = 1/φ, for the Unborn Axes (x = 1/τ) it is 1/φ², for Kelvin's cell (x = 1) it is 1/2; and for every pair x, 1/x the two heights add to x/(1 + x) + 1/(1 + x) = 1, which is the height of the pyramids that make a cube into Kepler's rhombic dodecahedron. Each height is measured in its own polar's half cube edge (the two polars have cubes of different sizes), so this is a statement about proportions: in proportion to its own cube, the polar of the golden bond divides the rhombic dodecahedron's pyramid between its two solids in the golden ratio. The polar of Delilah is a cube; the polar of Samson is an octahedron with low pyramids on four alternate faces; the

keyed solids are not convex and have no polar solid. (The polar solids are not claimed to fill space.)

7. Conclusion

> *The octet, the classical mixed filling of tetrahedra and octahedra, sits a third of the way along a slider that runs from Kepler's rhombic dodecahedron to the cube. Along it, one position makes the two pieces weigh the same: Samson and Delilah, where one piece doubles and the other halves.*

One flat cut per double cone turns the octet into a family of mixed space-fillers, every member convex, with the rhombic dodecahedron and the cube at its ends. Its special members are exact: two fillings with every edge equal, one where the two kinds share space equally, and the Delian bond, where the tetrahedron's piece doubles and the octahedron's halves. With paper #48 this completes DICTO's DICTONIC SOLIDS and BONDS: the golden bond of the icosahedron, its keyed form, and the Delian bond of the octet. Open: the same cut for the cube and octahedron lattices that #48's factor x reaches at its ends (octahedra with cuboctahedra), a single solid for each bond whose copies alone fill space, and printed sets of Samson and Delilah.

8. What is known, and what is new

> *The octet, its dual and the ways the cube, the octet and the rhombic dodecahedron are related are old. We did not find the slider itself, its equal-edged members or the Delian bond.*

Known. The octet (tetrahedral-octahedral honeycomb) [3]; the rhombic dodecahedron as a cube with six pyramids [2, 4] and, equally, as an octahedron with eight quarter-tetrahedra, and the cube as a tetrahedron with four corners; Knoll and Morgan [2] relate the cube packing, the octet and the rhombic dodecahedron packing by stretching, truncation, tetrahedra inscribed in cubes and duality, as separate steps. Weighted Voronoi (power, Laguerre) diagrams [6] give one-parameter families of convex fillings in general; the family here is one of them (the cut planes are the power bisectors of the tetrahedron and octahedron centres). The polar remark rests on the classical duality of the truncated octahedron (Kelvin's cell [14]) and the tetrakis hexahedron. Closest in spirit are the continuous families of tilings by tetrahedra and octahedra of Conway, Jiao and Torquato [7] and of Gabbrielli, Jiao and Torquato [8], obtained by retessellating the octet: there both kinds of tile stay regular and only their sizes and arrangement change, while here one piece stays regular and its partner becomes a cut rhombic dodecahedron or a cut cube.

Not found, in the sources listed in Appendix A: the one-parameter family through the octet from rhombic dodecahedra to cubes as such, its two equilateral members, the half-and-half member, the Delian bond (equal volumes, doubling and halving at once), and the golden division of the rhombic dodecahedron's pyramid by the polar golden bond. "Not found" is limited to those sources.

9. In the apps

Samson and Delilah are being added to Polyhedraverse among DICTO's pieces, with a Kaleidohedra world, DICTONIC, whose three bond buttons and one slider run through the golden, keyed and Delian bonds (https://dictospheres.com).

References

[1] DICTO, "The Golden Split: Icosahedra, Unborns and AXE Become Two Convex Space-Fillers", paper #48, DICTO DOI: 10.5281/zenodo.23596362 (2026).
[2] E. Knoll, S. Morgan, "Finding the Dual of the Tetrahedral-Octahedral Space Filler", Bridges: Mathematical Connections in Art, Music, and Science (2003) 205–212.
[3] "Tetrahedral-octahedral honeycomb", Wikipedia (viewed 23 October 2026).
[4] "Rhombic dodecahedron", Wikipedia (viewed 23 October 2026).
[5] DICTO, Shadow forms (study 2026-10-23-shadow-forms, commit 099b211): octet/octet_exact.py → octet-exact.txt (the family at six rational t) and octet/octet_special_exact.py → octet-special-exact.txt (the special members, exact in their number fields).
[6] F. Aurenhammer, "Power diagrams: properties, algorithms and applications", SIAM Journal on Computing 16 (1987) 78–96.
[7] J. H. Conway, Y. Jiao, S. Torquato, "New family of tilings of three-dimensional Euclidean space by tetrahedra and octahedra", Proceedings of the National Academy of Sciences USA 108 (2011) 11009–11012.
[8] R. Gabbrielli, Y. Jiao, S. Torquato, "Families of tessellations of space by elementary polyhedra via retessellations of face-centered-cubic and related tilings", Physical Review E 86 (2012) 041141.
[9] R. B. Fuller, "Synergetic building construction" (the octet truss), US Patent 2,986,241 (1961).
[10] P. L. Wantzel, "Recherches sur les moyens de reconnaître si un problème de géométrie peut se résoudre avec la règle et le compas", Journal de Mathématiques Pures et Appliquées 2 (1837) 366–372.
[11] T. L. Heath, A History of Greek Mathematics, vol. I, Clarendon Press, Oxford (1921), on the duplication of the cube.
[12] DICTO, "Aristotle's Pyramid", paper #44, DICTO DOI: 10.5281/zenodo.23529497 (2026).
[13] J. Kepler, Strena seu de nive sexangula (1611).
[14] W. Thomson (Lord Kelvin), "On the division of space with minimum partitional area", Philosophical Magazine 24 (1887) 503–514.

Appendix A. Sources checked for prior work

DICTO's reference library (kaleidohedra-references): nothing on this family; read for this check: Gabbrielli, Jiao and Torquato 2012 [8] (families of tilings by regular tetrahedra and octahedra, both kinds regular throughout), which builds on Conway, Jiao and Torquato 2011 [7]. Read in full: Knoll and Morgan 2003 [2]. Web, 23 October 2026: searches for cubes with four alternate corners truncated together with octahedra; rhombic dodecahedra truncated at their three-fold corners together with tetrahedra; power (Laguerre) diagrams of the tetrahedral and octahedral holes of the face-centred cubic lattice; tetrahedra and octahedra of equal volume, cube root of 2. Found only the classical octet and its relatives (Wikipedia: tetrahedral-octahedral, quarter cubic and tetragonal disphenoid honeycombs; MathWorld, space-filling polyhedra; Goldberg's catalogue as cited there). Not checked: Wolfram PolyhedronData, Stella / Great Stella libraries.

Appendix B. Data (octet edge 1, $V_T = \sqrt{2}/12$)

Position	t	Tetrahedron piece	Octahedron piece
rhombic dodecahedra	0	—	rhombic dodecahedron, $6V_T$
equilateral (a)	$(2\sqrt{6} - 3)/15 \approx 0.1266$	tetrahedron, edge $(2\sqrt{6} - 3)/5$	cut rhombic dodecahedron, same edge
the octet	1/3	tetrahedron, edge 1, V_T	octahedron, edge 1, $4V_T$
half and half	$1 - \sqrt[3]{6}/3 \approx 0.3943$	cut cube, $3V_T/2$	octahedron, $a = \sqrt[3]{6}/2$, $3V_T$
Delian bond	$1 - (2/3)\cdot 2^{-1/3} \approx 0.4709$	**Samson**, $2V_T$	**Delilah**, edge $2^{-1/3}$, $2V_T$
equilateral (b)	$5/3 - (2/3)\sqrt{2} \approx 0.7239$	cut cube, edge $\sqrt{2} - 1$	octahedron, edge $\sqrt{2} - 1$
cubes	1	cube, edge $1/\sqrt{2}$, $3V_T$	—