The DICTO Ogdoecontahedron



The Spheres of Dharma turned inside out by polar reciprocation: the dual of the Hecatodihedron (80 faces: 20 regular hexagons and 60 kites) as the second station of a reversed exact φ³ cycle, one kind of piece per layer, the great stellated dodecahedron locked in the outer gap, a dual pair in the inner gap, the cycle repeating inward to its core, the original and reversed cycles sharing one space, and the checks that keep each step exact

DICTO (artist, Osaka, Japan) · contact@dictospheres.com · https://dictospheres.com
Draft 2 · 16 October 2026 · DICTO DOI: on publication · Licence CC BY 4.0 · Data, scripts, exact checks and renders: https://github.com/DICTOR-Master/kaleidohedra (studies/2026-10-16-polar-dharma; shared checks in studies/_lib).

The Spheres of Dharma, the Hecatodihedron, the idea of turning the cycle inside out by polar reciprocation, the call for an even nest, the stars and the Hecatodihedron tried in the gaps, the dual pair and its reversed order, the golden rectangles, the endless core, the two cycles together as a toggle, the demand for stricter checks and the name Ogdoecontahedron are DICTO's. Computation, exact arithmetic and renders were done with Claude (Anthropic) as a research assistant.

Abstract. The Spheres of Dharma nest seven icosahedral solids exactly, from an icosahedron out to the icosahedron φ³ larger. DICTO asked what the cycle looks like turned inside out. Replacing every station by its polar reciprocal reverses the nesting and gives a new exact cycle: dodecahedron ⊂ icosidodecahedron ⊂ **Ogdoecontahedron** ⊂ pentakis dodecahedron ⊂ icosahedron ⊂ rhombic triacontahedron ⊂ dodecahedron φ³ larger (volume ratio φ⁹). The Ogdoecontahedron is the dual of the DICTO Hecatodihedron: 80 faces (20 regular hexagons, 60 kites), 102 corners, 180 edges, only two edge lengths, and 90 of its corners on one sphere, the inside-out image of the Hecatodihedron's 90 faces on one sphere. Alone it is not a patch of an icosahedral quasicrystal; packed in five periodic packings round a five-fold axis the copies that must be removed along the axis, together with the central copy, are exactly the canonical Fibonacci chain for |h| < 12.81 (finding #30, summarised in Section 2). Cutting each layer by cones over the faces of its inner solid gives **one kind of piece per layer** (12, 12, 20, 60, 20, 30), every volume exact in ℚ(√5) with remainder 0. The cycle inherits two loose ends. Tightening them gives an evenly stepped nest, still one kind per layer and exact. Left loose, the large outer gap holds the great stellated dodecahedron at exactly one size: its 20 points on the outer dodecahedron's corners, its 12 valleys on the triacontahedron's tips, cutting the gap into 60 tetrahedra and 60 pentagonal pyramids. The inner gap admits no star and no locked dual pair, but the truncated icosidodecahedron pair fits in reverse order (as polarity demands), scaled by 6/φ² and 2/15 with play √5·φ³/9. The golden rectangles that frame the whole are the outer dodecahedron's own small stellation, and the cycle repeats inward without end. The original and reversed cycles can share one space: a sphere of polarity exists that makes them share the icosahedron and dodecahedron, and another that makes them share the icosidodecahedron and triacontahedron, but no sphere does both. We also record the checks this study forced on us, because three of our first tables were wrong.

1. The Spheres of Dharma turned inside out

The Spheres of Dharma (DICTO DOI: 10.5281/zenodo.23310833) is an exact cycle of nested icosahedral solids: icosahedron ⊂ icosidodecahedron ⊂ dodecahedron ⊂ truncated icosahedron ⊂ Hecatodihedron ⊂ rhombic triacontahedron ⊂ icosahedron φ^3 larger, each layer filled by a few repeated pieces. Its seventh stage, closed as a convex solid, is the **DICTO Truncated-Icosahedral Hecatodihedron** (DICTO DOI: 10.5281/zenodo.23360858): the truncated icosahedron with a Shield on each hexagon, 102 faces (12 pentagons, 30 rhombi, 60 triangles), 80 corners.

DICTO's question was what happens to the cycle turned inside out. **Polar reciprocation** in a sphere about the common centre sends every plane to a point and every point to a plane; a solid goes to its dual, and inclusion reverses: if P ⊂ Q then Q* ⊂ P* (a classical fact of convex geometry; see e.g. Firsching 2015). So the reversed cycle is again a nest, read from the other end:

$$\mathbf{D \subset ID \subset Ogdoecontahedron \subset PD \subset I \subset RT \subset D \cdot \varphi^3}$$

(dodecahedron, icosidodecahedron, the Hecatodihedron's dual, pentakis dodecahedron, icosahedron, rhombic triacontahedron, dodecahedron). We computed every pole exactly in $\mathbb{Q}(\sqrt{5})$ from the snapped original corners. The outer dodecahedron is exactly φ^3 times the inner one, so the volume closes at $38 + 17\sqrt{5} = \varphi^9$. Duality swaps solids in pairs, so the icosidodecahedron and the triacontahedron, and the icosahedron and the dodecahedron, appear in both cycles in swapped places; the truncated icosahedron becomes the pentakis dodecahedron, and the Hecatodihedron becomes something new.




Figure 1. The reversed cycle, faithful to the original: (a) closed, the outer dodecahedron; (b) cut in half: under reciprocation the large layers become small and the small large, so the outer layer is far the largest; (c) pulled apart by layer.

2. The DICTO Ogdoecontahedron

The second station is the polar of the Hecatodihedron. DICTO named it for its faces, as he named the Hecatodihedron for its 102: the **Ogdoecontahedron**, from Greek ὀγδοήκοντα (eighty) and ἕδρα (face).

- **80 faces, 102 corners, 180 edges** (102 − 180 + 80 = 2). Each face is the dual of a Hecatodihedron corner. The 20 Shield apexes, where six faces meet, become **20 regular hexagons**; the 60 corners of the truncated icosahedron, where four faces meet, become **60 kites**.

- **Two edge lengths.** In the chain's scale (reciprocation in the unit sphere), the hexagon side squared is 2/9 − 2√5/27 (side 0.237881), and each kite has two sides of that length, next to the hexagons, and two of squared length 86/3249 + 2√5/361 (side 0.197124). The hexagons are regular (all angles 120°).
- **One sphere, inside out.** 90 corners lie exactly on the sphere of radius 2/3; the other 12 sit just outside, at squared radius (170 + 2√5)/361. The Hecatodihedron has 90 faces touching one sphere and its 12 pentagons a little inside it: reciprocation trades a tangent sphere of faces for a sphere of corners.
- **Built from the dual side.** The Hecatodihedron is the truncated icosahedron with a pyramid-like Shield added on each hexagon; its dual is the truncated icosahedron's dual, the pentakis dodecahedron, with its 20 six-fold corners **cut off flat** into regular hexagons. Adding a cap on one side is cutting a corner on the other.
- **Its place in the nest.** Its 20 hexagons lie in the planes of the icosidodecahedron's 20 triangles, each hexagon holding a triangle; its 60 kites lie in the faces of the pentakis dodecahedron around it. Volume (−1000 + 1240√5)/1539.
- **Its face arrangement is known; the solid is not.** Combinatorially (which faces meet which, ignoring shape), the Ogdoecontahedron is the pentakis dodecahedron with its 20 six-fold corners cut down to the edge midpoints; in the operator notation of the Antiprism software (Rossiter) it is the dual of L0 applied to the dodecahedron, so the Hecatodihedron has the arrangement of L0(D). We checked this by an exact isomorphism search on the face maps, with controls. What we did not find, in McCooey's catalogue of 553 solids scanned in full nor elsewhere, is this metric solid: regular hexagons with kites of 117.85°, 68.62°, 117.85° and 55.69°. The nearest catalogued relative is the 6-truncated pentakis dodecahedron (hexagons and pentagons), whose truncation, carried to the edge midpoints, gives this arrangement.




Figure 2. The DICTO Ogdoecontahedron, hexagons wisteria, kites cyan: (a) oblique; (b) along a three-fold axis, a regular hexagon in the centre; (c) along a five-fold axis, one of the 12 outer corners where five kites meet.

Periodic and aperiodic: the solid alone, and in packings (#30)

The solid alone. The regular hexagon is the tile of the periodic honeycomb; icosahedral symmetry is the symmetry of quasicrystals, which no periodic structure can have. In the single solid the two only sit side by side. Its kites are not Penrose kites (angles 117.8°, 68.6°, 117.8°, 55.7°), and its golden coordinates link it to the six-dimensional lattice of icosahedral quasicrystals no more than any golden solid. Our bridge study (studies/2026-10-16-ogdoecontahedron-bridge) found that it is neither a patch nor a cluster of the canonical icosahedral quasicrystal. Its corners reach the six-dimensional window only after nine golden inflations, when the solid spans thousands of tiling edges. None of its edges lies on a

symmetry axis, whereas every edge of that tiling does, and no face angle can be built from golden or Penrose rhombi. Its one tie is classical: its 30 two-fold corners are the icosidodecahedron's, the 30-atom shell of the Tsai and Mackay clusters. The study tested the canonical primitive tiling and three standard cluster shells only.

In packings: a Fibonacci spine (finding #30; DICTO DOI: 10.5281/zenodo.23396361). The link appears when copies of the solid are packed. The Ogdoecontahedron is centrally symmetric, and its 20 hexagon directions contain exactly five inscribed cubes. Joining copies face to face on the eight hexagons of one cube gives a periodic, body-centred cubic packing: five periodic packings, the five "voices". Placed in five 72° wedges round a five-fold axis, they form the icosahedral irrational twin (neighbouring voices related by 44.4775° about a shared three-fold axis). Copies that clash are removed in whole five-fold rings. Checked exactly in $\mathbb{Q}(\sqrt{5})$, the result is overlap-free to radius 10: 2756 sites, 385 removed, 7465 close pairs, 0 overlaps.

- **Golden heights (exact).** Every copy sits at a height $(m + n\varphi)u$ along the axis, with $u^2 = 8/81 - 8\sqrt{5}/405$.
- **The spine is the canonical Fibonacci chain (exact).** Apart from one ring, the removed copies nearest the axis lie in one plane, on a rank-2 lattice in which the axis has the irrational slope $-\varphi$. Together with the central copy, which is kept, they are exactly the canonical singular Fibonacci cut-and-project chain, with window $[0, W)$, $W^2 = 64/81 + 128\sqrt{5}/405$ ($W = 1.223451$), for $|h| < 12.81$. The gaps are $S = 1.22345$ and $L = 1.97958$, with $L/S = \varphi$.
- **The window has a mechanism (exact).** A ring of five copies overlaps itself exactly when it is closer to the axis than t_c, $t_c^2 = 8/9 + 8\sqrt{5}/45$ ($t_c = 1.134201$). Since $t_c < W$ and no plane lattice point lies between them for $|h| < 12.81$, the two windows select the same points there.
- **Beyond 12.81.** At $h = 12.81$ the core copy overlaps one other copy, which overlaps two more (exact). On finite patches (integer program, float clash test), keeping the core copy is cheaper by one ring, there and at the next three such points (-22.71, 34.77, -44.66). Minimal removal therefore thins the spine to a three-gap chain of the narrower window $[0, t_c)$, still aperiodic but no longer Fibonacci. Staying pure Fibonacci costs at least one extra ring at each such point (555 copies removed instead of 385 at radius 10).

So aperiodic order appears where periodic packings of this solid meet. The chain's lattice is a plane section of one packing, its internal coordinate is the distance from the axis, and its window is fixed by the ring clash at t_c. No aperiodic structure is put in: only five periodic packings of one solid meeting round an irrational (five-fold) axis, removal in whole five-fold rings, and the rule that hard copies may not overlap. Interfaces between periodic grains as one-dimensional quasicrystals, model sets near irrational lines and hard-core windows are known (Sutton 1988; Ranganathan, Srivastava & Lord 2006; Baake & Grimm 2013). The selection along a five-fold junction line, its self-clash window and the departure from it were not found, and are presented as a new exact instance of known ideas. The proofs, the fork at 12.81 and the seams are in the #30 paper.




Figure 3. From the #30 paper. (a) The five-fold chorus seen down the axis (small early build): the five voices in five colours, removed copies as ghosts. (b) The core plane as a cut-and-project picture (exact points, drawn in float): plane lattice points by height along the axis and distance t from it. Circled: removed in the radius-10 build. Pale band: the canonical window [0, W). Darker band: the ring-clash window [0, t_c). Star: the central copy. Diamond: the fork at height 12.81.

3. One kind of piece per layer

We cut each layer A ⊂ B by the cones from the centre over the faces of the inner solid A: each piece is cone(F) ∩ B beyond F, convex by construction (the Kion construction of Theaetetus, DICTO DOI: 10.5281/zenodo.23378070). Faces of A that already lie on B's surface give no piece. Every corner was solved exactly as the meeting point of three exact planes, and every volume summed exactly against its layer.

Layer	Pieces	Each piece	Exact volume
D → ID	12	a column on each dodecahedron face (two regular pentagons, 5 pentagonal and 5 triangular sides)	(−144185 + 64487√5)/243
ID → Ogdoecontahedron	12	a cap on each icosidodecahedron pentagon (its triangles lie in the hexagons)	(−155 + 101√5)/4617
Ogdoecontahedron → PD	20	a low pyramid on each regular hexagon (the kites lie in PD's faces)	(−22 + 10√5)/81
PD → I	60	an irregular wedge, 5 corners and 6 triangles	(−6560 + 2936√5)/513
I → RT	20	Kepler's pyramid on each icosahedron face	(−796 + 356√5)/3
RT → D·φ³	30	a roof on each rhombus, its ridge a dodecahedron edge	200 − 1336√5/15

In every layer, the sum of the pieces minus the layer volume is exactly 0.




Figure 4. One piece of each layer, seen from the side with the outward direction up, each drawn at its own scale; layer colours as in the cut views. (a)–(f) the faithful nest, inside to outside (O = Ogdoecontahedron): the column on a dodecahedron face, the cap on an icosidodecahedron pentagon, the low pyramid on a regular hexagon, the irregular wedge, Kepler's pyramid, and the roof on a rhombus. (g), (h) the two end pieces that change in the tight nest (Section 4): the near-antiprism column and the wedge on a rhombus.

4. Two loose ends, and the even nest

The faithful copy keeps the original closure (outer dodecahedron = inner × φ^3), and that leaves two stations floating: the triacontahedron could grow ×1.894 before touching the outer dodecahedron, and the inner dodecahedron ×1.232 before touching the icosidodecahedron. The four middle stations touch. DICTO asked for a more even nest. Tightening both ends, the inner dodecahedron grows by exactly $11/18 + 5\sqrt{5}/18$ until its corners touch the icosidodecahedron's triangles, and the outer one shrinks by exactly $5 - 2\sqrt{5}$ until the triacontahedron's twelve tips touch its face centres. The inradii then step evenly (0.495, 0.567, 0.623, 0.653, 0.714, 0.764, 0.898), the outer layer falls from 25.2 to 1.9, and every layer is still one kind of piece, exact with remainder 0: the inner layer becomes 12 near-antiprism columns, $(-35/729 + 71\sqrt{5}/2187)$ each, and the outer 30 wedges on the rhombi, $(-7352/3 + 1096\sqrt{5})$ each. The price is the golden closure: the tight cycle repeats at linear ratio 1.8146 (squared $49815/2 - 22275\sqrt{5}/2$), not a power of φ. We keep both versions.




Figure 5. Faithful and tight, cut in half: (a) the faithful nest, its large outer gap; (b) the tight nest, both loose ends closed.

5. The great stellated dodecahedron in the outer gap

DICTO asked whether a Kepler–Poinsot solid could sit in the faithful nest's large outer gap. We treated each star as a filled body (points of nonzero winding number; the densities at the centre come out 3, 3, 7, 7 as they should) and found the range of sizes that keeps the triacontahedron inside the star and the star inside the outer dodecahedron. The great icosahedron does not fit. The great and small stellated dodecahedra float between two sizes. The **great stellated dodecahedron fits at exactly one size**: its 20 points are the outer dodecahedron's 20 corners, and its 12 valleys, where five spikes meet, are the chain's icosahedron corners, which are also the triacontahedron's five-fold tips. We checked this exactly: each of the 12 icosahedron corners lies on five of the star's face planes.

The star is the chain's icosahedron with a spike on each face (Kepler; our In-betweens paper, DICTO DOI: 10.5281/zenodo.23345502, #27), and its volume is exactly 3 icosahedra, as follows from the standard formulas. Its surface is 60 triangles, each seen once from the centre, and the cones over them cut the gap into two exact layers: inside the star, around the triacontahedron, **60 congruent tetrahedra** (point, triacontahedron corner and two valleys), $(188/3 - 28\sqrt{5})$ each; outside it, up to the dodecahedron, **60 congruent pentagonal pyramids**, $(112/3 - 248\sqrt{5}/15)$ each. Both sum to their layers with remainder 0.

The fit is rhythmic rather than lucky. The squared ratio of the star's point radius to its valley radius is $3\varphi^3/\sqrt{5}$: apart from the fixed $\sqrt{3}$ and $\sqrt{5}$ that convert between three-fold and five-fold radii, it is φ^3, the step by which the whole cycle closes.




Figure 6. The great stellated dodecahedron in the outer gap: (a) the star alone, 60 tetrahedra (orange) around the triacontahedron; (b) cut in half; (c) closed by 60 pentagonal pyramids (indigo) to the outer dodecahedron.

6. The inner gap and a dual pair in reverse order

In the inner gap, between the dodecahedron and the icosidodecahedron, none of the four stars fits. The Hecatodihedron itself fits but floats: its size can vary ×1.232, touching 20 points at either end and never both. DICTO then asked for a dual pair instead. A pair X, Y locks when three contacts hold at once (dodecahedron on X, X on Y, Y on the icosidodecahedron). No icosahedral Archimedean–Catalan pair locks. The truncated icosidodecahedron does *not* fit inside its dual; as DICTO pointed out, in the reversed cycle the pair must go in reverse order too, and so it does: the **disdyakis triacontahedron inside the truncated icosidodecahedron**. Exactly: the disdyakis triacontahedron (from the dual of the standard truncated icosidodecahedron) is scaled by $9 - 3\sqrt{5} = 6/\varphi^2$, the truncated icosidodecahedron by 2/15, and the play between them is $5/9 + 2\sqrt{5}/9 = \sqrt{5}\cdot\varphi^3/9 \approx 1.0525$. The truncated icosidodecahedron's 12 decagons lie exactly in the icosidodecahedron's pentagons. The layers cut into 12 pieces of one kind, 120 of one kind and 50 caps of two kinds, all exact with remainder 0. The nearest pair to a lock is the rhombicosidodecahedron inside the deltoidal hexecontahedron, play exactly $62/123 + 137\sqrt{5}/615 \approx 1.00218$: short of locking by 0.2%.




Figure 7. The inner gap: (a) the truncated icosidodecahedron pair up to the icosidodecahedron, the decagons flush in its pentagons; (b) cut in half: dodecahedron, disdyakis triacontahedron layer (cyan), truncated icosidodecahedron layer (amber), caps (coral); (c) the rhombicosidodecahedron pair, cut.

7. The golden frame

DICTO asked whether golden rectangles would frame the whole. The 15 golden rectangles of an icosahedron (five perpendicular triples) clear the whole nest when the icosahedron is the

next turn of the cycle, the chain's icosahedron × φ^3 (the smallest icosahedron around the dodecahedron is not enough). Then each of the outer dodecahedron's 30 edges lies along one of the rectangles' 30 long edges, centred, and the long edge is φ^3 times as long (corner radius^2 $10 + 2\sqrt{5}$, midpoint radius^2 4). This is classical: the 30 long edges are the edges of the outer dodecahedron's **small stellated dodecahedron**, made by extending its edges (Kepler 1619), a pentagram chord being $1 + 2\varphi = \varphi^3$ times the pentagon side. The frame that holds the cycle is its outer dodecahedron's own first stellation.




Figure 8. The nest (with the star layers) inside the 15 golden rectangles of the next turn's icosahedron.

8. The core: the cycle repeats inward

What fills the inner dodecahedron? The cycle itself. Its outer dodecahedron is exactly φ^3 times its inner one, so the whole nest scaled by $1/\varphi^3$ fills the inner dodecahedron with no gap. Inside that sits the next turn, and so on to the centre. In the faithful version every turn brings back the large gap, so the great stellated dodecahedron locks at the heart of every turn: its points on that turn's dodecahedron corners and its valleys holding the next triacontahedron. The golden frame of Section 7 is the next turn outward, and the core is the next turn inward.




Figure 9. The infinite core, cut in half, in true proportion: (a) four turns, each a copy of the whole nest at $1/\varphi^3$ of the one around it, the great stellated dodecahedron (orange spikes) at the heart of every turn; (b) the centre, one turn in. Each deeper turn is its own render set in at exactly $1/\varphi^3$, so the picture is seamless and stays sharp.

9. Both cycles at once: the toggle

DICTO asked whether the original cycle and its reversed image could share one space. They can, and the sphere of reciprocation is the one free choice. Polarity in a sphere of radius ρ sends radius r to ρ^2/r, so in log-radius it is a reflection about $\log \rho$; the cycle's φ^3 repeat is a translation. Together they give mirror spheres every $\varphi^{3/2}$ in radius: the union of the two cycles reads the same outward and inward, a palindrome in scale, and the polarity swaps the two cycles.

- **Ends setting**, $\rho^2 = \varphi^2/4 = (3 + \sqrt{5})/8$, mirror sphere radius $\varphi/2$: the polarity sends the original icosahedron to the original dodecahedron, so the two cycles share I and D. The reversed turn nests almost entirely inside the original icosahedron.
- **Middle setting**, $\rho^2 = 3\varphi^2/4$, mirror sphere radius $\sqrt{3}\cdot\varphi/2$: they share the icosidodecahedron and the triacontahedron instead. The reversed cycle is then exactly 3 times its ends-setting size.
- **Never both.** Moving from one setting to the other scales ρ^2 by 3, while the cycle only repeats under powers of φ^3; 3 is not a power of φ, so no sphere gives both. Nor can the loose ends be bent to allow it: the original icosahedron can be scaled only within $[1/1.232241, 1.894427]$ before the nest breaks, and sharing both pairs would need a scale of $3/\varphi^3 \approx 0.708$ or 3.

Cut against each other, the two cycles form one solid of convex pieces, with every piece and layer summing exactly; that interlock, and the shapes of its pieces, are reported separately (studies/2026-10-16-interlock).




Figure 10. Both cycles in one space, cut in half, see-through: the original cycle warm, the reversed cycle cool, shared stations in dark ink. (a) Ends setting, sharing the icosahedron and dodecahedron; (b) middle setting, sharing the icosidodecahedron and triacontahedron.




Figure 11. Both settings on one radial scale (log radius, three turns of each cycle). Each station is drawn as the range from its face distance to its corner radius, with a dot at their geometric mean; grey lines join each station to its polar image through the toggle sphere; dashed lines are the mirror spheres, every $\varphi^{3/2}$; black marks the stations the two cycles share. TI = truncated icosahedron, H = Hecatodihedron, O = Ogdoecontahedron, PD = pentakis dodecahedron.

10. Method: the checks this study forced on us

Three of our first tables in this study were wrong, and each was caught before publication. Standard coordinate lists for the Archimedean solids come in two mirror golden frames, and most of ours were in the wrong one, so they did not share the chain's axes. A slack formula

was inverted. A float face count reported 103 kinds of piece in a layer of 120 congruent pieces. DICTO asked for rules that would prevent this, and every study script now calls a shared module (studies/_lib/geomcheck.py) that stops with an error instead of printing a wrong table:

1. **Frame:** every generated solid must be invariant under the chain's own five-fold and three-fold turns.
2. **Both orders:** every pair is tested both ways; polar reversal swaps the order of a dual pair.
3. **Known case first:** every formula reproduces a known answer (the dodecahedron's slack 1.232 in the icosidodecahedron; the triacontahedron locked in the star; star densities 3, 3, 7, 7) before it is used.
4. **Kinds by symmetry:** pieces are the same kind when the icosahedral group maps one onto the other, after reducing float hulls to their true corners.
5. **Exact gate:** a float result is a lead; a claim is made when the exact $\mathbb{Q}(\sqrt{5})$ remainder is 0. Exact corners come from plane triples that reproduce the point.
6. **Labels:** every number is marked exact or float, faithful or adjusted, and originals are never overwritten.

The module's tests replay each of our slips and show the check catching it.

11. Relation to known work, and status

Known and credited: polar reciprocation and the reversal of inclusion (convex geometry; Firsching 2015); the largest copy of one polytope in another (Croft 1980; Firsching 2015 computes it exactly; our fits are the concentric, same-orientation case); nested icosahedral shells in clusters, viruses and fullerenes (Takakura et al. 2007, the Tsai cluster of the Spheres of Dharma; Twarock, Valiunas and Zappa 2015; Dechant et al. 2014); the Kepler–Poinsot solids, the great stellated dodecahedron as an icosahedron with twenty spikes and the small stellated dodecahedron as the dodecahedron's extended edges (Kepler 1619; Poinsot 1810); the 15 golden rectangles of the icosahedron. Hafner (2007) dissects the small and great stellated dodecahedra into the triacontahedron and hexecontahedron using 60 tetrahedra, each 1/6 of a prolate golden rhombohedron; our 60 tetrahedra between the triacontahedron and the star are different pieces: each has exactly φ^2 times the volume of his on the same triacontahedron edge (checked in $\mathbb{Q}(\sqrt{5})$).

Not found in our reference library and web searches (16 October 2026): the reversed Spheres of Dharma as an exact nest with one piece kind per layer; the Ogdoecontahedron and the Hecatodihedron (#28) as metric solids (their face arrangements are known, Section 2); the tight nest; the great stellated dodecahedron's lock in the reversed cycle, which follows directly from classical facts and is presented as an observation about this cycle, not a new theorem; the inner-gap dual pairs and their exact scales and play; the exact pieces and volumes; the two settings of the toggle and the proof that they exclude each other.

12. Reproducibility

In studies/2026-10-16-polar-dharma: `polar.py` (the reversed chain, exact poles, closure, touching), `polar_pieces.py` (the cone cut; kinds by symmetry orbit), `polar_exact.py` (exact corners and volumes; arguments `tight`, `tid`, `rid`), `polar_tight.py`, `polar_stars.py` and `polar_gsd_exact.py`, `polar_gsd_pieces.py`, `polar_heca_gap.py`, `polar_round.py`, `polar_dual_pair.py`,

`polar_pair.py`, `polar_golden_rect.py`, `hstar_render.py`, `polar_turns.py`, `polar_hafner_exact.py`, `toggle_render.py`; the face-arrangement check is `ocd_dl0d.py` in studies/2026-10-16-ocd-pyrochlore; outputs in `data/`, renders by `render_polar.html`, `render_hstar.html` and the `shoot_*.mjs` scripts. Checks: `python3 studies/_lib/test_geomcheck.py`.

References

1. J. Kepler, *Harmonices mundi libri V*, Linz, 1619.
2. L. Poinsot, Mémoire sur les polygones et les polyèdres, *J. École Polytechnique* 10 (1810) 16–48.
3. M. Firsching, Computing maximal copies of polyhedra contained in a polyhedron, *Experimental Mathematics* 24 (2015) 98–105, doi:10.1080/10586458.2014.956374.
4. H. T. Croft, On maximal regular polyhedra inscribed in a regular polyhedron, *Proc. London Math. Soc.* (3) 41 (1980) 279–296, doi:10.1112/plms/s3-41.2.279.
5. I. Hafner, Dissection of small stellated dodecahedron and great stellated dodecahedron to rhombic triacontahedron and hexecontahedron, *Visual Mathematics* 9 (2007) no. 2.
6. R. Twarock, M. Valiunas, E. Zappa, Orbits of crystallographic embedding of non-crystallographic groups and applications to virology, *Acta Cryst. A* 71 (2015) 569–582, doi:10.1107/S2053273315015326.
7. P.-P. Dechant, J. Wardman, T. Keef, R. Twarock, Viruses and fullerenes: symmetry as a common thread?, *Acta Cryst. A* 70 (2014) 162–167, doi:10.1107/S2053273313034220.
8. A. Rossiter, Antiprism, polyhedron modelling software, https://www.antiprism.com.
9. D. McCooey, Visual Polyhedra, https://dmccooey.com/polyhedra (scanned 15 October 2026).
10. H. Takakura, C. P. Gómez, A. Yamamoto, M. de Boissieu, A. P. Tsai, Atomic structure of the binary icosahedral Yb–Cd quasicrystal, *Nature Materials* 6 (2007) 58–63, doi:10.1038/nmat1799.
11. DICTO, Spheres of Dharma, DICTO DOI: 10.5281/zenodo.23310833.
12. DICTO, The DICTO Truncated-Icosahedral Hecatodihedron, DICTO DOI: 10.5281/zenodo.23360858.
13. DICTO, Capturing the In-betweens, DICTO DOI: 10.5281/zenodo.23345502.
14. DICTO, Theaetetus, DICTO DOI: 10.5281/zenodo.23378070.
15. A. P. Sutton, Irrational tilt grain boundaries as one-dimensional quasicrystals, *Acta Metall.* 36 (1988) 1291–1299, doi:10.1016/0001-6160(88)90281-7.
16. S. Ranganathan, A. K. Srivastava, E. A. Lord, Coincidence-site lattices as rational approximants to irrational twins, *J. Mater. Sci.* 41 (2006) 7696–7703, doi:10.1007/s10853-006-0966-y.
17. M. Baake, U. Grimm, *Aperiodic Order, Vol. 1: A Mathematical Invitation*, Cambridge University Press (2013), doi:10.1017/CBO9781139025256.
18. DICTO, Is the DICTO Ogdoecontahedron a real periodic–aperiodic bridge? Study, 15 October 2026: https://github.com/DICTOR-Master/kaleidohedra/tree/3fd407f/studies/2026-10-16-ogdoecontahedron-bridge.
19. DICTO, A Fibonacci Spine in a Five-fold Chorus of Ogdoecontahedron Packings, DICTO DOI: 10.5281/zenodo.23396361.
20. DICTO, Kaleidohedra (apps), DICTO DOI: 10.5281/zenodo.23173809.