The Chain of Karma

The icosahedron and the dodecahedron as endless shells of each other

DICTO (artist, Osaka, Japan) · contact@dictospheres.com · https://dictospheres.com
Version 1 · 12 October 2026 · Licence CC BY 4.0 · Data, scripts, renders and checks are included with this record and at https://github.com/DICTOR-Master/kaleidohedra.
The Lotus seed, the Chain of Karma, the FUJI tip + mirror rule and the names Lotus seed, AXE, FUJI and CLEO are DICTO's. Design study and checking with Claude (Anthropic).

Abstract. We show that the regular icosahedron and the regular dodecahedron can each be filled exactly as a shell around the other, and that the two shells alternate without end. A dodecahedron of edge 1 is an icosahedron of edge 1/φ inside a shell of 12 pentagonal pyramids, 30 AXE and 20 CLEO; an icosahedron of edge φ² (the Lotus seed) is that dodecahedron inside a shell of 12 J11 (each an icosahedron with one pyramid cap removed), 30 AXE and 20 FUJI. Thus the nesting I(1/φ) ⊂ D(1) ⊂ I(φ²) ⊂ D(φ³) ⊂ … grows by φ³ per cycle, and each shell has one piece for every vertex, edge and face of the dual pair (12, 30, 20). The pieces are exact, at edge 1: AXE is a tetrahedron with five edges 1 and one 1/φ (two equilateral faces, two golden triangles 1, 1, 1/φ); CLEO is a triangular pyramid on an equilateral base of side 1/φ with side edges 1; FUJI is a pyramid of height 1/√3 on a flat-ended (truncated) triangle with sides 1 and 1/φ alternating and all angles 120°, with side edges 1, three equilateral and three golden faces, and a cube corner at its apex; the pentagonal pyramid has all edges 1. Every dihedral angle of the kit is, modulo 180°, an integer combination of two angles, α = arccos(√5/3) = 41.81° and δ = arctan(3 − √5) = 37.38°: the icosahedron's own is 180° − α = 138.19°, and FUJI's faces meet its base at α on the long sides and δ on the short. Because every icosahedron inside is again whole, each can be replaced by its own Lotus seed: the Chain of Karma, built and checked to 20,497 pieces. The gap pieces cannot be rebuilt from copies of themselves at one size (Dehn invariant); instead we give a growth rule for every pyramid, and so for AXE, CLEO, FUJI and the pentagonal pyramid: φ·P = P + a copy of size 1/φ turned over + a side ring with three times P's volume and Dehn invariant zero. By Sydler's theorem the ring is always equivalent to a cube, so the kit regenerates itself up to a cube-equivalent remainder; whether the ring can be cut into kit pieces remains open. A second exact cycle, through the icosidodecahedron, the fullerene cage and the rhombic triacontahedron, shares the dodecahedron with the Lotus seed and is described separately (Spheres of Dharma). The nesting of the dual pair at golden ratios goes back to Pacioli and Kepler and appears at every φ-power in Gray (2002); a different dodecahedral growth, with whole dodecahedra inside, is due to Al Raisi, Koca, Koca and Koc (2026). The exact shells and their endless alternation are new.

1. Two shells

Throughout, φ = (1 + √5)/2, edges are given relative to the dodecahedron D(1) of edge 1, and I(a), D(a) denote the regular icosahedron and dodecahedron of edge a. Two exact decompositions carry the whole construction:

$$D(1) = I(1/\varphi) + 12\ J2 + 30\ AXE + 20\ CLEO, \quad I(\varphi^2) = D(1) + 12\ J11 + 30\ AXE + 20\ FUJI.$$

In the first, the 12 pentagonal pyramids J2 (all edges 1) stand on the dodecahedron's faces with their points inward, and each point lands exactly on a corner of I(1/φ): the dodecahedron's inradius 1.113516 minus the pyramid's height 0.525731 is 0.587785, the circumradius of I(1/φ). In the second, each of the 12 outer icosahedra I(1) is cut by a face plane of D(1) into a J2 cap (inside D) and a J11 (outside). Volumes: 0.515028 + 12 × 0.301503 + 30 × 0.083333 + 20 × 0.051503 = 7.663119 = D(1), and 7.663119 + 12 × 1.880192 + 30 × 0.083333 + 20 × 0.321175 = 39.148928 = I(φ²) = φ⁶ I(1). Together they give the Lotus seed (DICTO Icosa-13, 113 pieces): I(φ²) = 12 I(1) + I(1/φ) + 60 AXE + 20 FUJI + 20 CLEO, whose 13 icosahedra are arranged as in Gray (2002). Each shell has one piece for every vertex, edge and face of the dual pair: 12, 30 and 20.

2. The Chain of Karma

Since the outer form of the Lotus seed is exactly I(φ²), it is a golden inflation rule. Replacing every icosahedron by a Lotus seed of its own size and turn, and keeping AXE, FUJI and CLEO whole, fills I(φ⁴), I(φ⁶), … exactly, upward without end, and in the same way downward. After n steps there are 13^n icosahedra and $100(13^n - 1)/12$ gap pieces, AXE, FUJI and CLEO in the ratio 3 : 1 : 1. Steps 1 to 3 were built and checked: 113, 1,569 and 20,497 pieces, volume equal to φ^{6n} times I(1) to within 3×10^{-9}. The icosahedra's share of the volume falls by $(12 + \varphi^{-3})/\varphi^6 = 0.6823$ each step (68.19%, 46.50%, 31.71%). Keeping only the icosahedra gives a fractal of similarity dimension D with $12\,\varphi^{-2D} + \varphi^{-3D} = 1$, D ≈ 2.6063, slightly above the Sierpinski icosahedron's log 12 / log φ² ≈ 2.5819 (12 corner copies only).




Figure 1. The Lotus seed pulled apart.




Figure 2. The Lotus seed closed: its surface carries the fullerene (C60) pattern, with the 20 flat-ended triangles (sides 1 and 1/φ alternating, all angles 120°) in green.




Figure 3. The Chain of Karma, two levels, pulled apart level by level.




Figure 4. Three levels: the Chain continues without end.

3. The gap pieces and the Dehn invariant

Every dihedral angle of the kit is, modulo 180°, an integer combination of α = arccos(√5/3) = 41.810° and δ = arctan(3 − √5) = 37.377°. This makes the Dehn invariant computable exactly (coordinates in Q(√5), integer relations by PSLQ). At a single size the gap pieces cannot be rebuilt from the kit: φ²·AXE, φ²·FUJI and φ²·CLEO pass the volume test (for example φ²·AXE has the volume of 9 AXE + 2 FUJI + 2 CLEO) but no whole-number combination of the edge-1 kit matches their Dehn invariants; the same program recovers the Lotus seed's own counts for φ²·I as a check. So the Chain of Karma is a hierarchy with gap pieces in every size step of φ², not a single-size tiling. With mixed sizes the screen rules nothing out; it is a necessary condition only. AXE and CLEO are, up to the factor 1/φ, the Mosseri–Sadoc tetrahedra t6 and t4 (Mosseri and Sadoc 1982; Kramer 1982, whose seven tiles add T2 = t2 ∪ t4). The six tetrahedra admit no stone inflation (Papadopolos and Ogievetsky 1999), and Danzer's ABCK tetrahedra all have Dehn invariant zero.




Figure 5. The kit at edge 1, every edge labelled: AXE, CLEO, FUJI, the pentagonal pyramid and J11, with their volumes. 60 AXE + 20 FUJI + 20 CLEO + 12 icosahedra of edge 1 + 1 of edge 1/φ fill the icosahedron of edge φ² exactly.

4. FUJI: tip, mirror, sides

DICTO's rule for a pyramid P (apex over a base). (1) *Tip:* cutting φ·P parallel to its base at P's height leaves P itself, so φ·P = P + a band. (2) *Mirror:* P(1/φ), turned upside down, sits in the

band with its base on the tip's base and its apex on the band's floor; the band's height is exactly P(1/φ)'s height. (3) *Sides:* the rest is a ring. Its volume is $(\varphi^3 - 1 - \varphi^{-3})\ P = 3P$, and its Dehn invariant $(\varphi - 1 - 1/\varphi)\ D(P) = 0$. Nothing here depends on the base, so the rule holds for every pyramid; AXE and CLEO are pyramids on any face, and FUJI and J2 on their bases (for the turned copy to fit, the base must contain its own point reflection scaled by 1/φ, as the regular pentagon and FUJI's hexagon do). By Sydler's theorem (1965) a region of Dehn invariant zero is scissors-congruent to a cube, so every piece of the kit regrows one φ-step larger from itself up to a cube-equivalent remainder. The sloping faces of the ring also tile with the kit's face shapes: each 60° trapezoid (top 1, bottom φ, legs 1/φ) is 2 triangles of side 1/φ, 2 of side $1/\varphi^2$ and one flat-ended triangle of size $1/\varphi^2$. Whether the ring itself can be cut into kit pieces is open. FUJI alone splits as a cube corner (the cube inside the dodecahedron; volume 1/6) plus three pieces each of CLEO's volume, and into 12 right-angled pieces whose faces slope at exactly 41.81° and 37.38°.




Figure 6. The pyramid growth rule, level by level: φ·P = P + a copy of size 1/φ turned over + a side ring with three times P's volume.

5. The dodecahedral Chain

DICTO's dodecahedral Chain starts from the first decomposition, D(1) = 12 J2 + I(1/φ) + 30 AXE + 20 CLEO. The icosahedron at the core grows by the Chain of Karma while the pyramid caps grow by the rule of section 4, and each cycle ends as a regular dodecahedron again: 63, 175 and 1,631 pieces for D(1), $D(\varphi^2)$ and $D(\varphi^4)$. Al Raisi, Koca, Koca and Koc (2026) grow the dodecahedron with whole dodecahedra inside; here it grows around icosahedra.




Figure 7. The dodecahedral Chain, its shell pulled apart.

6. A second cycle: the Spheres of Dharma

A second exact, endlessly repeating cycle passes through the icosidodecahedron, the fullerene cage and the rhombic triacontahedron: icosahedron φ^2 ⊃ triacontahedron ⊃ cage ⊃ dodecahedron ⊃ icosidodecahedron ⊃ icosahedron $1/\varphi$. It shares the dodecahedron of edge 1 and both icosahedra with the Lotus seed, and each half of either structure fills its side of that dodecahedron exactly, so the two can be exchanged half by half at every scale. It is described in its own paper (DICTO, Spheres of Dharma, #23).

7. The 600-cell, flattened

DICTO's route into four dimensions began with the DICTO Dodeca-13 (#13): a dodecahedron with twelve more on its faces, the 120-cell's first shell flattened, its 10.3° gaps being the flattening. The Lotus seed is the same for the 600-cell, the 120-cell's dual, as the icosahedron is the dodecahedron's. Seen down one vertex, the 600-cell (circumradius 1, edge $1/\varphi$) has shells of 1, 12 (icosahedron, radius 0.587785), 20 (dodecahedron), 12 (icosahedron), 30 (icosidodecahedron), …, alternating icosahedron and dodecahedron as the Chain's shells do. Two matches are exact (`cell600_match.py`): the Lotus seed's small icosahedron is the 600-cell's first shell (edge $1/\varphi$, circumradius 0.587785), and the centres of the Lotus seed's 13 icosahedra are the 600-cell's centre vertex and its 12 neighbours, scaled by φ^2 (radius 1.538842, spacing φ). The 600-cell is the icosahedral "12 around 1" packing that closes only in curved four-dimensional space; in flat space twelve around one leaves gaps (geometrical frustration, Sadoc and Mosseri). The Lotus seed reads as a flattened piece of the 600-cell, with AXE, FUJI and CLEO as exactly what flat space needs where curved space closes by itself, and the Chain of Karma repeats that flattening at every scale. The match is in projection; we do not claim the Lotus seed is a projection of the 600-cell.

The fullerene chain

DICTO (2026-10-11): the Lotus seed shows a fullerene architecture. Checked (`fullerene_surface.py`): its surface is 60 triangles of edge 1, grouped round the 12 corners as

pentagonal caps, and 20 flat-ended triangles (DICTO's term: triangles with their corners cut flat, sides 1 and 1/φ, the FUJI bases) in the places of the hexagons, so the Lotus seed has the face pattern of C60, the buckyball (Kroto et al. 1985), with flat-ended triangles for its hexagons: a truncated icosahedron whose cut corners are filled back by whole icosahedra. Chain step 2 keeps the 12 caps, has flat-ended triangles in two sizes (60 small, 20 of sides φ and φ^2) and 60 pairs of triangles folded over the edges: fullerenes inside fullerenes, every piece a triangle, whole or flat-ended. Counted as hexagons it has C180's 80, but it is not a Goldberg polyhedron (Goldberg 1937). The nesting recalls carbon onions, nested icosahedral fullerene shells (Ugarte 1992).

8. Periodic and aperiodic

The Chain of Karma is self-similar and not periodic: its symmetry is icosahedral and its scale steps are powers of φ. A periodic frame for icosahedra exists: an icosahedron with six edge roofs is a regular octahedron, so icosahedra with their roofs in the octahedra of the octet truss, and smaller icosahedra with corner and edge caps in its tetrahedra, fill space exactly (DICTO's icosahedral octet). Whether the Chain can grow inside that frame, joining the periodic and the self-similar orders, is open.

9. Self-similar, repeating, not periodic

Scaling is the growth step: the icosahedron and dodecahedron shells repeat by scaling, φ^3 per cycle, and the counts follow it exactly (13^n icosahedra after n steps of the Chain, gap pieces in the fixed proportion AXE : FUJI : CLEO = 3 : 1 : 1), as the growth matrix's largest (Perron–Frobenius) eigenvalue and its eigenvector say. **Fit** is the exact fit: each shell fills the space round the other, with volumes and Dehn invariants balancing. **Non-periodicity** is that no sliding of the whole lands it on itself. The three are linked: because the scale factor is irrational, a sliding repeat would force repeats at φ^3, φ^6, … times every period, which no single lattice of periods allows; and the same scaling makes every local arrangement recur.

Status. Proved: the shells, the exact fits and the counts. The Chain as built has a centre, so it is trivially not periodic. For a tiling of all space by a finite kit at bounded sizes, two further properties are expected once the kit closes (the FUJI and pyramid side rings, section 4): non-periodicity, via unique composition (Solomyak 1998), and repetitivity (every finite patch recurring within a bounded distance of every point).

10. Relation to known work, and status

Nesting of the icosahedron and dodecahedron. Golden nestings of the two go back to Pacioli and Kepler. Gray (2002) describes icosahedra at every φ-power, each containing a central icosahedron at φ^{-3} and twelve at φ^{-2} (the arrangement of the Lotus seed's icosahedra), and dodecahedra that "scale pack" with them; the gaps are not filled. The Sierpinski icosahedron keeps the twelve corner icosahedra at $1/\varphi^2$ and leaves the gaps empty. Al Raisi, Koca, Koca and Koc (2026) grow the dodecahedron by τ^n with whole dodecahedra inside, using modified Mosseri–Sadoc tiles, and note that icosahedra cannot be treated this way with those tiles.

Tiles. AXE and CLEO are the Mosseri–Sadoc tetrahedra t6 and t4 at scale 1/φ (Mosseri and Sadoc 1982; Kramer 1982); the six tetrahedra admit no stone inflation (Papadopolos and Ogievetsky 1999; Ogievetsky and Papadopolos 1999), and Koca et al. (2020) state that they do

not inflate the icosahedron past edge τ. Danzer's ABCK tetrahedra (1989), with Dehn invariant zero, and Sydler's theorem (1965) frame the remainder of the pyramid rule.

Nested icosahedral shells in matter. Concentric icosahedral shells are central to the structure of real materials and viruses, as point arrays: Mackay's icosahedral sphere packing (1962) and Frank's icosahedral order in liquids (1952); the Tsai-type cluster of icosahedral quasicrystals and their approximants, concentric dodecahedron, icosahedron and icosidodecahedron shells inside a rhombic triacontahedral envelope (Takakura et al. 2007; Deguchi et al. 2015; Ishimasa 2019); and icosahedral viruses (Caspar and Klug 1962), whose capsids and genomes follow nested icosahedral point arrays at fixed radial ratios, obtained from affine extensions of the icosahedral group or by projection from a six-dimensional lattice (Keef and Twarock 2009; Twarock, Valiunas and Zappa 2015), the same framework applying to nested fullerene cages (Dechant et al. 2014). The Chain of Karma differs in kind: its shells are solid and filled exactly, with whole regular icosahedra among the pieces, and the icosahedron and dodecahedron alternate as shells of each other without end.

Status. In the sources above and those in the study folder we found no description of this solid, exact, alternating filling. The search is not exhaustive; the claim of novelty is limited to it.

Remarks

The following are DICTO's own readings of the work, given as remarks; they are not mathematical claims.

On the two shells: "It is like yin and yang: neither one nor the other." Each solid holds the other at its centre, and each is the other's shell.

On the 600-cell: "It does strike me as remarkably similar to a 4D polytope."

On section 9: "The pattern that doesn't repeat is exactly the pattern that is repeating." DICTO relates the Chain to Kyushindo, the philosophy of Kenshiro Abbe, with its three principles 万物流転 *banbutsu ruten* (all things in constant flux), 律動 *ritsudō* (rhythmic motion) and 調和 *chōwa* (harmony): change, rhythm and harmony, each contained in the others. In section 9's terms they correspond to non-periodicity, scaling and exact fit.

References

1. R. Mosseri and J.-F. Sadoc, in: P. H. Gaskell, J. M. Parker, E. A. Davis (eds.), The Structure of Non-Crystalline Materials 1982 (Taylor & Francis, London, 1983) p. 137. P. Kramer, Non-periodic central space filling with icosahedral symmetry using copies of seven elementary cells, *Acta Cryst.* A38 (1982) 257–264.
2. Z. Papadopolos and O. Ogievetsky, On inflation rules for Mosseri–Sadoc tilings (1999), arXiv:math-ph/9911005; O. Ogievetsky and Z. Papadopolos, On quasiperiodic space tilings, inflation and Dehn invariants (1999), arXiv:math-ph/9910006.
3. N. O. Koca, R. Koc, M. Koca and A. Al-Siyabi, Dodecahedral structures with Mosseri–Sadoc tiles (2020), arXiv:2009.07048.
4. L. Danzer, Three-dimensional analogs of the planar Penrose tilings and quasicrystals, *Discrete Math.* 76 (1989) 1–7.
5. J.-P. Sydler, Conditions nécessaires et suffisantes pour l'équivalence des polyèdres de l'espace euclidien à trois dimensions, *Comment. Math. Helv.* 40 (1965) 43–80.
6. R. W. Gray, icosahedron and dodecahedron scaling pages (2002), rwgrayprojects.com.
7. R. Al Raisi, N. O. Koca, M. Koca, R. Koc, Modified Mosseri–Sadoc tiles from D6 (2026), arXiv:2604.03988.

8. B. Solomyak, Nonperiodicity implies unique composition for self-similar translationally finite tilings, *Discrete Comput. Geom.* 20 (1998) 265–279.
9. H. W. Kroto, J. R. Heath, S. C. O'Brien, R. F. Curl, R. E. Smalley, C60: Buckminsterfullerene, *Nature* 318 (1985) 162–163.
10. M. Goldberg, A class of multi-symmetric polyhedra, *Tohoku Math. J.* 43 (1937) 104–108.
11. D. Ugarte, Curling and closure of graphitic networks under electron-beam irradiation, *Nature* 359 (1992) 707–709.
12. A. L. Mackay, A dense non-crystallographic packing of equal spheres, *Acta Cryst.* 15 (1962) 916–918.
13. F. C. Frank, Supercooling of liquids, *Proc. R. Soc. Lond. A* 215 (1952) 43–46.
14. H. Takakura, C. P. Gómez, A. Yamamoto, M. de Boissieu, A. P. Tsai, Atomic structure of the binary icosahedral Yb–Cd quasicrystal, *Nature Materials* 6 (2007) 58–63.
15. K. Deguchi et al., Crystal structure of superconducting 1/1 cubic Au–Ge–Yb approximant with Tsai-type cluster (2015), arXiv:1501.01722.
16. T. Ishimasa, Icosahedral quasicrystal, 1/1 and 2/1 approximants in Zn-based ternary alloys containing Au and Yb/Tb (2019), arXiv:1910.13078.
17. D. L. D. Caspar, A. Klug, Physical principles in the construction of regular viruses, *Cold Spring Harb. Symp. Quant. Biol.* 27 (1962) 1–24.
18. T. Keef, R. Twarock, Affine extensions of the icosahedral group with applications to the three-dimensional organisation of simple viruses, *J. Math. Biol.* 59 (2009) 287–313.
19. R. Twarock, M. Valiunas, E. Zappa, Orbits of crystallographic embedding of non-crystallographic groups and applications to virology, *Acta Cryst. A* 71 (2015) 569–582.
20. P.-P. Dechant, J. Wardman, T. Keef, R. Twarock, Viruses and fullerenes: symmetry as a common thread?, *Acta Cryst. A* 70 (2014) 162–167.