

Problem Set 1

Barrera Munoz Diego

Due: October 8, 2026

Contents

Question 1: Education	2
Question 2: Political Economy	5

Question 1: Education

A school counselor was curious about the average of IQ of the students in her school and took a random sample of 25 students' IQ scores. The following is the data set:

```
1 y <- c(105, 69, 86, 100, 82, 111, 104, 110, 87, 108, 87, 90, 94, 113, 112, 98,
      80, 97, 95, 111, 114, 89, 95, 126, 98)
```

1. Find a 90% confidence interval for the average student IQ in the school (As a hint, you first need the mean and SD to create a CI):

Answer: Since we don't know the population standard deviation (σ), and the size of our sample is small, we will be working with a t distribution to create a CI.

First, we need to calculate the sample size (n), sample mean ($\bar{y}$), standard deviation (s) and the standard error (SE).

$$SE = \frac{s}{\sqrt{n}}$$

```
1 n <- length(y)
2 y_bar <- mean(y)
3 s <- sd(y)
4 se <- s/sqrt(n)
5 cat(" 'y' has",n,"observations, \n The sample mean is :",y_bar,
6     "\n The standard deviation is: ", s,
7     "\n And the standard error is: ", se)
```

```
'y' has 25 observations,
The sample mean is : 98.44
The standard deviation is: 13.09287
And the standard error is: 2.618575
```

Then we need to calculate the critical value (t). We can use this R code:

```
1 critical_t <- qt(p = 0.95, df = n - 1) # df = 25 - 1 = 24
2 critical_t
```

```
[1] 1.710882
```

Note that we use $p = 0.95$ in the code because we want 5% in the lower tail and 5% in the upper tail, leaving 90% in the centre of the distribution.

Then the 90% CI will be:

$$CI = \bar{y} \pm t(SE)$$

$$CI_{90\%} = 98.44 \pm 1.710882(2.618575) = [93.96, 102.92]$$

The CI can also be obtained with this R code:

```
1 ci90 <- c(y_bar - critical_t * se,
2           y_bar + critical_t * se)
3 print(c("Lower" = round(ci90, 2)[1], "Upper" = round(ci90, 2)[2]))
```

```
Lower Upper
93.96 102.92
```

In repeated random sampling, about 90% of intervals made this way would contain the true school mean.

2. Next, the school counselor was curious whether the average student IQ in her school is higher than the average IQ score (100) among all the schools in the country.

Using the same sample, conduct the appropriate hypothesis test with $\alpha = 0.05$.

Answer: To perform an appropriate hypothesis test, first we need to examine the data and formulate the hypotheses. The sample is small, the students were selected randomly, and IQ scores are measured on an interval scale. The school counselor wants to test whether the average IQ score is greater than 100. Therefore, our null and alternative hypotheses are:

- $H_0: \mu \leq 100$
- $H_a: \mu > 100$

Then, for the test statistic, we will use a right-tailed, one-sample t -test because we want to test whether the population mean μ is *higher* than 100:

$$t = \frac{\bar{y} - \mu_0}{SE}$$

$$t = \frac{98.44 - 100}{2.618575} = -0.5957$$

For a one-sided test with $\alpha = 0.05$ and $df = 24$:

$$t_{0.95,24} \approx 1.711$$

Since $-1.711 < -0.5957 < 1.711$ we **fail to reject the null hypothesis** and therefore there is not enough evidence that the school's mean IQ is greater than 100 (with an $\alpha = 0.05$).

We can verify our results with this R code:

```
1 t.test(y, mu = 100, alternative = "greater")
```

One Sample t-test

```
data: y
t = -0.59574, df = 24, p-value = 0.7215
alternative hypothesis: true mean is greater than 100
95 percent confidence interval:
93.95993      Inf
sample estimates:
mean of x
98.44
```

With this code we can observe that the p-value is 0.7215.

Since 0.7215 is not less than 0.05 (α value), we fail to reject the null hypothesis.

Question 2: Political Economy

Researchers are curious about what affects the amount of money communities spend on addressing homelessness. The following variables constitute our data set about social welfare expenditures in the USA.

State	50 states in US
Y	per capita expenditure on shelters/housing assistance in state
X1	per capita personal income in state
X2	Number of residents per 100,000 that are "financially insecure" in state
X3	Number of people per thousand residing in urban areas in state
Region	1=Northeast, 2= North Central, 3= South, 4=West

Explore the `expenditure` data set and import data into R.

```
1 expenditure <- read.table("https://raw.githubusercontent.com/ASDS-TCD/StatsI_2026/main/datasets/expenditure.txt", header=T)
```

Answer: First, we need to look at the structure of the data and check whether the data set contains any missing values (NAs):

```
1 print(head(expenditure)) # Look at the first rows in the data set.
2 print(dim(expenditure)) # Check the structure (dimensions) of the data set.
3 print(sum(is.na(expenditure))) # Check if the data set has NA values.
```


```
> print(head(expenditure))
STATE Y   X1  X2  X3 Region
1    ME 61 1704 388 399      1
2    NH 68 1885 272 598      1
3    VT 72 1745 397 370      1
4    MA 72 2394 458 868      1
5    RI 62 1966 157 899      1
6    CT 91 2817 162 690      1
> print(dim(expenditure))
[1] 50  6
> print(sum(is.na(expenditure)))
[1] 0
```

The data contain 50 states (rows), six columns and no missing values (NAs).

- Please plot the relationships among Y , $X1$, $X2$, and $X3$? What are the correlations among them (you just need to describe the graph and the relationships among them)?

Answer: We can use a scatterplot matrix to compare every pair of numerical variables with this R code:

```
1 variables <- expenditure[c("Y", "X1", "X2", "X3")] # Extract only the
   required variables to another data set.
2
3 pdf("PS01_figures/scatterplot_matrix.pdf", width = 7, height = 5)
4 ggpairs(variables) # Plot the variables.
5 dev.off()
```



With this graph we can observe each variable plotted against the other ones, showing the relationships between six distinct pairs of variables. Some pairs of variables, including $X1-X3$ and $Y-X1$, appear to have moderate positive correlations. But if we want to better understand their relationships, we need to calculate the correlation coefficient for each pair of variables.

With the correlation coefficients, we can observe that Y (spending) has a moderate positive relationship with $X1$ (income), $X2$ (financial insecurity), and $X3$ (urbanisation). States with higher values of these variables tend to spend more, although the dots are quite spread out and therefore harder to fit into a linear regression model.

$X1$ and $X3$ have the strongest relationship of the six ($r = 0.595$): higher-income states tend to be more urban. $X1$ and $X2$ ($r = 0.206$), and $X2$ and $X3$ ($r = 0.221$), have weak positive relationships.

These are associations; correlation does not establish causation.


- Please plot the relationship between Y and *Region*? On average, which region has the highest per capita expenditure on housing assistance?

Answer: Since *Region* is a categorical variable, we will use a boxplot to compare spending across the four groups. First, we need to create a factor variable, *Region_name*, from *Region*, which is currently coded using the numbers 1 to 4.

```
1 expenditure$Region_name <- factor(expenditure$Region, # Set Region as
  factors as a new variable.
2                                     levels = 1:4,
3                                     labels = c("Northeast",
4                                                "North Central",
5                                                "South",
6                                                "West"))
```

Then we create the boxplot:

```
1 region_means <- tapply(expenditure$Y, expenditure$Region_name, mean) #
  Get the mean for each region.
2
3 pdf("PS01_figures/spending_by_region.pdf", width = 7, height = 7) #
  Create image.
4 boxplot(Y ~ Region_name, data = expenditure,
5         col = "lightgrey", xlab = "Region",
6         ylab = "Per capita expenditure (Y)")
7 points(1:4, region_means, col = "red", pch = 18) # Add a red dot where
  the mean is in the boxplot for each region.
8 legend("topleft", legend = "Mean", col = "red", pch = 18,
9       bty = "n")
10 dev.off()
```



```
1 print(round(region_means, 2))
```

Northeast	North Central	South	West
79.44	83.92	69.19	88.31

The **West** region has the highest average expenditure, at 88.31. The South region has the lowest, at 69.19. The wide spread in the West indicates that spending varies considerably across its states, particularly compared with the North Central region, which has the second-highest average expenditure.


- Please plot the relationship between Y and $X1$? Describe this graph and the relationship. Reproduce the above graph including one more variable *Region* and display different regions with different types of symbols and colors.

Answer: To create this plot, we will use the *ggplot2* package:

```
1 library(ggplot2)
```

First, we put $X1$ (Income) on the horizontal axis and Y (Spending) on the vertical axis with this R code:

```
1 income_plot <- ggplot(expenditure, aes(x = X1, y = Y)) +
2   geom_point(colour = "blue", size = 2) +
3   labs(x = "Per capita personal income (X1)",
4        y = "Per capita expenditure (Y)") +
5   theme_minimal()
6 ggsave("PS01_figures/spending_and_income.pdf", plot = income_plot,
7        width = 7, height = 4)
```




We can observe that there is a **moderate positive relationship** ($r = 0.532$). States with higher per capita income tend to spend more per person on housing assistance. Income and spending are positively associated, but states with similar incomes can have different spending levels.

Adding Region to the same graph:

First, we use the mapping *Region_name* → colour and shape inside *aes()*. Different shapes make the regions easier to distinguish. Then we select four different colours and shapes. Thus, the R code that produces the new plot is:

```
1 region_income_plot <- ggplot(expenditure, aes(x = X1, y = Y, colour =  
2   Region_name, shape = Region_name)) +  
3   geom_point(size = 2.5) +  
4   scale_colour_manual(values = c("blue", "darkgreen", "orange", "red")) +  
5   scale_shape_manual(values = c(15, 16, 17, 18)) +  
6   labs(x = "Per capita personal income (X1)",  
7        y = "Per capita expenditure (Y)",  
8        colour = "Region", shape = "Region") +  
9   theme_minimal()  
10  ggsave("PS01_figures/income_and_spending_by_region.pdf",  
        plot = region_income_plot, width = 7, height = 4)
```



With the updated plot we can observe that many Southern states (orange triangles) are in the lower-income, lower-spending part of the graph. Western states (red diamonds) have a wide spread in spending, including the lowest and highest values.

Note that the groups overlap, so *Region* does not fully explain the relationship.