

# 18-Month Upgrade Presentation

Divakar Kumar

OxWaSP, University of Warwick

13 April 2016

# Outline

## Motivation

- Problem specification
- Langevin Diffusion

## ReScaLE Methodology

- Rejection sampling on diffusion path
- Quasi-stationarity
- ReScaLE Algorithm

## Example-sampling from Cauchy density

- Some outputs
- Current Challenges & Further Research
- Thesis Structure

## Bibliography

# Problem specification

- **How to simulate from an intractable distribution  $\pi$ ?**
- Specifically, we might be interested in Bayesian inference of parameter  $x$  in parameter space.

$$\pi(x) = p(x) \prod_{i=1}^N f_i(x)$$

- MCMC Approach
  - Expensive calculation of product in every iteration in MCMC.
  - Memory bottleneck.
- Break  $\longrightarrow$  *Compute Posterior*  $\longrightarrow$  *Recombine*<sup>1</sup>
- Gradient Method

---

<sup>1</sup>Scott, S. L., Blocker, A. W., Bonassi, F. V., Chipman, H. A., George, E. I., and McCulloch, R. E.

# Problem specification

- **How to simulate from an intractable distribution  $\pi$ ?**
- Specifically, we might be interested in Bayesian inference of parameter  $x$  in parameter space.

$$\pi(x) = p(x) \prod_{i=1}^N f_i(x)$$

- MCMC Approach
  - Expensive calculation of product in every iteration in MCMC.
  - Memory bottleneck.
- Break  $\longrightarrow$  *Compute Posterior*  $\longrightarrow$  *Recombine*<sup>1</sup>
- Gradient Method

---

<sup>1</sup>Scott, S. L., Blocker, A. W., Bonassi, F. V., Chipman, H. A., George, E. I., and McCulloch, R. E.

# Problem specification

- **How to simulate from an intractable distribution  $\pi$ ?**
- Specifically, we might be interested in Bayesian inference of parameter  $x$  in parameter space.

$$\pi(x) = p(x) \prod_{i=1}^N f_i(x)$$

- MCMC Approach
  - Expensive calculation of product in every iteration in MCMC.
  - Memory bottleneck.
- Break  $\longrightarrow$  *Compute Posterior*  $\longrightarrow$  *Recombine*<sup>1</sup>
- Gradient Method

---

<sup>1</sup>Scott, S. L., Blocker, A. W., Bonassi, F. V., Chipman, H. A., George, E. I., and McCulloch, R. E. (2013). Bayes and Big Data : The Consensus Monte Carlo Algorithm. pages 1–22

# Problem specification

- **How to simulate from an intractable distribution  $\pi$ ?**
- Specifically, we might be interested in Bayesian inference of parameter  $x$  in parameter space.

$$\pi(x) = p(x) \prod_{i=1}^N f_i(x)$$

- MCMC Approach
  - Expensive calculation of product in every iteration in MCMC.
  - Memory bottleneck.
- Break  $\longrightarrow$  *Compute Posterior*  $\longrightarrow$  *Recombine*<sup>1</sup>
- Gradient Method

---

<sup>1</sup>Scott, S. L., Blocker, A. W., Bonassi, F. V., Chipman, H. A., George, E. I., and McCulloch, R. E.

# Problem specification

- **How to simulate from an intractable distribution  $\pi$ ?**
- Specifically, we might be interested in Bayesian inference of parameter  $x$  in parameter space.

$$\pi(x) = p(x) \prod_{i=1}^N f_i(x)$$

- MCMC Approach
  - Expensive calculation of product in every iteration in MCMC.
  - Memory bottleneck.
- Break  $\longrightarrow$  *Compute Posterior*  $\longrightarrow$  *Recombine*<sup>1</sup>
- Gradient Method

---

<sup>1</sup>Scott, S. L., Blocker, A. W., Bonassi, F. V., Chipman, H. A., George, E. I., and McCulloch, R. E.

# Gradient Method

## Langevin Diffusion

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t \quad X_0 = x, t \in [0, T]$$

- Invariant distribution of above diffusion is  $\pi$ .<sup>1</sup>
- How do we simulate?
  - Metropolis Adjusted Langevin Algorithm (MALA)<sup>2</sup>
  - Euler-Maruyama Scheme

$$X_{t+\delta} = X_t + \frac{1}{2} \nabla \log \pi(X_t) \delta + \epsilon, \quad \epsilon \sim N(0, \delta)$$

- Pitfalls:
  - Non-exact method
  - Infinite time horizon for convergence.
  - Storage bottleneck.
  - High computational cost.
  - Discretisation biases.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

<sup>2</sup>Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations. *Bernoulli*, 2(4):pp. 341–363



# Gradient Method

## Langevin Diffusion

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t \quad X_0 = x, t \in [0, T]$$

- Invariant distribution of above diffusion is  $\pi$ .<sup>1</sup>
- How do we simulate?
  - Metropolis Adjusted Langevin Algorithm (MALA)<sup>2</sup>
  - Euler-Maruyama Scheme

$$X_{t+\delta} = X_t + \frac{1}{2} \nabla \log \pi(X_t) \delta + \epsilon, \quad \epsilon \sim N(0, \delta)$$

- Pitfalls:
  - Non-exact method
  - Infinite time horizon for convergence.
  - Storage bottleneck.
  - High computational cost.
  - Discretisation biases.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data.  
*Upcoming paper*, (2):1–20

<sup>2</sup>Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations.  
*Bernoulli*, 2(4):pp. 341–363

# Gradient Method

## Langevin Diffusion

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t \quad X_0 = x, t \in [0, T]$$

- Invariant distribution of above diffusion is  $\pi$ .<sup>1</sup>
- How do we simulate?
  - Metropolis Adjusted Langevin Algorithm (MALA)<sup>2</sup>
  - Euler-Maruyama Scheme

$$X_{t+\delta} = X_t + \frac{1}{2} \nabla \log \pi(X_t) \delta + \epsilon, \quad \epsilon \sim N(0, \delta)$$

- Pitfalls:
  - Non-exact method
  - Infinite time horizon for convergence.
  - Storage bottleneck.
  - High computational cost.
  - Discretisation biases.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data.  
*Upcoming paper*, (2):1–20

<sup>2</sup>Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations.  
*Bernoulli*, 2(4):pp. 341–363

# Gradient Method

## Langevin Diffusion

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t \quad X_0 = x, t \in [0, T]$$

- Invariant distribution of above diffusion is  $\pi$ .<sup>1</sup>
- How do we simulate?
  - Metropolis Adjusted Langevin Algorithm (MALA)<sup>2</sup>
  - Euler-Maruyama Scheme

$$X_{t+\delta} = X_t + \frac{1}{2} \nabla \log \pi(X_t) \delta + \epsilon, \quad \epsilon \sim N(0, \delta)$$

- Pitfalls:
  - Non-exact method
  - Infinite time horizon for convergence.
  - Storage bottleneck.
  - High computational cost.
  - Discretisation biases.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

<sup>2</sup>Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations. *Bernoulli*, 2(4):pp. 341–363

# Gradient Method

## Langevin Diffusion

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t \quad X_0 = x, t \in [0, T]$$

- Invariant distribution of above diffusion is  $\pi$ .<sup>1</sup>
- How do we simulate?
  - Metropolis Adjusted Langevin Algorithm (MALA)<sup>2</sup>
  - Euler-Maruyama Scheme

$$X_{t+\delta} = X_t + \frac{1}{2} \nabla \log \pi(X_t) \delta + \epsilon, \quad \epsilon \sim N(0, \delta)$$

- Pitfalls:
  - Non-exact method
  - Infinite time horizon for convergence.
  - Storage bottleneck.
  - High computational cost.
  - Discretisation biases.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

<sup>2</sup>Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations. *Bernoulli*, 2(4):pp. 341–363

# Gradient Method

## Langevin Diffusion

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t \quad X_0 = x, t \in [0, T]$$

- Invariant distribution of above diffusion is  $\pi$ .<sup>1</sup>
- How do we simulate?
  - Metropolis Adjusted Langevin Algorithm (MALA)<sup>2</sup>
  - Euler-Maruyama Scheme

$$X_{t+\delta} = X_t + \frac{1}{2} \nabla \log \pi(X_t) \delta + \epsilon, \quad \epsilon \sim N(0, \delta)$$

- Pitfalls:
  - Non-exact method
  - Infinite time horizon for convergence.
  - Storage bottleneck.
  - High computational cost.
  - Discretisation biases.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

<sup>2</sup>Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations. *Bernoulli*, 2(4):pp. 341–363

# Gradient Method

## Langevin Diffusion

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t \quad X_0 = x, t \in [0, T]$$

- Invariant distribution of above diffusion is  $\pi$ .<sup>1</sup>
- How do we simulate?
  - Metropolis Adjusted Langevin Algorithm (MALA)<sup>2</sup>
  - Euler-Maruyama Scheme

$$X_{t+\delta} = X_t + \frac{1}{2} \nabla \log \pi(X_t) \delta + \epsilon, \quad \epsilon \sim N(0, \delta)$$

- Pitfalls:
  - Non-exact method
  - Infinite time horizon for convergence.
  - Storage bottleneck.
  - High computational cost.
  - Discretisation biases.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

<sup>2</sup>Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations. *Bernoulli*, 2(4):pp. 341–363

# Gradient Method

## Langevin Diffusion

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t \quad X_0 = x, t \in [0, T]$$

- Invariant distribution of above diffusion is  $\pi$ .<sup>1</sup>
- How do we simulate?
  - Metropolis Adjusted Langevin Algorithm (MALA)<sup>2</sup>
  - Euler-Maruyama Scheme

$$X_{t+\delta} = X_t + \frac{1}{2} \nabla \log \pi(X_t) \delta + \epsilon, \quad \epsilon \sim N(0, \delta)$$

- Pitfalls:
  - Non-exact method
  - Infinite time horizon for convergence.
  - Storage bottleneck.
  - High computational cost.
  - Discretisation biases.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

<sup>2</sup>Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations. *Bernoulli*, 2(4):pp. 341–363

# Gradient Method

## Langevin Diffusion

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t \quad X_0 = x, t \in [0, T]$$

- Invariant distribution of above diffusion is  $\pi$ .<sup>1</sup>
- How do we simulate?
  - Metropolis Adjusted Langevin Algorithm (MALA)<sup>2</sup>
  - Euler-Maruyama Scheme

$$X_{t+\delta} = X_t + \frac{1}{2} \nabla \log \pi(X_t) \delta + \epsilon, \quad \epsilon \sim N(0, \delta)$$

- Pitfalls:
  - Non-exact method
  - Infinite time horizon for convergence.
  - Storage bottleneck.
  - High computational cost.
  - Discretisation biases.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

<sup>2</sup>Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations. *Bernoulli*, 2(4):pp. 341–363



# How do we simulate Exactly?

- Exact Algorithm <sup>1</sup>: Rejection sampling on diffusion path space
  - Difficult to draw according to target measure  $\mathbb{Q}$ .
  - Choose a proposal measure  $\mathbb{W}$  which is easier to draw from. Requires:
    - $\mathbb{Q}$  is absolutely continuous w.r.t  $\mathbb{W}$  with

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) \leq M$$

- Accept each sample  $X$  with probability  $\frac{1}{M} \frac{d\mathbb{Q}}{d\mathbb{W}}(X)$
- R-N Derivative for Langevin diffusion

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) = \frac{d\mathbb{Q}|_{X_t=y}}{d\mathbb{W}|_{X_t=y}}(X) \frac{\rho_{0,t}(\cdot, y)}{w_{0,t}(\cdot, y)}$$

- The transition density

$$\rho_{0,t}(\cdot, y) = w_{0,t}(\cdot, y) \mathbb{E}_{\mathbb{W}|X_t=y} \left( \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \right)$$

- The transition density  $\rho_{0,t}(\cdot, \cdot) \longrightarrow \pi$  as  $t \longrightarrow \infty$ .

---

<sup>1</sup>Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098

# How do we simulate Exactly?

- Exact Algorithm <sup>1</sup>: Rejection sampling on diffusion path space
  - Difficult to draw according to target measure  $\mathbb{Q}$ .
  - Choose a proposal measure  $\mathbb{W}$  which is easier to draw from. Requires:
    - $\mathbb{Q}$  is absolutely continuous w.r.t  $\mathbb{W}$  with

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) \leq M$$

- Accept each sample  $X$  with probability  $\frac{1}{M} \frac{d\mathbb{Q}}{d\mathbb{W}}(X)$
- R-N Derivative for Langevin diffusion

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) = \frac{d\mathbb{Q}|_{X_t=y}}{d\mathbb{W}|_{X_t=y}}(X) \frac{\rho_{0,t}(\cdot, y)}{w_{0,t}(\cdot, y)}$$

- The transition density

$$\rho_{0,t}(\cdot, y) = w_{0,t}(\cdot, y) \mathbb{E}_{\mathbb{W}|X_t=y} \left( \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \right)$$

- The transition density  $\rho_{0,t}(\cdot, \cdot) \longrightarrow \pi$  as  $t \longrightarrow \infty$ .

---

<sup>1</sup>Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098

# How do we simulate Exactly?

- Exact Algorithm <sup>1</sup>: Rejection sampling on diffusion path space
  - Difficult to draw according to target measure  $\mathbb{Q}$ .
  - Choose a proposal measure  $\mathbb{W}$  which is easier to draw from. Requires:
    - $\mathbb{Q}$  is absolutely continuous w.r.t  $\mathbb{W}$  with

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) \leq M$$

- Accept each sample  $X$  with probability  $\frac{1}{M} \frac{d\mathbb{Q}}{d\mathbb{W}}(X)$
- R-N Derivative for Langevin diffusion

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) = \frac{d\mathbb{Q}|_{X_t=y}}{d\mathbb{W}|_{X_t=y}}(X) \frac{\rho_{0,t}(\cdot, y)}{w_{0,t}(\cdot, y)}$$

- The transition density

$$\rho_{0,t}(\cdot, y) = w_{0,t}(\cdot, y) \mathbb{E}_{\mathbb{W}|X_t=y} \left( \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \right)$$

- The transition density  $\rho_{0,t}(\cdot, \cdot) \longrightarrow \pi$  as  $t \longrightarrow \infty$ .

---

<sup>1</sup>Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098

# How do we simulate Exactly?

- Exact Algorithm <sup>1</sup>: Rejection sampling on diffusion path space
  - Difficult to draw according to target measure  $\mathbb{Q}$ .
  - Choose a proposal measure  $\mathbb{W}$  which is easier to draw from. Requires:
    - $\mathbb{Q}$  is absolutely continuous w.r.t  $\mathbb{W}$  with

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) \leq M$$

- Accept each sample  $X$  with probability  $\frac{1}{M} \frac{d\mathbb{Q}}{d\mathbb{W}}(X)$
- R-N Derivative for Langevin diffusion

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) = \frac{d\mathbb{Q}|_{X_t=y}}{d\mathbb{W}|_{X_t=y}}(X) \frac{\rho_{0,t}(\cdot, y)}{w_{0,t}(\cdot, y)}$$

- The transition density

$$\rho_{0,t}(\cdot, y) = w_{0,t}(\cdot, y) \mathbb{E}_{\mathbb{W}|X_t=y} \left( \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \right)$$

- The transition density  $\rho_{0,t}(\cdot, \cdot) \longrightarrow \pi$  as  $t \longrightarrow \infty$ .

---

<sup>1</sup>Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098

# How do we simulate Exactly?

- Exact Algorithm <sup>1</sup>: Rejection sampling on diffusion path space
  - Difficult to draw according to target measure  $\mathbb{Q}$ .
  - Choose a proposal measure  $\mathbb{W}$  which is easier to draw from. Requires:
    - $\mathbb{Q}$  is absolutely continuous w.r.t  $\mathbb{W}$  with

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) \leq M$$

- Accept each sample  $X$  with probability  $\frac{1}{M} \frac{d\mathbb{Q}}{d\mathbb{W}}(X)$
- R-N Derivative for Langevin diffusion

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) = \frac{d\mathbb{Q}|_{X_t=y}}{d\mathbb{W}|_{X_t=y}}(X) \frac{\rho_{0,t}(\cdot, y)}{w_{0,t}(\cdot, y)}$$

- The transition density

$$\rho_{0,t}(\cdot, y) = w_{0,t}(\cdot, y) \mathbb{E}_{\mathbb{W}|X_t=y} \left( \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \right)$$

- The transition density  $\rho_{0,t}(\cdot, \cdot) \longrightarrow \pi$  as  $t \longrightarrow \infty$ .

---

<sup>1</sup>Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098

# How do we simulate Exactly?

- Exact Algorithm <sup>1</sup>: Rejection sampling on diffusion path space
  - Difficult to draw according to target measure  $\mathbb{Q}$ .
  - Choose a proposal measure  $\mathbb{W}$  which is easier to draw from. Requires:
    - $\mathbb{Q}$  is absolutely continuous w.r.t  $\mathbb{W}$  with

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) \leq M$$

- Accept each sample  $X$  with probability  $\frac{1}{M} \frac{d\mathbb{Q}}{d\mathbb{W}}(X)$
- R-N Derivative for Langevin diffusion

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) = \frac{d\mathbb{Q}|_{X_t=y}}{d\mathbb{W}|_{X_t=y}}(X) \frac{\rho_{0,t}(\cdot, y)}{w_{0,t}(\cdot, y)}$$

- The transition density

$$\rho_{0,t}(\cdot, y) = w_{0,t}(\cdot, y) \mathbb{E}_{\mathbb{W}|X_t=y} \left( \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \right)$$

- The transition density  $\rho_{0,t}(\cdot, \cdot) \longrightarrow \pi$  as  $t \longrightarrow \infty$ .

---

<sup>1</sup>Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098

# How do we simulate Exactly?

- Exact Algorithm <sup>1</sup>: Rejection sampling on diffusion path space
  - Difficult to draw according to target measure  $\mathbb{Q}$ .
  - Choose a proposal measure  $\mathbb{W}$  which is easier to draw from. Requires:
    - $\mathbb{Q}$  is absolutely continuous w.r.t  $\mathbb{W}$  with

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) \leq M$$

- Accept each sample  $X$  with probability  $\frac{1}{M} \frac{d\mathbb{Q}}{d\mathbb{W}}(X)$
- R-N Derivative for Langevin diffusion

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) = \frac{d\mathbb{Q}|_{X_t=y}}{d\mathbb{W}|_{X_t=y}}(X) \frac{\rho_{0,t}(\cdot, y)}{w_{0,t}(\cdot, y)}$$

- The transition density

$$\rho_{0,t}(\cdot, y) = w_{0,t}(\cdot, y) \mathbb{E}_{\mathbb{W}|X_t=y} \left( \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \right)$$

- The transition density  $\rho_{0,t}(\cdot, \cdot) \longrightarrow \pi$  as  $t \longrightarrow \infty$ .

---

<sup>1</sup>Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098.

# How do we simulate Exactly?

- Exact Algorithm <sup>1</sup>: Rejection sampling on diffusion path space
  - Difficult to draw according to target measure  $\mathbb{Q}$ .
  - Choose a proposal measure  $\mathbb{W}$  which is easier to draw from. Requires:
    - $\mathbb{Q}$  is absolutely continuous w.r.t  $\mathbb{W}$  with

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) \leq M$$

- Accept each sample  $X$  with probability  $\frac{1}{M} \frac{d\mathbb{Q}}{d\mathbb{W}}(X)$
- R-N Derivative for Langevin diffusion

$$\frac{d\mathbb{Q}}{d\mathbb{W}}(X) = \frac{d\mathbb{Q}|_{X_t=y}}{d\mathbb{W}|_{X_t=y}}(X) \frac{\rho_{0,t}(\cdot, y)}{w_{0,t}(\cdot, y)}$$

- The transition density

$$\rho_{0,t}(\cdot, y) = w_{0,t}(\cdot, y) \mathbb{E}_{\mathbb{W}|X_t=y} \left( \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \right)$$

- The transition density  $\rho_{0,t}(\cdot, \cdot) \longrightarrow \pi$  as  $t \longrightarrow \infty$ .

---

<sup>1</sup>Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098.



# Transition Density <sup>1</sup>

For  $\mu(x) = \frac{1}{2} \nabla \log \pi(x)$ ,

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x)^{\frac{1}{2}} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$
$$l := \inf_x \frac{\mu^2 + \mu'}{2}(x) \quad \phi(X_s) := \frac{((\mu(X_s))^2 + \mu'(X_s))}{2} - l$$

- It is still difficult to draw according to  $\rho_{0,t}(0, x)$ .
- Drop  $\pi(x)^{\frac{1}{2}}$  and converge to wrong density  $\pi(x)^{\frac{1}{2}}$ .
- Double the drift!

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x) \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- Drop  $\pi(x)$  again and converge to correct density  $\pi(x)$  !

---

<sup>1</sup>Dacunha-Castelle, D. and Florens-Zmirou, D. (1986). Estimation of the Coefficients of a Diffusion from Discrete Observations. *Stochastics*, 19(4):263–284

# Transition Density <sup>1</sup>

For  $\mu(x) = \frac{1}{2} \nabla \log \pi(x)$ ,

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x)^{\frac{1}{2}} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$
$$l := \inf_x \frac{\mu^2 + \mu'}{2}(x) \quad \phi(X_s) := \frac{((\mu(X_s))^2 + \mu'(X_s))}{2} - l$$

- It is still difficult to draw according to  $\rho_{0,t}(0, x)$ .
- Drop  $\pi(x)^{\frac{1}{2}}$  and converge to wrong density  $\pi(x)^{\frac{1}{2}}$ .
- Double the drift!

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x) \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- Drop  $\pi(x)$  again and converge to correct density  $\pi(x)$  !

---

<sup>1</sup>Dacunha-Castelle, D. and Florens-Zmirou, D. (1986). Estimation of the Coefficients of a Diffusion from Discrete Observations. *Stochastics*, 19(4):263–284

# Transition Density <sup>1</sup>

For  $\mu(x) = \frac{1}{2} \nabla \log \pi(x)$ ,

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x)^{\frac{1}{2}} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$
$$l := \inf_x \frac{\mu^2 + \mu'}{2}(x) \quad \phi(X_s) := \frac{((\mu(X_s))^2 + \mu'(X_s))}{2} - l$$

- It is still difficult to draw according to  $\rho_{0,t}(0, x)$ .
- Drop  $\pi(x)^{\frac{1}{2}}$  and converge to wrong density  $\pi(x)^{\frac{1}{2}}$ .
- Double the drift!

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x) \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- Drop  $\pi(x)$  again and converge to correct density  $\pi(x)$  !

---

<sup>1</sup>Dacunha-Castelle, D. and Florens-Zmirou, D. (1986). Estimation of the Coefficients of a Diffusion from Discrete Observations. *Stochastics*, 19(4):263–284

# Transition Density <sup>1</sup>

For  $\mu(x) = \frac{1}{2} \nabla \log \pi(x)$ ,

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x)^{\frac{1}{2}} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$
$$l := \inf_x \frac{\mu^2 + \mu'}{2}(x) \quad \phi(X_s) := \frac{((\mu(X_s))^2 + \mu'(X_s))}{2} - l$$

- It is still difficult to draw according to  $\rho_{0,t}(0, x)$ .
- Drop  $\pi(x)^{\frac{1}{2}}$  and converge to wrong density  $\pi(x)^{\frac{1}{2}}$ .
- Double the drift!

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x) \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- Drop  $\pi(x)$  again and converge to correct density  $\pi(x)$  !

---

<sup>1</sup>Dacunha-Castelle, D. and Florens-Zmirou, D. (1986). Estimation of the Coefficients of a Diffusion from Discrete Observations. *Stochastics*, 19(4):263–284

# Transition Density <sup>1</sup>

For  $\mu(x) = \frac{1}{2} \nabla \log \pi(x)$ ,

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x)^{\frac{1}{2}} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$
$$l := \inf_x \frac{\mu^2 + \mu'}{2}(x) \quad \phi(X_s) := \frac{((\mu(X_s))^2 + \mu'(X_s))}{2} - l$$

- It is still difficult to draw according to  $\rho_{0,t}(0, x)$ .
- Drop  $\pi(x)^{\frac{1}{2}}$  and converge to wrong density  $\pi(x)^{\frac{1}{2}}$ .
- Double the drift!

$$\rho_{0,t}(0, x) \propto \exp \left\{ -\frac{(x)^2}{2t} \right\} \pi(x) \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- Drop  $\pi(x)$  again and converge to correct density  $\pi(x)$  !

---

<sup>1</sup>Dacunha-Castelle, D. and Florens-Zmirou, D. (1986). Estimation of the Coefficients of a Diffusion from Discrete Observations. *Stochastics*, 19(4):263–284

# Quasi-stationarity

$$q_{0,t}(0, x) = \exp \left\{ -\frac{(x)^2}{2t} \right\} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- $q_{0,t}(0, x)$  is the quasi-stationary density of a killed Brownian motion<sup>1</sup> with state dependent killing rate  $\phi(X_s)$ .
- **Problem-1:** How to continuously sample trajectory of a Brownian motion?
- **Problem-2:** How to simulate the quasi-stationary density of a killed Brownian motion?
- **Solution-1:** Need to simulate finite dimensional distribution of sample path of Brownian motion.
- **Solution-2:** Need a mechanism to simulate from the quasi-stationary density of a killed Brownian motion.
  - ScaLE Algorithm uses SMC approach to sample from QSD.
  - ReScaLE uses Glynn & Blanchet method to sample from QSD.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data.  
*Upcoming paper*, (2):1–20

# Quasi-stationarity

$$q_{0,t}(0, x) = \exp \left\{ -\frac{(x)^2}{2t} \right\} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- $q_{0,t}(0, x)$  is the quasi-stationary density of a killed Brownian motion<sup>1</sup> with state dependent killing rate  $\phi(X_s)$ .
- **Problem-1:** How to continuously sample trajectory of a Brownian motion?
- **Problem-2:** How to simulate the quasi-stationary density of a killed Brownian motion?
- **Solution-1:** Need to simulate finite dimensional distribution of sample path of Brownian motion.
- **Solution-2:** Need a mechanism to simulate from the quasi-stationary density of a killed Brownian motion.
  - ScaLE Algorithm uses SMC approach to sample from QSD.
  - ReScaLE uses Glynn & Blanchet method to sample from QSD.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data.  
*Upcoming paper*, (2):1–20

# Quasi-stationarity

$$q_{0,t}(0, x) = \exp \left\{ -\frac{(x)^2}{2t} \right\} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- $q_{0,t}(0, x)$  is the quasi-stationary density of a killed Brownian motion<sup>1</sup> with state dependent killing rate  $\phi(X_s)$ .
- **Problem-1:** How to continuously sample trajectory of a Brownian motion?
- **Problem-2:** How to simulate the quasi-stationary density of a killed Brownian motion?
- **Solution-1:** Need to simulate finite dimensional distribution of sample path of Brownian motion.
- **Solution-2:** Need a mechanism to simulate from the quasi-stationary density of a killed Brownian motion.
  - ScaLE Algorithm uses SMC approach to sample from QSD.
  - ReScaLE uses Glynn & Blanchet method to sample from QSD.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20



# Quasi-stationarity

$$q_{0,t}(0, x) = \exp \left\{ -\frac{(x)^2}{2t} \right\} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- $q_{0,t}(0, x)$  is the quasi-stationary density of a killed Brownian motion<sup>1</sup> with state dependent killing rate  $\phi(X_s)$ .
- **Problem-1:** How to continuously sample trajectory of a Brownian motion?
- **Problem-2:** How to simulate the quasi-stationary density of a killed Brownian motion?
- **Solution-1:** Need to simulate finite dimensional distribution of sample path of Brownian motion.
- **Solution-2:** Need a mechanism to simulate from the quasi-stationary density of a killed Brownian motion.
  - ScaLE Algorithm uses SMC approach to sample from QSD.
  - ReScaLE uses Glynn & Blanchet method to sample from QSD.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

# Quasi-stationarity

$$q_{0,t}(0, x) = \exp \left\{ -\frac{(x)^2}{2t} \right\} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- $q_{0,t}(0, x)$  is the quasi-stationary density of a killed Brownian motion<sup>1</sup> with state dependent killing rate  $\phi(X_s)$ .
- **Problem-1:** How to continuously sample trajectory of a Brownian motion?
- **Problem-2:** How to simulate the quasi-stationary density of a killed Brownian motion?
- **Solution-1:** Need to simulate finite dimensional distribution of sample path of Brownian motion.
- **Solution-2:** Need a mechanism to simulate from the quasi-stationary density of a killed Brownian motion.
  - ScaLE Algorithm uses SMC approach to sample from QSD.
  - ReScaLE uses Glynn & Blanchet method to sample from QSD.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

# Quasi-stationarity

$$q_{0,t}(0, x) = \exp \left\{ -\frac{(x)^2}{2t} \right\} \mathbf{E}_{\mathbb{W}|X_t=x} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- $q_{0,t}(0, x)$  is the quasi-stationary density of a killed Brownian motion<sup>1</sup> with state dependent killing rate  $\phi(X_s)$ .
- **Problem-1:** How to continuously sample trajectory of a Brownian motion?
- **Problem-2:** How to simulate the quasi-stationary density of a killed Brownian motion?
- **Solution-1:** Need to simulate finite dimensional distribution of sample path of Brownian motion.
- **Solution-2:** Need a mechanism to simulate from the quasi-stationary density of a killed Brownian motion.
  - ScaLE Algorithm uses SMC approach to sample from QSD.
  - ReScaLE uses Glynn & Blanchet method to sample from QSD.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

# Quasi-stationarity

$$q_{0,t}(0, \mathbf{x}) = \exp \left\{ -\frac{(\mathbf{x})^2}{2t} \right\} \mathbf{E}_{\mathbb{W} | X_t = \mathbf{x}} \left( \exp \left\{ -\int_0^t \phi(X_s) ds \right\} \right)$$

- $q_{0,t}(0, \mathbf{x})$  is the quasi-stationary density of a killed Brownian motion<sup>1</sup> with state dependent killing rate  $\phi(X_s)$ .
- **Problem-1:** How to continuously sample trajectory of a Brownian motion?
- **Problem-2:** How to simulate the quasi-stationary density of a killed Brownian motion?
- **Solution-1:** Need to simulate finite dimensional distribution of sample path of Brownian motion.
- **Solution-2:** Need a mechanism to simulate from the quasi-stationary density of a killed Brownian motion.
  - ScaLE Algorithm uses SMC approach to sample from QSD.
  - ReScaLE uses Glynn & Blanchet method to sample from QSD.

---

<sup>1</sup>Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20

# Solution-1:

## Coloring Scheme <sup>1</sup>

Let  $\tau_1, \dots, \tau_k$  be the Poisson Process with rate  $M$  where  $M$  is such that  $\sup_x \phi(x) \leq M$ . Let  $X_{\tau_1}, \dots, X_{\tau_k}$  be the realised skeleton of a Brownian motion  $\{X_t : t \geq 0\}$  at times  $\tau_1, \dots, \tau_k$ . If process is killed at  $\tau_j$  with probability  $\frac{\phi(X_{\tau_j})}{M}$ . Then,

$$\mathbb{P}(\text{Process survived until time } t) = \exp \left\{ - \int_0^t \phi(X_s) ds \right\}$$

- Suggests to simulate  $\tau_1, \dots, \tau_k$  from homogeneous Poisson Process of rate  $M$  and decide to kill the process at time of event  $\tau_j$  with probability  $\frac{\phi(X_{\tau_j})}{M}$ .

---

<sup>1</sup>Kingman, J. F. C. (1993). *Poisson Processes*.  
The Clarendon Press Oxford University Press, New York

# Solution-1:

## Coloring Scheme <sup>1</sup>

Let  $\tau_1, \dots, \tau_k$  be the Poisson Process with rate  $M$  where  $M$  is such that  $\sup_x \phi(x) \leq M$ . Let  $X_{\tau_1}, \dots, X_{\tau_k}$  be the realised skeleton of a Brownian motion  $\{X_t : t \geq 0\}$  at times  $\tau_1, \dots, \tau_k$ . If process is killed at  $\tau_j$  with probability  $\frac{\phi(X_{\tau_j})}{M}$ . Then,

$$\mathbb{P}(\text{Process survived until time } t) = \exp \left\{ - \int_0^t \phi(X_s) ds \right\}$$

- Suggests to simulate  $\tau_1, \dots, \tau_k$  from homogeneous Poisson Process of rate  $M$  and decide to kill the process at time of event  $\tau_j$  with probability  $\frac{\phi(X_{\tau_j})}{M}$ .

---

<sup>1</sup>Kingman, J. F. C. (1993). *Poisson Processes*.  
The Clarendon Press Oxford University Press, New York

# Solution-2: Algorithm by Glynn and Blanchet<sup>1</sup>

## Estimating Quasi-Stationary distribution

- 1 Initialize the probability vector  $\pi = \pi_0$  on the non-absorbing states of Markov chain.
- 2 Select a non-absorbing state of the Markov chain  $x_0$  and set  $X_0 = x_0$ .
- 3 Simulate the Markov chain normally starting with  $X_0$  until absorption. Update  $\pi$  by counting the number of visits to each state until absorption.
- 4 Choose an initial position according to normalized vector  $\pi$  and goto step 3.
- 5 Steps 3. and 4. are repeated many times to get an estimate of quasi-stationary dist.

---

<sup>1</sup>Blanchet, J., Glynn, P., and Zheng, S. (2012). Empirical Analysis of a Stochastic Approximation Approach for Computing Quasi-stationary Distributions. *Evolve - A bridge between Probability*, pages 19–37

# Solution-2: Algorithm by Glynn and Blanchet<sup>1</sup>

## Estimating Quasi-Stationary distribution

- 1 Initialize the probability vector  $\pi = \pi_0$  on the non-absorbing states of Markov chain.
- 2 Select a non-absorbing state of the Markov chain  $x_0$  and set  $X_0 = x_0$ .
- 3 Simulate the Markov chain normally starting with  $X_0$  until absorption. Update  $\pi$  by counting the number of visits to each state until absorption.
- 4 Choose an initial position according to normalized vector  $\pi$  and goto step 3.
- 5 Steps 3. and 4. are repeated many times to get an estimate of quasi-stationary dist.

---

<sup>1</sup>Blanchet, J., Glynn, P., and Zheng, S. (2012). Empirical Analysis of a Stochastic Approximation Approach for Computing Quasi-stationary Distributions. *Evolve - A bridge between Probability*, pages 19–37



# Solution-2: Algorithm by Glynn and Blanchet<sup>1</sup>

## Estimating Quasi-Stationary distribution

- 1 Initialize the probability vector  $\pi = \pi_0$  on the non-absorbing states of Markov chain.
- 2 Select a non-absorbing state of the Markov chain  $x_0$  and set  $X_0 = x_0$ .
- 3 Simulate the Markov chain normally starting with  $X_0$  until absorption. Update  $\pi$  by counting the number of visits to each state until absorption.
- 4 Choose an initial position according to normalized vector  $\pi$  and goto step 3.
- 5 Steps 3. and 4. are repeated many times to get an estimate of quasi-stationary dist.

---

<sup>1</sup>Blanchet, J., Glynn, P., and Zheng, S. (2012). Empirical Analysis of a Stochastic Approximation Approach for Computing Quasi-stationary Distributions. *Evolve - A bridge between Probability*, pages 19–37

# Solution-2: Algorithm by Glynn and Blanchet<sup>1</sup>

## Estimating Quasi-Stationary distribution

- 1 Initialize the probability vector  $\pi = \pi_0$  on the non-absorbing states of Markov chain.
- 2 Select a non-absorbing state of the Markov chain  $x_0$  and set  $X_0 = x_0$ .
- 3 Simulate the Markov chain normally starting with  $X_0$  until absorption. Update  $\pi$  by counting the number of visits to each state until absorption.
- 4 Choose an initial position according to normalized vector  $\pi$  and goto step 3.
- 5 Steps 3. and 4. are repeated many times to get an estimate of quasi-stationary dist.

---

<sup>1</sup>Blanchet, J., Glynn, P., and Zheng, S. (2012). Empirical Analysis of a Stochastic Approximation Approach for Computing Quasi-stationary Distributions. *Evolve - A bridge between Probability*, pages 19–37

# Solution-2: Algorithm by Glynn and Blanchet<sup>1</sup>

## Estimating Quasi-Stationary distribution

- 1 Initialize the probability vector  $\pi = \pi_0$  on the non-absorbing states of Markov chain.
- 2 Select a non-absorbing state of the Markov chain  $x_0$  and set  $X_0 = x_0$ .
- 3 Simulate the Markov chain normally starting with  $X_0$  until absorption. Update  $\pi$  by counting the number of visits to each state until absorption.
- 4 Choose an initial position according to normalized vector  $\pi$  and goto step 3.
- 5 Steps 3. and 4. are repeated many times to get an estimate of quasi-stationary dist.

---

<sup>1</sup>Blanchet, J., Glynn, P., and Zheng, S. (2012). Empirical Analysis of a Stochastic Approximation Approach for Computing Quasi-stationary Distributions. *Evolve - A bridge between Probability*, pages 19–37

# Solution-2: Algorithm by Glynn and Blanchet<sup>1</sup>

## Estimating Quasi-Stationary distribution

- 1 Initialize the probability vector  $\pi = \pi_0$  on the non-absorbing states of Markov chain.
- 2 Select a non-absorbing state of the Markov chain  $x_0$  and set  $X_0 = x_0$ .
- 3 Simulate the Markov chain normally starting with  $X_0$  until absorption. Update  $\pi$  by counting the number of visits to each state until absorption.
- 4 Choose an initial position according to normalized vector  $\pi$  and goto step 3.
- 5 Steps 3. and 4. are repeated many times to get an estimate of quasi-stationary dist.

---

<sup>1</sup>Blanchet, J., Glynn, P., and Zheng, S. (2012). Empirical Analysis of a Stochastic Approximation Approach for Computing Quasi-stationary Distributions. *Evolve - A bridge between Probability*, pages 19–37

# ReScaLE Algorithm - Pseudocode

## Algorithm 2.1: ReScaLE Algorithm( $\mu, x_0$ )

1.  $l \leftarrow \inf_{x \in \mathbb{R}} \frac{\mu^2 + \mu'}{2}, \phi \leftarrow \frac{\mu^2 + \mu'}{2} - l, M \leftarrow \sup_{x \in \mathbb{R}} \phi(x)$
  2.  $t_0 \leftarrow 0; X_{t_0} \leftarrow x_0$
  3.
    - do**  $\left\{ \begin{array}{l} (t_1, t_2, \dots) \sim \text{Poisson Process of rate } M \text{ starting at } t_0 \\ (X_{t_1}, X_{t_2}, \dots) \sim \text{Brownian Motion started at position } X_{t_0} \\ \text{Kill the process at } X_{t_i} \text{ with probability } \phi(X_{t_i})/M \\ \text{exit once kill occurs} \end{array} \right.$
  4. **starting time**  $\sim U[0, t_{\text{kill}}]$
  5. **starting value**  $\sim \text{Brownian Bridge conditioned on neighbors of starting time}$
  6. **GOTO** 2. with  $t_0 \leftarrow t_{\text{kill}}; X_{t_0} \leftarrow \text{starting value}$
- return**  $((X_{t_1}, X_{t_2}, \dots))$

# An Illustration of ReScaLE




Figure: A run of ReScaLE

# An Illustration of ReScaLE






Figure: A run of ReScaLE

# An Example



:  $\phi$ -function




: Density Comparison

Figure: Implementation of ReScaLE algorithm to Cauchy density




# Different Initial Positions


$X_0 = 1$




$X_0 = 2$



$X_0 = -1$



$X_0 = -3$



# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - **Choose Starting Time**  $\sim U[\frac{t_{kill}}{2}, t_{kill}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - **Choose Starting Time**  $\sim U[\frac{t_{kill}}{2}, t_{kill}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - **Choose Starting Time**  $\sim U[\frac{t_{kill}}{2}, t_{kill}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - **Choose Starting Time**  $\sim U[\frac{t_{kill}}{2}, t_{kill}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - Choose **Starting Time**  $\sim U[\frac{t_{kill}}{2}, t_{kill}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - Choose **Starting Time**  $\sim U[\frac{t_{kill}}{2}, t_{kill}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - Choose **Starting Time**  $\sim U[\frac{t_{kill}}{2}, t_{kill}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.



# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - Choose **Starting Time**  $\sim U[\frac{t_{kill}}{2}, t_{kill}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - **Choose Starting Time**  $\sim \mathcal{U}[\frac{t_{\text{kill}}}{2}, t_{\text{kill}}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - **Choose Starting Time**  $\sim \mathcal{U}[\frac{t_{\text{kill}}}{2}, t_{\text{kill}}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - **Choose Starting Time**  $\sim \mathcal{U}[\frac{t_{\text{kill}}}{2}, t_{\text{kill}}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Current Challenges & Further Research

- No formal proof of **Glynn & Blanchet algorithm** for CTMC on general state space.
- So far ReScaLE captures restricted class of models only!
  - What happens if  $\phi := \frac{\mu^2 + \mu'}{2} - l$  is unbounded?
  - What if  $\mu = \nabla \log \pi$  is not differentiable e.g. double exponential?
  - What if  $\pi$  has bounded support?
- ReScaLE shows strong affinity towards its chosen initial position.
  - How to construct an 'adaptive' ReScaLE?
  - It takes infinite time horizon for convergence; can we speed it up?
  - **Choose Starting Time**  $\sim \mathcal{U}[\frac{t_{\text{kill}}}{2}, t_{\text{kill}}]$
- Computational aspects
  - Rate of convergence analysis.
  - Parallel execution of ReScaLE and its theory.
  - Optimal version of ReScaLE.

# Thesis Structure

## 1 Chapter One - Motivation

- Sampling from intractable distributions - problem specification and literature reviews of some current methodologies.
- Literature review of ScaLE method.
- How the method links to Big Data Problem?

## 2 Chapter Two - ReScaLE methodology

- Ways to simulate from quasi-stationary distribution.
- Proof of G-B algorithm in general.
- Formal analysis of ReScaLE.
- Adaptive ReScaLE ?
- ReScaLE for general  $\phi$ -function.

## 3 Chapter Three - Computational aspects

- Adaptive ReScaLE algorithm.
- Parallel execution of ReScaLE.
- Comparison of convergence analysis.

## 4 Chapter Four - An application to real data.

## 5 Chapter Five - Conclusion and further research

# Thesis Structure

## 1 Chapter One - Motivation

- Sampling from intractable distributions - problem specification and literature reviews of some current methodologies.
- Literature review of ScaLE method.
- How the method links to Big Data Problem?

## 2 Chapter Two - ReScaLE methodology

- Ways to simulate from quasi-stationary distribution.
- Proof of G-B algorithm in general.
- Formal analysis of ReScaLE.
- Adaptive ReScaLE ?
- ReScaLE for general  $\phi$ -function.

## 3 Chapter Three - Computational aspects

- Adaptive ReScaLE algorithm.
- Parallel execution of ReScaLE.
- Comparison of convergence analysis.

## 4 Chapter Four - An application to real data.

## 5 Chapter Five - Conclusion and further research

# Thesis Structure

## 1 Chapter One - Motivation

- Sampling from intractable distributions - problem specification and literature reviews of some current methodologies.
- Literature review of ReScaLE method.
- How the method links to Big Data Problem?

## 2 Chapter Two - ReScaLE methodology

- Ways to simulate from quasi-stationary distribution.
- Proof of G-B algorithm in general.
- Formal analysis of ReScaLE.
- Adaptive ReScaLE ?
- ReScaLE for general  $\phi$ -function.

## 3 Chapter Three - Computational aspects

- Adaptive ReScaLE algorithm.
- Parallel execution of ReScaLE.
- Comparison of convergence analysis.

## 4 Chapter Four - An application to real data.

## 5 Chapter Five - Conclusion and further research



# Thesis Structure

## 1 Chapter One - Motivation

- Sampling from intractable distributions - problem specification and literature reviews of some current methodologies.
- Literature review of ScaLE method.
- How the method links to Big Data Problem?

## 2 Chapter Two - ReScaLE methodology

- Ways to simulate from quasi-stationary distribution.
- Proof of G-B algorithm in general.
- Formal analysis of ReScaLE.
- Adaptive ReScaLE ?
- ReScaLE for general  $\phi$ -function.

## 3 Chapter Three - Computational aspects

- Adaptive ReScaLE algorithm.
- Parallel execution of ReScaLE.
- Comparison of convergence analysis.

## 4 Chapter Four - An application to real data.

## 5 Chapter Five - Conclusion and further research

# Thesis Structure

- 1 Chapter One - Motivation
  - Sampling from intractable distributions - problem specification and literature reviews of some current methodologies.
  - Literature review of ScaLE method.
  - How the method links to Big Data Problem?
- 2 Chapter Two - ReScaLE methodology
  - Ways to simulate from quasi-stationary distribution.
  - Proof of G-B algorithm in general.
  - Formal analysis of ReScaLE.
  - Adaptive ReScaLE ?
  - ReScaLE for general  $\phi$ -function.
- 3 Chapter Three - Computational aspects
  - Adaptive ReScaLE algorithm.
  - Parallel execution of ReScaLE.
  - Comparison of convergence analysis.
- 4 Chapter Four - An application to real data.
- 5 Chapter Five - Conclusion and further research


## Towards not being afraid of the big bad data set

Gareth Roberts

(joint work with Paul Fearnhead, Adam Johansen & Murray Pollock)

[Gareth.o.Roberts@warwick.ac.uk](mailto:Gareth.o.Roberts@warwick.ac.uk)

June 2nd, 2015



CRISM



00:50



56:31



# Bibliography

- [1] Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098.
- [2] Blanchet, J., Glynn, P., and Zheng, S. (2012). Empirical Analysis of a Stochastic Approximation Approach for Computing Quasi-stationary Distributions. *Evolve - A bridge between Probability*, pages 19–37.
- [3] Dacunha-Castelle, D. and Florens-Zmirou, D. (1986). Estimation of the Coefficients of a Diffusion from Discrete Observations. *Stochastics*, 19(4):263–284.
- [4] Kingman, J. F. C. (1993). *Poisson Processes*. The Clarendon Press Oxford University Press, New York.
- [5] Roberts, G. O., Pollock, M., Fearnhead, P., and Johansen, A. M. (2016). An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20.
- [6] Roberts, G. O. and Tweedie, R. L. (1996). Exponential Convergence of Langevin Distributions and Their Discrete Approximations. *Bernoulli*, 2(4):pp. 341–363.
- [7] Scott, S. L., Blocker, A. W., Bonassi, F. V., Chipman, H. A., George, E. I., and McCulloch, R. E. (2013). Bayes and Big Data : The Consensus Monte Carlo Algorithm. pages 1–22.