

# Resampled Scalable Langevin Exact Method

*A method to simulate from an intractable distribution*

**Divakar Kumar, Dr. Murray Pollock, Prof. Gareth Roberts**

Department of statistics, University of Warwick

D.Kumar@warwick.ac.uk



## Abstract

The *Scalable Langevin Exact (ScaLE)* is a diffusion based approach to simulate from an intractable distribution. The ScaLE is a recent alternative to gradient based Langevin MCMC schemes such as MALA which circumvents the need to use Metropolis type correction. Consequently, due to this circumvention this methodology is highly applicable to ‘Big Data’ problems. The ScaLE method approximates the intractable distribution of interest with the *quasi-stationary* distribution of a ‘killed’ *Brownian motion*. ReScaLE method uses a different simulation mechanism to sample from the quasi-stationary density of a ‘killed’ Brownian motion.

## Introduction

*Re-sampled Scalable Langevin Exact (ReScaLE)* method is a novel method of sampling from an intractable distribution of interest. Contrary to ScaLE, ReScaLE method uses an alternative approach to simulate from the quasi-stationary distribution of a killed Brownian motion whose invariant distribution is given by concerned intractable distribution. The aim of this poster is to furnish an introduction ReScaLE methodology as a tool to sample from an intractable distribution of interest. ReScaLE method uses a recent advancement in the quasi-stationary literature proposed by Glynn & Blanchet.

## Main Objective

- **HOW TO SIMULATE FROM AN INTRACTABLE DISTRIBUTION  $\pi$  ?.**
- **Simulate the stationary distribution of Langevin diffusion**

$$dX_t = \frac{1}{2} \nabla \log \pi(X_t) dt + dB_t, \quad X_0 = x_0, t \in [0, T]. \quad (1)$$

## Path-Space Rejection Sampling for diffusion

1. Propose path  $X$  from a measure  $\mathbb{W}$  for the target measure  $\mathbb{Q}$  such that  $\frac{d\mathbb{Q}}{d\mathbb{W}}(X) \leq M$ .
2. Accept the path  $X$  with probability

$$P_{\mathbb{W}}(X) := \frac{1}{M} \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \quad (2)$$

## Exactly simulating the trajectories of Langevin diffusion

$$p_{0,t}(\cdot, y) = w_{0,t}(\cdot, y) \mathbb{E}_{\mathbb{W}_{|X_t=y}} \left( \frac{d\mathbb{Q}}{d\mathbb{W}}(X) \right) \quad (3)$$

For  $\mu(x) = \frac{1}{2} \nabla \log \pi(x)$ , the transition density is:

$$p_{0,t}(x_0 = 0, x) \propto \exp \left\{ -\frac{x^2}{2t} \right\} \left\{ \pi(x) \right\}^{\frac{1}{2}} \mathbb{E}_{x_0, x} \left( \exp \left\{ -\int_0^t \phi_{\mu}(X_s) ds \right\} \right) \rightarrow \pi. \quad (4)$$

where

$$l := \inf_x \frac{\mu^2 + \mu'}{2}(x) \quad \phi_{\mu}(X_s) := \frac{(\mu(X_s)^2 + \mu'(X_s))}{2} - l \quad (5)$$

## Double the drift! - Drop $\pi(x)$

$$p_{0,t}(x_0 = 0, x) \propto \exp \left\{ -\frac{x^2}{2t} \right\} \left\{ \pi(x) \right\} \mathbb{E}_{x_0, x} \left( \exp \left\{ -\int_0^t \phi_{2\mu}(X_s) ds \right\} \right) \rightarrow \pi^2. \quad (6)$$

## Killed Brownian Motion

The quasi-stationary density of a killed Brownian motion with killing rate  $\phi_{2\mu}$  is

$$q_{0,t}(0, x) := \exp \left\{ -\frac{x^2}{2t} \right\} \mathbb{E}_{x_0, x} \left( \exp \left\{ -\int_0^t \phi_{2\mu}(X_s) ds \right\} \right). \quad (7)$$

## Problems:

1. **Problem-1:** How to continuously sample trajectory of a Brownian motion?
2. **Problem-2:** How to simulate the quasi-stationary density of a killed Brownian motion?

## Sampling from the QSD of Brownian motion

### Result - 1:

Let  $\tau_1, \dots, \tau_k$  be the Poisson process with rate  $M$  where  $M$  is such that  $\sup_x \phi(x) \leq M$ . Let  $X_{\tau_1}, \dots, X_{\tau_k}$  be the realised skeleton of a Brownian motion  $\{X_t : t \geq 0\}$  at times  $\tau_1, \dots, \tau_k$ . If process is killed at  $\tau_j$  with probability  $\frac{\phi(X_{\tau_j})}{M}$ . Then,

$$\mathbb{P}(\text{Process survived until time } t) = \exp \left\{ -\int_0^t \phi(X_s) ds \right\}$$


### Result - 2: Glynn & Blanchet’s approach of estimating the QSD

1. Initialize the probability vector  $\pi = \pi_0$  on the non-absorbing states of Markov chain.
2. Select a non-absorbing state of the Markov chain  $x_0$  and set  $X_0 = x_0$ .
3. Simulate the Markov chain normally starting with  $X_0$  until absorption. Update  $\pi$  by counting the number of visits to each state until absorption.
4. Choose an initial position according to normalized vector  $\pi$  and goto step 3.
5. Steps 3. and 4. are repeated many times to get an estimate of quasi-stationary dist.

## ReScaLE method - Pseudocode


### Algorithm 0.1: RESCALE ALGORITHM( $\mu, x_0$ )

1.  $l \leftarrow \inf_{x \in \mathbb{R}} \frac{\mu^2 + \mu'}{2}, \phi \leftarrow \frac{\mu^2 + \mu'}{2} - l, M \leftarrow \sup_{x \in \mathbb{R}} \phi(x)$
  2.  $t_0 \leftarrow 0; X_{t_0} \leftarrow x_0$
  3.  $\left\{ \begin{array}{l} (t_1, t_2, \dots) \sim \text{Poisson Process of rate } M \text{ starting at } t_0 \\ (X_{t_1}, X_{t_2}, \dots) \sim \text{Brownian Motion started at position } X_{t_0} \\ \text{Kill the process at } X_{t_i} \text{ with probability } \phi(X_{t_i})/M \\ \text{exit once kill occurs} \end{array} \right.$
  4. **starting time**  $\sim U[0, t_{\text{kill}}]$
  5. **starting value**  $\sim$  Brownian Bridge conditioned on neighbors of **starting time**
  6. **GOTO** 2. with  $t_0 \leftarrow t_{\text{kill}}; X_{t_0} \leftarrow \text{starting value}$
- return**  $((X_{t_1}, X_{t_2}, \dots))$



**Figure 1:** An illustration of ReScaLE methodology

An application of ReScaLE methodology to sample from Cauchy density



## Current challenges and further research

- No formal proof exists for the regenerative algorithm by Glynn and Blanchet for CTMC on general state space.
- Current ReScaLE method shows persistence issues.
- How to make the method ‘adaptive’?
- How to ‘speed-up’ the method for faster convergence to quasi-stationary density?

## References

- [1] A. Beskos, O. Papaspiliopoulos, and G. O. Roberts. Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098, 2006.
- [2] Jose Blanchet, Peter Glynn, and Shuheng Zheng. Empirical Analysis of a Stochastic Approximation Approach for Computing Quasi-stationary Distributions. *Evolve - A bridge between Probability*, pages 19–37, 2012.
- [3] Gareth O. Roberts, Murray Pollock, Paul Fearnhead, and Adam M Johansen. An Unbiased and Scalable Monte Carlo Method for Bayesian Inference for Big Data. *Upcoming paper*, (2):1–20, 2016.
- [4] Gareth O Roberts and Richard L Tweedie. Exponential Convergence of Langevin Distributions and Their Discrete Approximations. *Bernoulli*, 2(4):pp. 341–363, 1996.

## Acknowledgements

This is a joint work with Dr. Murray Pollock & Prof. Gareth Roberts under the Oxford-Warwick Statistics Program (OxWaSP) - a joint DPhil program being run between the department of statistics, University of Oxford and University of Warwick, UK.