

Validating Causal Steering in Protein Language Models: Proxy Accuracy Does Not Guarantee Steering Success

Dmitriy Ivkov
University of Michigan, Ann Arbor
divkov@umich.edu

Abstract

Activation steering changes model output by modifying internal activations. Protein steering studies often use scoring proxies because generated sequences cannot be tested in a wet-lab assay during every model evaluation. We ask whether agreement between a proxy and experimental labels also indicates that the proxy will provide a useful steering target. We test five protein properties in ESM2-650M with difference-of-means steering vectors and a fixed six-requirement evaluation protocol. The binding-affinity proxy has the strongest held-out correlation in the study ($r = 0.80$), but its steering direction has no measurable effect. A weaker catalytic-activity proxy ($r = 0.22$) produces an effect that repeats across three random seeds. We find a similar mismatch for individual attention heads: strong correlation with known 3D contacts does not imply a large effect under ablation. Two additional checks change how we interpret other results. A residue-exclusion check gives different conclusions across seeds, and an apparent expression-yield effect uses a direction that overlaps with the intrinsic-disorder direction. These results show that proxy validation tests the score, while direct intervention tests are needed to evaluate the steering direction.

1 Introduction

Difference-of-means activation steering modifies a model’s residual stream during generation. The steering direction is obtained by subtracting the mean activation of a low-property group from the mean activation of a high-property group. This method has reproduced a known thermostability effect in ESM2-650M [3].

That result leaves an important question open when the method is applied to a new protein property. Generated proteins cannot be evaluated with a wet-lab assay during every model run, so steering studies often depend on a scoring proxy. A strong correlation between the proxy and experimental labels shows that the score contains information about the property. However, it does not show that a difference-of-means vector built from that property will change the model in the intended way.

We test five properties under the same protocol, and four of them reach the steering stage. The order of their proxy correlations does not match the order of their steering effects. The binding-affinity proxy has the strongest held-out correlation, but its steering vector has no detectable effect. The catalytic-activity proxy has a much weaker correlation, yet its effect is consistent across intervention strengths and random seeds.

We test a related assumption at the level of individual model components. An attention head that correlates strongly with known protein contacts is not unusually important when it is ablated. Together, these experiments distinguish evidence that a model represents a property from evidence that a particular intervention controls that property.

This paper evaluates the reliability of steering evidence rather than introducing a new steering algorithm. We apply one protocol to five properties, investigate why two results fail or remain inconclusive, and document two analysis choices that would have changed our conclusions.

2 Related Work

Huang et al. [3] introduced difference-of-means activation steering for protein language models and validated

the method on thermostability. We apply the same intervention to five new properties and test whether proxy accuracy predicts steering success.

Vig et al. [8] identified attention heads whose attention patterns correlate with 3D protein contacts. The original study described this evidence as associative. We reproduce their analysis, test the most enriched head in our data by causal ablation, and extend the analysis to all 480 heads in the model.

Adams et al. [1] used linear probes to identify thermostability and localization information in sparse-autoencoder features from ESM-2. Their steering experiment tested whether a selected latent produced more coherent sequence changes than a random latent. It did not compare the result with an experimental property label. Our study addresses a related but different question: whether correlational evidence predicts the result of a downstream intervention.

3 Methods

3.1 Study design

We study binding affinity, catalytic activity, intrinsic disorder, immunogenicity, and expression yield. We evaluate each property through the same sequence of steps. First, we validate a property-specific score against experimental labels. If the score passes that check, we construct a steering vector, generate sequences at several intervention strengths, and compare the learned vector with a matched random control. The immunogenicity study stops after proxy validation because none of its candidate scores generalizes to the experimental endpoint.

3.2 Activation steering

For each target property, we divide real protein sequences into high-property and low-property groups. At every layer of ESM2-650M, we compute the mean residual-stream activation for each group and subtract the two means. The resulting vector is added to the residual stream during sequence generation. A scalar α controls the intervention strength.

Each generated sequence is evaluated with the property-specific compositional proxy. We compare the property-derived steering direction with a random direction of the same norm. This comparison tests whether the measured change depends on the learned direction rather than only on the size of the intervention.

3.3 Six evaluation requirements

We fix six requirements before testing the new properties:

1. The property-derived direction must outperform a random direction of the same norm in a paired bootstrap test with 10,000 resamples.
2. The effect must increase monotonically across at least three intervention strengths in a range where generation remains stable.
3. The effect must remain after the most frequently substituted residues are excluded from the score.
4. The scoring proxy must agree with experimental labels on held-out data before steering is attempted.
5. The method must outperform an existing baseline when a suitable baseline is available.
6. An evaluation of a new property must contain at least 150 held-out sequences.

The third requirement is a check for compositional collapse. A steering direction can appear successful if it repeatedly inserts one residue that directly increases the proxy score. Excluding the dominant substitutions tests whether the effect extends beyond that narrow change.

We use three assessment outcomes. If proxy validation fails, we reject the proxy and do not attempt steering. If the main steering test succeeds but the residue-exclusion result fails or changes across seeds, we classify the evidence as inconclusive. We call a result supported only when it satisfies every applicable requirement.

Table 1: Results for the five target properties. Proxy r is the Pearson correlation with experimental labels on held-out data, measured before steering.

Target	Dataset	Proxy r	Steering result	Assessment
Binding affinity	ProteinGym DMS (KRAS/DARPin)	0.80	no measurable effect	Unsupported
Catalytic activity	DLKcat (cross-protein)	0.22	repeats across 3 seeds	Supported
Intrinsic disorder	DisProt	0.45	main effect 3/3; check 2/3	Inconclusive
Immunogenicity	IEDB (4-tier)	≤ 0.10	steering not attempted	Proxy rejected
Expression yield	eSol	0.31	fails residue-exclusion check	Inconclusive

3.4 Attention-head ablation

We also examine whether correlational evidence predicts the importance of individual attention heads. Following Vig et al. [8], we measure contact enrichment in their protein language model and select the most enriched head in our data. We ablate that head and compare the change in masked-residue prediction with a low-correlation control head. We then repeat the ablation for all 480 heads in the model and rank them by measured causal effect.

4 Results

4.1 Results across five target properties

Table 1 summarizes the five target properties. Binding affinity has the strongest proxy correlation but no measurable steering effect. Catalytic activity has the weakest proxy correlation among the four properties that reach the steering stage, yet its effect is larger and repeats across seeds.

4.2 Binding affinity

We use a ProteinGym deep-mutational-scanning assay [6] of KRAS binding to a DARPin. The proxy is built from position-specific mutational-sensitivity weights learned on the training split. It scores a sequence according to whether the wild-type residue is preserved at positions where mutations have a large measured effect.

The proxy reaches $r = 0.80$ on held-out data. A weight-shuffle control gives $r = -0.001$, which indicates that the position-specific weights carry the signal. The proxy also generalizes to positions excluded from weight fitting ($r = 0.54$) and from single to double mutants ($r = 0.70$).

Despite this validation, the steering direction has no detectable effect at any tested strength. Generation remains stable, and no strength produces degenerate sequences. Output collapse therefore does not explain the result.

4.3 Catalytic activity

For catalytic activity, we use a cross-protein enzyme turnover dataset from DLKcat [4]. The proxy measures the fraction of glycine minus the fraction of arginine.

The proxy reaches held-out $r = 0.22$, the weakest result among the four targets that reach the steering stage. Even so, the steering effect is significant at every tested strength in the stable range. It increases monotonically, remains after residue exclusion, and repeats across three random seeds.

4.4 Intrinsic disorder

We use DisProt annotations [7]. The intrinsic-disorder proxy uses the published TOP-IDP amino-acid scale [2]. It reaches held-out $r = 0.45$. A sliding-window comparison with per-residue disorder labels gives AUC 0.71.




Figure 1: Proxy correlation and steering effect size for the four targets that reach the steering stage. Effect size is Cohen’s d at $\alpha = 0.5$ for the learned direction relative to the matched random direction. The four points show no positive relation between the two measurements.

The main steering effect is significant and monotonic on all three random seeds. The residue-exclusion check is less stable. It passes on two seeds, retaining 36–42% of the original effect, but fails on the third, where it retains 10% and its confidence interval crosses zero. The effect is repeatable, but the evidence against a simple compositional artifact is not equally strong on every seed.

4.5 Immunogenicity

For immunogenicity, we use records from the Immune Epitope Database [9] and evaluate candidate proxies at four levels: peptide MHC-II binding affinity, mass-spectrometry-confirmed peptide presentation, measured T-cell response, and full-length source antigens. Candidate proxies correlate with the surrogate binding-affinity endpoint at $r = 0.37$ – 0.43 but fall to $r \leq 0.10$ on the measured T-cell-response endpoint.

One proxy for full-length antigens initially reaches $r = 0.38$. Further testing shows that organism identity explains 58% of this signal. When organism identity is held out during cross-validation, the correlation changes to -0.32 . Because no proxy passes the validation requirement, we do not run a steering experiment.

4.6 Expression yield

For expression yield, we use the eSol dataset [5]. The proxy is based on the absolute deviation of net charge from neutrality, following a published recombinant-solubility discriminant [10]. It reaches held-out $r = 0.31$ – 0.34 and produces a significant, monotonic steering effect.

The effect disappears when the dominant residue substitutions are excluded from scoring. One of those residues is part of the proxy definition. The result is therefore inconclusive under the fixed protocol.




Figure 2: Dose-response curves for catalytic activity and intrinsic disorder. Both effects increase monotonically across the tested intervention strengths. Error bars show 95% bootstrap confidence intervals.

4.7 Attention-head ablation

In our replication of Vig et al. [8], the most contact-enriched head has 12.9-fold enrichment over background. Ablating it produces a change in masked-residue prediction that is statistically indistinguishable from ablating a low-correlation control head.

The full sweep supports the same interpretation, but it does not provide a precise rank for this head. The evaluation uses 104 sampled positions, so many heads have identical measured effects. The selected head shares its value with 131 other heads, forming a tied block that spans ranks 286–417. At the same resolution, 166 heads show no measurable effect. Strong contact correlation therefore does not imply a large or unique contribution to masked-residue prediction.

5 Failure Analysis and Reliability Checks

5.1 The binding-affinity groups differ at few residues

The binding-affinity sequences are point mutants of one 188-residue reference protein. The high- and low-binding groups differ by only one or two residues per sequence. Their mean per-residue compositional distance is 0.003, compared with 0.02–0.05 for the three cross-protein datasets.

This difference offers a possible explanation for the failed intervention. The proxy uses position-specific weights, so it can predict binding labels from a small number of mutations. In contrast, the steering direction averages activations over groups of nearly identical sequences. The few mutations that distinguish the groups may contribute too little to that average. Our evidence supports this explanation as a possibility, not as a demonstrated mechanism.

5.2 The expression-yield direction overlaps with disorder

We compare the expression-yield steering vector with every other target vector. Its cosine similarity with the intrinsic-disorder direction is +0.38 across all 33 layers and rises to +0.56–0.67 in the three deepest layers. This overlap shows that the two directions are not orthogonal and may capture some of the same compositional variation. It does not show that disorder causes the expression-yield result. Because the expression-yield result also fails residue exclusion, we do not count it as separate evidence for a new steering effect. The comparison requires one additional embedding pass and is inexpensive relative to a complete steering sweep.




Figure 3: Intrinsic-disorder effects before and after residue exclusion for three random seeds. The main effect is stable, but the confidence interval for the residue-excluded score crosses zero on seed 1.

5.3 Unstable generation made five-layer steering look stronger

In a follow-up thermostability experiment, we compared steering all 33 layers with steering only five layers previously identified as important. An early analysis selected the best intervention strength from the full tested range, including $\alpha \geq 1.0$. At the selected strength, generation under the 33-layer intervention had already collapsed, while generation under the 5-layer intervention had not. The 5-layer version appeared better only because it failed at a higher strength.

We restricted the comparison to a range where both interventions produced stable sequences and reran the complete experiment. The 5-layer intervention then retained 43–59% of the full effect. This is a useful partial result, but it does not support the original claim that five layers perform as well as all 33.

5.4 A robustness criterion changed across seeds

The intrinsic-disorder effect repeats across all three seeds, but its residue-exclusion check passes on only two. A single run could therefore support either a pass or a failure for that criterion. Repeating only the main effect is not enough when a separate artifact check is used to justify the result.

6 Discussion

These results show why representation and intervention should be evaluated separately. A proxy can accurately summarize experimental labels without defining a direction that changes model output in the intended way. Likewise, an attention head can align with a biological pattern without making a large, individually measurable contribution to the model’s prediction.

Proxy validation remains necessary because it rules out scores that do not track the target, as the immunogenicity experiment shows. The problem arises when evidence about the score is also treated as evidence about the steering vector. The binding-affinity experiment shows that the score and the vector can depend on different sources of signal.

The experiments suggest several practical checks. Before constructing a steering vector, researchers should measure how much the high- and low-property groups differ. Comparisons between interventions should use the same range of strengths, and every intervention should produce stable sequences throughout that range. Important results should be repeated across seeds, including the artifact checks used to support them. When several properties are studied together, similarities between their steering vectors can reveal that two apparent effects may not be independent.

Every successful result in this study is causal only with respect to the model’s activation space. The intervention changes generated sequences and outperforms a matched random direction, but no generated protein was tested in a biological assay. A successful protocol result means that the intervention changes a validated proxy. It does not establish that the generated protein has the intended biological property.

7 Limitations

All experiments use ESM2-650M at one model scale. The relationship between proxy accuracy and steering effect could differ in other models or at other scales. The proxies are compositional heuristics rather than learned predictors. A stronger non-compositional proxy could behave differently.

We selected each final proxy from several candidates using the same held-out split on which we report its correlation. The reported correlations are therefore likely optimistic and should be treated as upper bounds. This issue affects the absolute values, although it does not reverse the ordering between the binding-affinity and catalytic-activity proxies.

We ran multiple significance tests without a formal correction. Agreement across intervention strengths and seeds reduces this risk but does not remove it. Seed robustness was checked for only two targets. We also did not evaluate generated sequences with a structural plausibility model or a wet-lab assay.

8 Conclusion

Across these experiments, higher proxy accuracy does not guarantee more successful steering. The binding-affinity result provides the clearest counterexample, and the attention-head ablation shows a related mismatch at the level of a model component. Correlational evidence can show that a model represents a property, but reliable steering claims still require direct tests of the intervention, its artifacts, and its stability.

LLM usage disclosure

Claude (Anthropic) assisted with code writing, experiment scripts, literature searches, and manuscript drafting. The author chose the study design, ran and checked the experiments, and verified each numerical claim against committed outputs.

Code and data availability

Code, data-processing scripts, saved results, and instructions for reproducing the experiments are available at <https://github.com/DIvkov575/esmplm-steering>.

References

- [1] Etowah Adams, Liam Bai, Minji Lee, Yiyang Yu, and Mohammed AlQuraishi. From mechanistic interpretability to mechanistic biology: Training, evaluating, and interpreting sparse autoencoders on protein language models. In *Proceedings of the 42nd International Conference on Machine Learning*, volume 267 of *Proceedings of Machine Learning Research*, pages 460–476, 2025.
- [2] Andrew Campen, Ryan M. Williams, Celeste J. Brown, Jingwei Meng, Vladimir N. Uversky, and A. Keith Dunker. Top-idp-scale: a new amino acid scale measuring propensity for intrinsic disorder. *Protein & Peptide Letters*, 15(9):956–963, 2008. doi: 10.2174/092986608785849164.

- [3] Long-Kai Huang, Rongyi Zhu, Bing He, and Jianhua Yao. Steering protein language models. *arXiv preprint arXiv:2509.07983*, 2025. Accepted to ICML 2025.
- [4] Feiran Li, Le Yuan, Hongzhong Lu, Gang Li, Yu Chen, Martin K. M. Engqvist, Eduard J. Kerkhoven, and Jens Nielsen. Deep learning-based kcat prediction enables improved enzyme-constrained model reconstruction. *Nature Catalysis*, 5(8):662–672, 2022. doi: 10.1038/s41929-022-00798-z.
- [5] Tatsuya Niwa, Bei-Wen Ying, Katsuyo Saito, WenZhen Jin, Shoji Takada, Takuya Ueda, and Hideki Taguchi. Bimodal protein solubility distribution revealed by an aggregation analysis of the entire ensemble of escherichia coli proteins. *Proceedings of the National Academy of Sciences*, 106(11):4201–4206, 2009. doi: 10.1073/pnas.0811922106.
- [6] Pascal Notin, Aaron Kollasch, Daniel Ritter, Lood Van Niekerk, Steffanie Paul, Han Spinner, Nathan Rollins, Ada Shaw, Rose Orenbuch, Ruben Weitzman, Jonathan Frazer, Mafalda Dias, Dinko Franceschi, Yarin Gal, and Debora Marks. Proteingym: Large-scale benchmarks for protein fitness prediction and design. In *Advances in Neural Information Processing Systems*, volume 36, pages 64331–64379, 2023. doi: 10.52202/075280-2810.
- [7] Federica Quaglia et al. Disprot in 2022: improved quality and accessibility of protein intrinsic disorder annotation. *Nucleic Acids Research*, 50(D1):D480–D487, 2022. doi: 10.1093/nar/gkab1082.
- [8] Jesse Vig, Ali Madani, Lav R. Varshney, Caiming Xiong, Richard Socher, and Nazneen Fatema Rajani. Bertology meets biology: Interpreting attention in protein language models. In *International Conference on Learning Representations*, 2021.
- [9] Randi Vita, Swapnil Mahajan, James A. Overton, Sandeep Kumar Dhanda, Sheridan Martini, Jason R. Cantrell, Daniel K. Wheeler, Alessandro Sette, and Bjoern Peters. The immune epitope database (iedb): 2018 update. *Nucleic Acids Research*, 47(D1):D339–D343, 2019. doi: 10.1093/nar/gky1006.
- [10] Derek L. Wilkinson and Roger G. Harrison. Predicting the solubility of recombinant proteins in escherichia coli. *Nature Biotechnology*, 9(5):443–448, 1991. doi: 10.1038/nbt0591-443.