Example: Multiple Joins


1



+ L	0.1	0.9
	+L	_


P(R,T)


Join R












+t | 0.09

-t | 0.81

-t |0.2



L L	_	
	\subseteq	
	_	





0.024	0.056	0.002	0.018	0.027	0.063	0.081	0.729
+	-	+	-	+	-	+	_
÷	+t	-t	-t	+t	+t	-t	-t
<u>_</u>	+r	+r	+L	-Ľ	J-	_L	_L

_	
$ \mathcal{I} $	
$\overline{\Gamma}$	
Ω.	

P(L|T)

0.3	0.7	0.1	6.0
+	-	+	_
+t	+t	-t	- t

+ 0.1

-l | 0.9


0.9	

0.3	0.7	0.1	6.0
+	-	+	_
†	+	-t	-t

$\left\langle \varkappa \right\rangle$	

Operation 2: Eliminate

- Second basic operation: marginalization
- Take a factor and sum out a variable
- Shrinks a factor to a smaller one
- A projection operation



[e:	
$\overline{}$,
9	
III	
a	
×	
\subseteq	
_ `	


0.08	0.02	60'0	0.81
+¢	-t	+t	-t
+L	+r	_L	-r


	7	
1	7	

 $\operatorname{sum} R$

0.17	0.83	
+t	-t	

Multiple Elimination





8