
CS 471/571 (Fall 2023): Introduction to Artificial Intelligence

Lecture 17: Bayes Nets

Thanh H. Nguyen

Source: <http://ai.berkeley.edu/home.html>



Reminder:

- Written assignment 3:
 - Deadline: Nov 08, 2023
- Programming project 3:
 - Will be posted tomorrow
 - Deadline: Nov 20, 2023

Probabilistic Models

- Models describe how (a portion of) the world works
- **Models are always simplifications**
 - May not account for every variable
 - May not account for all interactions between variables
 - “All models are wrong; but some are useful.”
 - George E. P. Box
- What do we do with probabilistic models?
 - We (or our agents) need to reason about unknown variables, given evidence
 - Example: explanation (diagnostic reasoning)
 - Example: prediction (causal reasoning)




Probability Recap

- Conditional probability $P(x|y) = \frac{P(x, y)}{P(y)}$
- Product rule $P(x, y) = P(x|y)P(y)$
- Chain rule
$$\begin{aligned} P(X_1, X_2, \dots, X_n) &= P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots \\ &= \prod_{i=1}^n P(X_i|X_1, \dots, X_{i-1}) \end{aligned}$$



Bayes Rule



Bayes Rule

- Two ways to factor a joint distribution over two variables:

$$P(x, y) = P(x|y)P(y) = P(y|x)P(x)$$

That's my rule!

- Dividing, we get:

$$P(x|y) = \frac{P(y|x)P(x)}{P(y)}$$

- Why is this at all helpful?
 - Lets us build one conditional from its reverse
 - Often one conditional is tricky but the other one is simple
- In the running for most important AI equation!



Quiz

- Given:

$P(W)$

R	P
sun	0.8
rain	0.2


$P(D|W)$

D	W	P
wet	sun	0.1
dry	sun	0.9
wet	rain	0.7
dry	rain	0.3

- What is $P(W \mid \text{dry})$?



Independence




Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- This says that their joint distribution *factors* into a product two simpler distributions
- Another form: $\forall x, y : P(x|y) = P(x)$
- We write: $X \perp\!\!\!\perp Y$
- Independence is a simplifying *modeling assumption*
 - What could we assume for {Weather, Traffic, Cavity, Toothache}?



Example: Independence?

$P_1(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

$P(T)$

T	P
hot	0.5
cold	0.5

$P_2(T, W)$

T	W	P
hot	sun	0.3
hot	rain	0.2
cold	sun	0.3
cold	rain	0.2

$P(W)$



W	P
sun	0.6
rain	0.4



Example: Independence

- N fair, independent coin flips:

$P(X_1)$		$P(X_2)$		\dots		$P(X_n)$	
H	0.5	H	0.5			H	0.5
T	0.5	T	0.5			T	0.5



Conditional Independence

- Unconditional (absolute) independence very rare
- *Conditional independence* is our most basic and robust form of knowledge about uncertain environments.
- X is conditionally independent of Y given Z

$$X \perp\!\!\!\perp Y | Z$$

if and only if:

$$\forall x, y, z : P(x, y | z) = P(x | z)P(y | z)$$


or, equivalently, if and only if

$$\forall x, y, z : P(x | z, y) = P(x | z)$$



Conditional Independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:
 - $P(+\text{catch} \mid +\text{toothache}, +\text{cavity}) = P(+\text{catch} \mid +\text{cavity})$
- The same independence holds if I don't have a cavity:
 - $P(+\text{catch} \mid +\text{toothache}, -\text{cavity}) = P(+\text{catch} \mid -\text{cavity})$
- Catch is *conditionally independent* of Toothache given Cavity:
 - $P(\text{Catch} \mid \text{Toothache}, \text{Cavity}) = P(\text{Catch} \mid \text{Cavity})$
- **Equivalent statements:**
 - $P(\text{Toothache} \mid \text{Catch}, \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity})$
 - $P(\text{Toothache}, \text{Catch} \mid \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity}) P(\text{Catch} \mid \text{Cavity})$
 - One can be derived from the other easily





Conditional Independence

- What about this domain:
 - Traffic
 - Umbrella
 - Raining



Conditional Independence

- What about this domain:
 - Fire
 - Smoke
 - Alarm



Conditional Independence and the Chain Rule

- Chain rule: $P(X_1, X_2, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots$

- Trivial decomposition:

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain}, \text{Traffic})$$

- With assumption of conditional independence:

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain})$$

- Bayes'nets / graphical models help us express conditional independence assumptions



Ghostbusters Chain Rule

- Each sensor depends only on where the ghost is
- That means, the two sensors are conditionally independent, given the ghost position
- T: Top square is red
B: Bottom square is red
G: Ghost is in the top

- Givens:

$$P(+g) = 0.5$$


$$P(-g) = 0.5$$

$$P(+t \mid +g) = 0.8$$

$$P(+t \mid -g) = 0.4$$

$$P(+b \mid +g) = 0.4$$

$$P(+b \mid -g) = 0.8$$




$$P(T,B,G) = P(G) P(T \mid G) P(B \mid G)$$

T	B	G	P(T,B,G)
+t	+b	+g	0.16
+t	+b	-g	0.16
+t	-b	+g	0.24
+t	-b	-g	0.04
-t	+b	+g	0.04
-t	+b	-g	0.24
-t	-b	+g	0.06
-t	-b	-g	0.06




Bayes' Nets: Big Picture




Bayes' Nets: Big Picture

- Two problems with using full joint distribution tables as our probabilistic models:
 - Unless there are only a few variables, the joint is WAY too big to represent explicitly
 - Hard to learn (estimate) anything empirically about more than a few variables at a time
- Bayes' nets:** a technique for describing complex joint distributions (models) using simple, local distributions (conditional probabilities)
 - More properly called **graphical models**
 - We describe how variables locally interact
 - Local interactions chain together to give global, indirect interactions
 - For about 10 min, we'll be vague about how these interactions are specified



Example Bayes' Net: Insurance



Example Bayes' Net: Car

