

FRACTURE DETECTED

69.1%


confidence

Diagnosis: Oblique Displaced

Imaging Analysis



Original X-ray



AI Explanation (Grad-CAM)

Diagnosis Details


Severity:

Moderate-Severe

Margin:

43.7% over 2nd class

Class Probabilities



Patient Summary

The AI analysis has detected what appears to be a Oblique Displaced fracture. This is classified as Moderate-Severe (The bone pieces have shifted out of place.).

Treatment Guidelines

- Closed or open reduction typically required
- May need internal fixation (pins, plates)
- Extended immobilization (8-12 weeks)
- Regular imaging to monitor alignment

Model Ensemble Breakdown

Model	Prediction	Confidence
Maxvit	Oblique Displaced	99.9%
Hypercolumn Cbam Densenet169	Oblique Displaced	99.4%
Rad Dino	Oblique Displaced	77.2%
Yolo	Oblique	99.7%

Gemini AI Explanation

Diagnosis: Oblique Displaced • Powered by Gemini

I understand you've received a diagnosis of an "Oblique Displaced" fracture. Let's break down what that means in a way that's easy to understand.

Oblique Displaced Fracture:

- * Oblique: This means the fracture line runs at an angle across the bone, rather than straight across it. Imagine cutting a piece of wood diagonally instead of straight across.
- * Displaced: This indicates that the broken pieces of the bone have shifted out of their normal position. They are no longer perfectly aligned.

The confidence level for this diagnosis is 0.69, which means the system is reasonably confident in this assessment.

This type of fracture is currently rated as moderate to severe.

Regarding Treatment and Prognosis:

The details suggest that a procedure called reduction is typically needed. This means a doctor will work to put the bone fragments back into their proper alignment.

- * This reduction can be done closed, meaning without surgery, or open, which involves surgery.
- * In some cases, internal fixation might be necessary. This involves using devices like pins or plates to hold the bone pieces together while they heal.
- * You can expect a period of extended immobilization, meaning the bone will need to be kept still for about 8 to 12 weeks.
- * Throughout this time, regular imaging (like X-rays) will be done to make sure the bone fragments are staying in the correct position as they heal.

The long-term outlook is generally good, especially with surgical correction, and the typical healing time is 8 to 12 weeks.

Please remember, this information is based on the diagnostic details provided. It's important to discuss these findings and the specific treatment plan with your doctor, as they can provide personalized care and answer all your questions.