

Examining Treatment Strategies for Cholera Incorporating Spatial Dynamics

Group: Plague Doctors

Jessa Mallare, Sid Reed, Daniel Segura, Aref Jadda

McMaster University


Instructor: Dr. David Earn

April 8, 2019

- Treatments have not always gone as planned in history
- Cholera


Some Biology on Cholera

- *Vibrio cholerae*
- Colonize the small intestines
- 10 percent of infected individuals develop symptoms
- Causes severe dehydration



Outbreaks in London (19th Century)

- 1832, 1849, 1854, 1866
- Miasma Theory
- John Snow



John Snow

Developing a Single-Patch Model

- Entire population (N) included
- 3 Compartments : S, I, R
- Compartment values are proportional
- Environment (Water)

SIRW Model Assumptions

- Birth Rate = Natural Death Rate and is constant
- Homogenous susceptibility to cholera across population
- No waning immunity
- No latency period
- Only infected individuals can infect the water sources
- Water source is still

SIRW Model



- Using the method of Next Generation Matrix (van den Driessche and Watmough, 2002)

$$F = \begin{pmatrix} \beta_I & \beta_w \\ 0 & 0 \end{pmatrix}$$
$$V = \begin{pmatrix} \frac{1}{\gamma + \mu} & 0 \\ \frac{1}{\gamma + \mu} & \frac{1}{\sigma} \end{pmatrix}$$

- R_0 is computed as the spectral radius of FV^{-1} :

$$\begin{aligned} \mathcal{R}_0 &= \rho(FV^{-1}) \\ &= \frac{\beta_I + \beta_w}{\gamma + \mu} \approx 1.1 - 2.7 \end{aligned}$$

- Two equilibria:


① DFE: $(S, I, R) = (1, 0, 0)$

② EE: $(S^*, I^*, R^*) = (\frac{1}{R_0}, \frac{\mu}{\gamma + \mu}(1 - S^*), I^*)$


- The DFE is stable when $\mathcal{R}_0 < 1$.
- The EE is globally stable when $\mathcal{R}_0 > 1$ (Tien and Earn, 2010).

SIWR Model Phase Portrait

Base Model Without Vital Dynamics $\mathcal{R}_0 = 2.4$




Base Model With Vital Dynamics $\mathcal{R}_0 = 2.25$




Final Size

- Assuming $\mu = 0$ and $\mathcal{R}_0 > 1$, final size formula* still holds:
- $Z = 1 - \exp(-\mathcal{R}_0 Z - \frac{\beta_w}{\sigma} w_0)$ (Tien and Earn, 2010)




Effects of the 19th Century Treatments

- Added parameter death caused by cholera (α)



Effects of the 19th Century Treatments



- Estimated death rate of cholera in the 19th century ranges from a small percentage up to fifty percent
- Including disease induced death is “beneficial” if cholera death rate is high (Why?)
- Improper sanitation increases peak prevalence

Multi-Patch Model

$$\frac{dS_i}{dt} = \mu N - \mu S_i - \beta_i S_i I_i - \phi \beta_i S_i \sum_j^n I_j - \beta_w S_i W_i - \psi \beta_w S_i \sum_j^n W_j$$

$$\frac{dI_i}{dt} = \beta_i S_i I_i + \beta_i \phi S_i \sum_j^n I_j + \beta_w S_i W_i + \beta_i \psi S_i \sum_j^n W_j - I_i(\gamma + \mu + \alpha)$$

$$\frac{dR_i}{dt} = \gamma I_i - \mu R_i$$

$$\frac{dW_i}{dt} = \xi I_i + \beta_i \psi I_i \sum_j^n W_j - \sigma W_i$$


Multi-Patch Model Assumptions

- No dispersal of individuals
- Infected individuals can infect the susceptible in neighboring patches
- All patches neighbouring i have the same transmission rate to patch i


Treatment Strategies For Cholera

- 1 Sanitation of Water
- 2 Vaccinations
- 3 Antibiotics


Sanitation of Water




Vaccinations



Antibiotics



Comparing the Treatment Strategies



- Parameters chosen from literature
- Lowest peak prevalence for antibiotic treatment

Conclusions and Further Research

- 19th century outbreaks
- Different Treatments have different costs and outcomes
- Significance of the using multi-patch model
- Further research on the spread of water borne diseases

Thank you!

References I

Paneth N, Vinten-Johansen P, Brody H, Rip M.

A rivalry of foulness: official and unofficial investigations of the London cholera epidemic of 1854.

Am J Public Health. 1998;88(10):1545–1553.

Tien JH, Earn DJ.

Multiple transmission pathways and disease dynamics in a waterborne pathogen model.

Bull Math Biol. 2010;72(6):1506–1533.

Codeco CT.

Endemic and epidemic dynamics of cholera: the role of the aquatic reservoir.

BMC Infect Dis. 2001;1:1.

References II

Chao DL, Longini IM, Morris JG.

Modeling cholera outbreaks.

Curr Top Microbiol Immunol. 2014;379:195–209.

van den Driessche P.

In: Brauer F, van den Driessche P, Wu J, editors. Spatial Structure: Patch Models. Berlin, Heidelberg: Springer Berlin Heidelberg; 2008. p. 179–189.

Tien JH, Poinar HN, Fisman DN, Earn DJ.

Herald waves of cholera in nineteenth century London.

J R Soc Interface. 2011;8(58):756–760.

Tuite AR, Tien J, Eisenberg M, Earn DJ, Ma J, Fisman DN.

Cholera epidemic in Haiti, 2010: using a transmission model to explain spatial spread of disease and identify optimal control interventions.

Ann Intern Med. 2011;154(9):593–601.

References III

Grad YH, Miller JC, Lipsitch M.

Cholera modeling: challenges to quantitative analysis and predicting the impact of interventions.

Epidemiology. 2012;23(4):523–530.

Lee EC, Kelly MR, Ochocki BM, Akinwumi SM, Hamre KES, Tien JH, et al.

Model distinguishability and inference robustness in mechanisms of cholera transmission and loss of immunity.

J Theor Biol. 2017;420:68–81.

Luo J, Wang J, Wang H.

Seasonal forcing and exponential threshold incidence in cholera dynamics.

Discrete and Continuous Dynamical Systems. 2017;22(6):2261–2290.

References IV

Riley S, Eames K, Isham V, Mollison D, Trapman P.

Five challenges for spatial epidemic models.

Epidemics. 2015;10:68–71.

WHO. Cholera; 2019.

Available from:

<https://www.who.int/news-room/fact-sheets/detail/cholera>.

Moe CL, Rheingans RD.

Global challenges in water, sanitation and health.

J Water Health. 2006;4 Suppl 1:41–57.

Graves PM, Deeks JJ, Demicheli V, Jefferson T.

Vaccines for preventing cholera: killed whole cell or other subunit vaccines (injected).

Chochrane Database Syst Rev. 2010;.