Examining Treatment Strategies for Cholera Incorporating Spatial Dynamics

Group: Plague Doctors Jessa Mallare, Sid Reed, Daniel Segura, Aref Jadda

McMaster University

Instructor: Dr. David Earn

April 8, 2019


Introduction

• Treatments have not always gone as planned in history

Cholera

Some Biology on Cholera

- Vibrio cholerae
- Colonizes the small intestine
- 10 percent of infected individuals develop symptoms
- Causes severe dehydration



Outbreaks in London (19^{th} Century)

- 1832, 1849, 1854, 1866
- Miasma Theory
- John Snow




Developing a Single-Patch Model

- Entire population (N) included
- 3 Compartments : S, I, R
- Compartment values are proportional
- Water compartment for Cholera

SIRW Model Assumptions

- Birth Rate = Natural Death Rate and is constant
- Population equally succeptible to infection
- No waning immunity
- No latency period
- Only infected individuals can infect the water sources
- Water sources can infect individuals

SIRW Model



R_0 Calculation

 Using the method of Next Generation Matrix (van den Driessche and Watmough, 2002)

$$F = \begin{pmatrix} \beta_i & \beta_w \\ 0 & 0 \end{pmatrix}$$
$$V = \begin{pmatrix} \frac{1}{\gamma + \mu} & 0 \\ \frac{1}{\gamma + \mu} & \frac{1}{\sigma} \end{pmatrix}$$


• R_0 is computed as the spectral radius of FV^{-1} :


$$\mathcal{R}_0 = \rho(FV^{-1})$$
$$= \frac{\beta_i + \beta_w}{\gamma + \mu} \approx 1.1 - 2.7$$

Equilibria and Stability

- Two equilibria:
 - **1 DFE**: (S, I, R) = (1, 0, 0)
 - **2** EE: $(S^*, I^*, R^*) = (\frac{1}{R_0}, \frac{\mu}{\gamma + \mu}(1 S^*), I^*)$
- The DFE is stable when $\mathcal{R}_0 < 1$.
- ullet The EE is globally stable when $\mathcal{R}_0>1$ (Tien and Earn, 2010).

SIWR Model Phase Portrait






Final Size


• Assuming $\mu = 0$ and $\mathcal{R}_0 > 1$, final size formula* still holds:

•
$$Z = 1 - \exp\left(-\mathcal{R}_0 Z - \frac{\beta_w}{\sigma} w_0\right)$$




Effects of the 19th Century Treatments

• Added parameter for death rate from Cholera: (α)



Effects of the 19th Century Treatments




- Estimated death rate from cholera in the 19th century estimated to be up to 50%
- Including disease induced death is "beneficial" if death rate by cholera is high (Why?)
- Improper sanitation increases peak prevalence


Treatment Strategies For Cholera

- Sanitation of Water
- Vaccinations
- Antibiotics


Sanitation of Water




Vaccinations



Antibiotics



Comparing the Treatment Strategies



- Parameters chosen from literature
- Lowest peak prevalence for vaccination treatment

Multi-Patch Model

$$\frac{dS_i}{dt} = \mu N - \mu S_i - \beta_i S_i I_i - \beta_w S_i W_i + r(S)$$

$$\frac{dI_i}{dt} = \beta_i S_i I_i + \beta_w S_i W_i - I_i (\gamma + \mu + \alpha)$$

$$\frac{dR_i}{dt} = \gamma I_i - \mu R_i + r(R)$$

$$\frac{dW_i}{dt} = \xi I_i - \sigma W_i + r(W)$$


Multi-Patch Model Assumptions

- No dispersal of infected individuals
- All patches have the same set of parameter values

Spatial Dynamics (Base)

Base Sanitation Antibiotics Vaccination

Comparing the Treatment Strategies In Multipatch Model



Conclusions and Further Research

- 19th century outbreaks poorly handeled without antibiotics/vaccines
- Different Treatments have different costs and outcomes
- Further research on improving water sanitation effectivness to reduce reliance on vaccines/antibiotics

Thank you!

References I

- Paneth, N., Vinten-Johansen, P., Brody, H. & Rip, M. A rivalry of foulness: official and unofficial investigations of the London cholera epidemic of 1854. *Am J Public Health* **88**, 1545–1553 (1998).
- Chao, D. L., Longini, I. M. & Morris, J. G. Modeling cholera outbreaks. *Curr. Top. Microbiol. Immunol.* **379**, 195–209 (2014).
- Tien, J. H., Poinar, H. N., Fisman, D. N. & Earn, D. J. Herald waves of cholera in nineteenth century London. *J R Soc Interface* **8**, 756–760 (2011).
- Tuite, A. R. *et al.* Cholera epidemic in Haiti, 2010: using a transmission model to explain spatial spread of disease and identify optimal control interventions. *Ann. Intern. Med.* **154**, 593–601 (2011).

References II

- Grad, Y. H., Miller, J. C. & Lipsitch, M. Cholera modeling: challenges to quantitative analysis and predicting the impact of interventions. *Epidemiology* **23**, 523–530 (2012).
- Wilford, J. N. How epidemics helped shape the modern metropolis. New York Times (2008).
- Sciarra, C., Rinaldo, A., Laio, F. & Pasetto, D. Mathematical modeling of cholera epidemics in South Sudan. *arXiv e-prints*. arXiv: 1801.03125 [q-bio.PE] (2018).
- Codeco, C. T. Endemic and epidemic dynamics of cholera: the role of the aquatic reservoir. *BMC Infect. Dis.* **1,** 1 (2001).
- Tien, J. H. & Earn, D. J. Multiple transmission pathways and disease dynamics in a waterborne pathogen model. *Bull. Math. Biol.* **72**, 1506–1533 (2010).

References III

- Van den Driessche, P. in *Mathematical Epidemiology* (eds Brauer, F., van den Driessche, P. & Wu, J.) 179–189 (Springer Berlin Heidelberg, Berlin, Heidelberg, 2008).
- Kaper, J. B., Morris, J. G. & Levine, M. M. Cholera. *Clinical Microbiology Reviews* **8**, 48–86 (1995).
- Li, P. cholera: Amend, Augment and Aid Analysis of John Snow's Cholera Map. (2019).
- Lee, E. C. *et al.* Model distinguishability and inference robustness in mechanisms of cholera transmission and loss of immunity. *J. Theor. Biol.* **420**, 68–81 (2017).
- Azman, A. S., Rudolph, K. E., Cummings, D. A. & Lessler, J. The incubation period of cholera: a systematic review. *J. Infect.* **66**, 432–438 (2013).

References IV

- Kong, J. D., Davis, W. & Wang, H. Dynamics of a cholera transmission model with immunological threshold and natural phage control in reservoir. *Bull. Math. Biol.* **76**, 2025–2051 (2014).
- Luo, J., Wang, J. & Wang, H. Seasonal forcing and exponential threshold incidence in cholera dynamics. *Discrete and Continuous Dynamical Systems* **22**, 2261–2290 (2017).
- Riley, S., Eames, K., Isham, V., Mollison, D. & Trapman, P. Five challenges for spatial epidemic models. *Epidemics* **10**, 68–71 (2015).
- WHO. Cholera. https://www.who.int/news-room/fact-sheets/detail/cholera (2019).
- Moe, C. L. & Rheingans, R. D. Global challenges in water, sanitation and health. *J Water Health* **4 Suppl 1**, 41–57 (2006).
- Earn, D. J., Andrews, P. W. & Bolker, B. M. Population-level effects of suppressing fever. *Proc. Biol. Sci.* **281**, 20132570 (Mar. 2014).

References V

- Mukerjee, S. Preliminary studies on the development of a live oral vaccine for anti-cholera immunizaion. *Bull. World Health Organ.* **29**, 753–766 (1963).
- Graves, P., Deeks, J., Demicheli, V. & Jefferson, T. Vaccines for preventing cholera: killed whole cell or other subunit vaccines (injected). *Chochrane Database Syst. Rev.* (2010).
- Nelson, E. J., Nelson, D. S., Salam, M. A. & Sack, D. A. Antibiotics for both moderate and severe cholera. *N. Engl. J. Med.* **364**, 5–7 (2011).
- Saha, D., Karim, M. M., Khan, W. & Ahmed, S. Single-dose azithromycin for the treatment of cholera in adults. *N. Engl. J. Med.* **364**, 2452–2462 (2006).

References VI



Fung, I. C., Fitter, D. L., Borse, R. H., Meltzer, M. I. & Tappero, J. W. Modelling the effect of water, sanitation, and hygiene and oral cholera vaccine implementation in Haiti. *Am. J. Trop. Med. Hyg.* **89**, 633–640 (2013).



Taylor, D., Kahawita, T., Cairncross, S. & Ensink, J. H. J. The Impact of Water, Sanitation and Hygiene Interventions to Control Cholera: A Systematic Review. *PLoS ONE* **10** (2015).