

# Is Sharing Caring?

Elucidating the Effects of the  
Presence of CRISPR-Cas Systems  
on Rates of Horizontal Gene  
Transfer Using Network Analysis

Siddharth Reed  
MolBiol 4C12 Thesis



Golding Lab,  
Biology Department,  
McMaster University

March 31, 2019

# Table of Contents

---

1. CRISPR-Cas systems
2. Horizontal Gene Transfer
3. Phylogenomic Networks
4. Do CRRISPR Systems Affect Horizontal Gene Transfer?
5. My Project
6. Results


# CRISPR-Cas systems

# What Are They?

---


# What Are They?

- Adaptive Bacterial Immune System




# What Are They?

- Adaptive Bacterial Immune System
- Protects against foreign DNA



# What Are They?

- Adaptive Bacterial Immune System
- Protects against foreign DNA
- Requires Cas proteins and CRISPR loci




# Diversity & Ubiquity




# Diversity & Ubiquity

- 45% of bacteria have CRISPR loci ( $n = 6782$ )<sup>2</sup>




# Diversity & Ubiquity

- 45% of bacteria have CRISPR loci ( $n = 6782$ )<sup>2</sup>
- 3 Main Types, multiple subtypes<sup>3</sup>




# Diversity & Ubiquity

- 45% of bacteria have CRISPR loci ( $n = 6782$ )<sup>2</sup>
- 3 Main Types, multiple subtypes<sup>3</sup>
- CRISPR arrays represent unique life history of an organism




# Diversity & Ubiquity

- 45% of bacteria have CRISPR loci ( $n = 6782$ )<sup>2</sup>
- 3 Main Types, multiple subtypes<sup>3</sup>
- CRISPR arrays represent unique life history of an organism
- 11% – 28% are false or orphaned CRISPR loci<sup>4</sup>




# Biotech Application

# Biotech Application




# Horizontal Gene Transfer

# Mechanisms






# Mechanisms




- Conjugation: Transfer of DNA through cell-cell connections<sup>6</sup>

# Mechanisms




- Conjugation: Transfer of DNA through cell-cell connections<sup>6</sup>
- Transformation: Incorporation of free-floating DNA into the genome<sup>6</sup>

# Mechanisms



- Conjugation: Transfer of DNA through cell-cell connections<sup>6</sup>
- Transformation: Incorporation of free-floating DNA into the genome<sup>6</sup>
- Transduction: Transfer of DNA through phage<sup>6</sup>

# Mechanisms




- Conjugation: Transfer of DNA through cell-cell connections<sup>6</sup>
- Transformation: Incorporation of free-floating DNA into the genome<sup>6</sup>
- Transduction: Transfer of DNA through phage<sup>6</sup>
- **CRISPR-Cas directly affects Transduction and Transformation<sup>6</sup>**


# Pan-Genomes

---


# Pan-Genomes



# Pan-Genomes



7



8

# Rate Influencing Factors

---



# Rate Influencing Factors

---

- Amount of exogenous DNA/cell density/phage density

# Rate Influencing Factors

---

- Amount of exogenous DNA/cell density/phage density
- Selective pressures

# Rate Influencing Factors

---

- Amount of exogenous DNA/cell density/phage density
- Selective pressures
- Metabolic costs

# Rate Influencing Factors


---

- Amount of exogenous DNA/cell density/phage density
- Selective pressures
- Metabolic costs
- Sequence compatibility

# Applications

---

# Applications




# Phylogenomic Networks

# What is A Network?

---




# What is A Network?




- Useful mathematical abstraction of real world system

# What is A Network?




- Useful mathematical abstraction of real world system
- Nodes can have attributes

# What is A Network?



- Useful mathematical abstraction of real world system
- Nodes can have attributes
- Directed or Undirected Edges

# What is A Network?




- Useful mathematical abstraction of real world system
- Nodes can have attributes
- Directed or Undirected Edges
- Weighted or Unweighted Edges

# Prokaryotic “Net of Life”


---

# Prokaryotic “Net of Life”




# Construction


## 1. Parametric methods




## 2. Phylogenetic methods



### 2a. Implicit phylogenetic methods



### 2b. Explicit phylogenetic methods



# Do CRRISPR Systems Affect Horizontal Gene Transfer?



Yes

# CRISPR Cost Complexity

# CRISPR Cost Complexity

---

- Cost tradeoff factors:

# CRISPR Cost Complexity

---

- Cost tradeoff factors:
  - Metabolic maintenance<sup>1</sup>

# CRISPR Cost Complexity

---

- Cost tradeoff factors:
  - Metabolic maintenance<sup>1</sup>
  - Environmental pressures<sup>13</sup>

# CRISPR Cost Complexity

---

- Cost tradeoff factors:
  - Metabolic maintenance<sup>1</sup>
  - Environmental pressures<sup>13</sup>
  - Off-target effects (autoimmune)<sup>14</sup>

# CRISPR Cost Complexity

---

- Cost tradeoff factors:
  - Metabolic maintenance<sup>1</sup>
  - Environmental pressures<sup>13</sup>
  - Off-target effects (autoimmune)<sup>14</sup>
  - Anti-CRISPR systems<sup>3</sup>

# CRISPR Cost Complexity

---

- Cost tradeoff factors:
  - Metabolic maintenance<sup>1</sup>
  - Environmental pressures<sup>13</sup>
  - Off-target effects (autoimmune)<sup>14</sup>
  - Anti-CRISPR systems<sup>3</sup>
  - Phage virulence/density<sup>3</sup>



# CRISPR Cost Complexity

---

- Cost tradeoff factors:
  - Metabolic maintenance<sup>1</sup>
  - Environmental pressures<sup>13</sup>
  - Off-target effects (autoimmune)<sup>14</sup>
  - Anti-CRISPR systems<sup>3</sup>
  - Phage virulence/density<sup>3</sup>
  - Prophage abundance<sup>15</sup>

# Curbing CRISPR Cost

## Curbing CRISPR Cost

- CRISPRs themselves can be transferred  $\implies$  population level immunity<sup>16</sup>

## Curbing CRISPR Cost

---

- CRISPRs themselves can be transferred  $\implies$  population level immunity<sup>16</sup>
- Selective CRISPR inactivation<sup>1</sup>

## Curbing CRISPR Cost

---

- CRISPRs themselves can be transferred  $\implies$  population level immunity<sup>16</sup>
- Selective CRISPR inactivation<sup>1</sup>
- CRISPR can enhance transduction-mediated HGT<sup>15</sup>

## Previous Findings

---

## Previous Findings

---

- Gophna et al. (2015) found no relation between the presence of CRISPR systems and HGT over short evolutionary timescales<sup>17</sup>

## Previous Findings

---

- Gophna et al. (2015) found no relation between the presence of CRISPR systems and HGT over short evolutionary timescales<sup>17</sup>
  - Assume all singletons arose from HGT



## Previous Findings

---

- Gophna et al. (2015) found no relation between the presence of CRISPR systems and HGT over short evolutionary timescales<sup>17</sup>
  - Assume all singletons arose from HGT
  - Used GC% to identify HGT

## Previous Findings

---

- Gophna et al. (2015) found no relation between the presence of CRISPR systems and HGT over short evolutionary timescales<sup>17</sup>
  - Assume all singletons arose from HGT
  - Used GC% to identify HGT
- Contradicted by a former undergraduate thesis student

## Previous Findings

---

- Gophna et al. (2015) found no relation between the presence of CRISPR systems and HGT over short evolutionary timescales<sup>17</sup>
  - Assume all singletons arose from HGT
  - Used GC% to identify HGT
- Contradicted by a former undergraduate thesis student
  - Can see inhibitory effects of CRISPR on HGT over short evolutionary time scales

## Previous Findings

---

- Gophna et al. (2015) found no relation between the presence of CRISPR systems and HGT over short evolutionary timescales<sup>17</sup>
  - Assume all singletons arose from HGT
  - Used GC% to identify HGT
- Contradicted by a former undergraduate thesis student
  - Can see inhibitory effects of CRISPR on HGT over short evolutionary time scales
  - Higher gene indel rates for CRISPR containing genera than non-CRISPR containing outgroups

# My Project

# Hypothesis

---

## Null Hypothesis

Bacterial strains or genera with known CRISPR systems will show no significant differences in network statistics compared to those strains or genera without known CRISPR systems.

# Hypothesis

---

## Null Hypothesis

Bacterial strains or genera with known CRISPR systems will show no significant differences in network statistics compared to those strains or genera without known CRISPR systems

## Alternative Hypothesis

Bacterial strains or genera with known CRISPR systems will show a significant difference in at least 1 network statistic compared to those strains or genera without known CRISPR systems.

# Objectives

---



# Objectives

---

## Within Network Comparisons

For genera with CRISPR containing strains, compare the node statistics of CRISPR-containing strain to non-CRISPR-containing strains.

# Objectives

---

## Within Network Comparisons

For genera with CRISPR containing strains, compare the node statistics of CRISPR-containing strain to non-CRISPR-containing strains.

## Gene Indel Rates vs. Network Statistics

Compare gene InDel rates to node/network statistics for CRISPR-containing and non-CRISPR-containing strains/genera.

# Workflow


---



Get Fastas


# Workflow

---




# Workflow

---




# Workflow


---



# Workflow




# Workflow







# Workflow




# Workflow




# Workflow




# Workflow



# Workflow



# Workflow



# Network Statistics

---

# Network Statistics

---

- **Average Node Degree:**  $\frac{1}{|N_u|} \sum_{uv} w_{uv}$  where  $N_u$  is the set of nodes incident to  $u$



## Network Statistics

---

- **Average Node Degree:**  $\frac{1}{|N_u|} \sum_{uv} w_{uv}$  where  $N_u$  is the set of nodes incident to  $u$
- **Average Edge Weight:**  $\frac{1}{N_c} \sum_i w_i$ , The average edge weight for all nodes with CRISPR or without CRISPR

# Network Statistics

---

- **Average Node Degree:**  $\frac{1}{|N_u|} \sum_{uv}^{N_u} w_{uv}$  where  $N_u$  is the set of nodes incident to  $u$
- **Average Edge Weight:**  $\frac{1}{N_c} \sum_i w_i$ , The average edge weight for all nodes with CRISPR or without CRISPR
- **Node Clustering Coefficient:**  $\frac{1}{k_u(k_u-1)} \sum_{vw}^{T(u)} (\hat{w}_{uw} \hat{w}_{vw} \hat{w}_{uv})^{\frac{1}{3}}$   
where  $T(u)$  is the set of triangles containing  $u$ <sup>18</sup>

## Network Statistics

- **Average Node Degree:**  $\frac{1}{|N_u|} \sum_{uv} w_{uv}$  where  $N_u$  is the set of nodes incident to  $u$
- **Average Edge Weight:**  $\frac{1}{N_c} \sum_i w_i$ , The average edge weight for all nodes with CRISPR or without CRISPR
- **Node Clustering Coefficient:**  $\frac{1}{k_u(k_u-1)} \sum_{vw}^{T(u)} (\hat{w}_{uw} \hat{w}_{vw} \hat{w}_{uv})^{\frac{1}{3}}$   
where  $T(u)$  is the set of triangles containing  $u$ <sup>18</sup>
- **Node Assortativity:**  $\frac{Tr(M) - ||M^2||}{1 - ||M^2||}$  Where  $M$  is the mixing matrix of a given attribute and  $||M||$  is the sum of all elements of  $M$ .<sup>19</sup>

## Network Statistics

- **Average Node Degree:**  $\frac{1}{|N_u|} \sum_{uv} w_{uv}$  where  $N_u$  is the set of nodes incident to  $u$
- **Average Edge Weight:**  $\frac{1}{N_c} \sum_i w_i$ , The average edge weight for all nodes with CRISPR or without CRISPR
- **Node Clustering Coefficient:**  $\frac{1}{k_u(k_u-1)} \sum_{vw}^{T(u)} (\hat{w}_{uw} \hat{w}_{vw} \hat{w}_{uv})^{\frac{1}{3}}$   
where  $T(u)$  is the set of triangles containing  $u$ <sup>18</sup>
- **Node Assortativity:**  $\frac{Tr(M) - ||M^2||}{1 - ||M^2||}$  Where  $M$  is the mixing matrix of a given attribute and  $||M||$  is the sum of all elements of  $M$ .<sup>19</sup>
- **Network Modularity:**  $Q = \frac{1}{2m} \sum_{uv} [W_{uv} - \frac{k_u k_v}{2m}] \delta(u, v)$  where  $m$  is the total weight of all edges,  $k_u$  is the degree of  $u$  and  $\delta(u, v)$  is 1 if  $u$  and  $v$  both have or do not have CRISPR systems and 0 otherwise.  $Q \in [-1, 1]$ <sup>20</sup>

# Results

# Limitations & Caveats

---

## Limitations & Caveats

---

- **Ignored Singletons:** Genes that did not cluster into any families were ignored from future steps, but may have still represented horizontally transferred genes

## Limitations & Caveats

---

- **Ignored Singletons:** Genes that did not cluster into any families were ignored from future steps, but may have still represented horizontally transferred genes
- **Ignored Some Gene Families:** For time considerations, only 1500 gene trees were generated for each genus



## Limitations & Caveats

---

- **Ignored Singletons:** Genes that did not cluster into any families were ignored from future steps, but may have still represented horizontally transferred genes
- **Ignored Some Gene Families:** For time considerations, only 1500 gene trees were generated for each genus
- **Significance Testing:** Samples are not necessarily independent in a network, further node statistics can only be tested for genera with  $> 20$  CRISPR and non-CRISPR OTUs.

## Limitations & Caveats

---

- **Ignored Singletons:** Genes that did not cluster into any families were ignored from future steps, but may have still represented horizontally transferred genes
- **Ignored Some Gene Families:** For time considerations, only 1500 gene trees were generated for each genus
- **Significance Testing:** Samples are not necessarily independent in a network, further node statistics can only be tested for genera with  $> 20$  CRISPR and non-CRISPR OTUs.
- **Taxonomic Mistakes:** Inconsistencies in taxonomic labelling can result in ignored or misplaced OTUs.

## Limitations & Caveats

---

- **Ignored Singletons:** Genes that did not cluster into any families were ignored from future steps, but may have still represented horizontally transferred genes
- **Ignored Some Gene Families:** For time considerations, only 1500 gene trees were generated for each genus
- **Significance Testing:** Samples are not necessarily independent in a network, further node statistics can only be tested for genera with  $> 20$  CRISPR and non-CRISPR OTUs.
- **Taxonomic Mistakes:** Inconsistencies in taxonomic labelling can result in ignored or misplaced OTUs.
- **Multifurcation Error:** Some species trees contained multifurcations, which were resolved randomly to generate a bifurcating tree. Estimating this error by examining variance over different resolutions is possible.

# Possible Future Directions

---

## Possible Future Directions

---

- **Inferring direction:** Directed networks have a host of available analytic tools undirected networks do not

## Possible Future Directions

---

- **Inferring direction:** Directed networks have a host of available analytic tools undirected networks do not
- **Gene function analysis:** Considering the transfer dynamics of different functional classes of genes

## Possible Future Directions

---

- **Inferring direction:** Directed networks have a host of available analytic tools undirected networks do not
- **Gene function analysis:** Considering the transfer dynamics of different functional classes of genes
- **Studying movement of CRISPR systems:** Studying how frequently CRISPR systems themselves are transferred from arrays, cas genes

## Possible Future Directions

---

- **Inferring direction:** Directed networks have a host of available analytic tools undirected networks do not
- **Gene function analysis:** Considering the transfer dynamics of different functional classes of genes
- **Studying movement of CRISPR systems:** Studying how frequently CRISPR systems themselves are transferred from arrays, cas genes
- **Intergenic comparisons:** Combine any set of fasta files for OTUs for analyzing transfer dynamics



## Possible Future Directions

---

- **Inferring direction:** Directed networks have a host of available analytic tools undirected networks do not
- **Gene function analysis:** Considering the transfer dynamics of different functional classes of genes
- **Studying movement of CRISPR systems:** Studying how frequently CRISPR systems themselves are transferred from arrays, cas genes
- **Intergenic comparisons:** Combine any set of fasta files for OTUs for analyzing transfer dynamics
- **Continuous CRISPR activity:** Labelling nodes by estimated CRISPR activity (array length, transcriptomic data, etc.)

## Possible Future Directions

---

- **Inferring direction:** Directed networks have a host of available analytic tools undirected networks do not
- **Gene function analysis:** Considering the transfer dynamics of different functional classes of genes
- **Studying movement of CRISPR systems:** Studying how frequently CRISPR systems themselves are transferred from arrays, cas genes
- **Intergenic comparisons:** Combine any set of fasta files for OTUs for analyzing transfer dynamics
- **Continuous CRISPR activity:** Labelling nodes by estimated CRISPR activity (array length, transcriptomic data, etc.)
- **Considering bacterial ecology and environments:** Consider geographically close OTUs or differences between networks due to environmental factors

# Conclusion

# Thanks

---

Thank you to





- Dr. G. Brian Golding
- Dr. Ben Evans
- The Golding lab
  - Caitlin Simopoulos
  - Daniella Lato
  - Zachery Dickson
  - Sam Long
  - Geoge Long
  - Lucy Zhang
  - Brianne Laverty
  - Nicole Zhang
- Everyone here for listening



All code used for this project is available at [https://github.com/DJSiddharthVader/thesis\\_SidReed](https://github.com/DJSiddharthVader/thesis_SidReed)




## References (1)

---

-  Devashish Rath et al. "The CRISPR-Cas immune system: Biology, mechanisms and applications". In: *Biochimie* 117 (2015). Special Issue: Regulatory RNAs, pp. 119–128. ISSN: 0300-9084.
-  GRissa, I. and Drevet, C. and Couvin, D. *CRISPRdb*. <http://crispr.i2bc.paris-saclay.fr/>. Online; accessed 22 October 2018. 2017.
-  J. Bondy-Denomy and A. R. Davidson. "To Acquire Or Resist:The Complex Biological Effects Of CRISPR-Cas systems". In: *Trends Microbio.* 22.4 (2014), pp. 218–25.
-  Quan Zhang and Yuzhen Ye. "Not all predicted CRISPR–Cas systems are equal: isolated cas genes and classes of CRISPR like elements". In: *BMC Bioinformatics* 18.1 (Feb. 2017), p. 92. ISSN: 1471-2105.






## References (2)

---

-  K. S. Makarova et al. “Evolution and classification of the CRISPR-Cas systems”. In: *Nat. Rev. Microbiol.* 9.6 (2011), pp. 467–477.
-  Ovidiu Popa and Tal Dagan. “Trends and barriers to lateral gene transfer in prokaryotes”. In: *Current Opinion in Microbiology* 14.5 (2011). Antimicrobials/Genomics, pp. 615–623. ISSN: 1369-5274.
-  L. C. Guimaraes et al. “Inside the Pan-genome - Methods and Software Overview”. In: *Curr. Genomics* 16.4 (2015), pp. 245–252.
-  David A. Rasko et al. “The Pangenome Structure of Escherichia coli: Comparative Genomic Analysis of E. coli Commensal and Pathogenic Isolates”. In: *Journal of Bacteriology* 190.20 (2008), pp. 6881–6893. ISSN: 0021-9193.




## References (3)

---

-  Björn Berglund. “Environmental dissemination of antibiotic resistance genes and correlation to anthropogenic contamination with antibiotics”. In: *Infection Ecology & Epidemiology* 5.1 (2015), p. 28564.
-  J. A. Bondy and U. S. R. Murty. *Graph theory with applications*. Wiley, 2002.
-  V. Kunin et al. “The net of life: reconstructing the microbial phylogenetic network”. In: *Genome Res.* 15.7 (2005), pp. 954–959.
-  Matt Ravenhall et al. “Inferring Horizontal Gene Transfer”. In: *PLOS Computational Biology* 11.5 (May 2015), pp. 1–16.
-  Senka Dzidic and Vladimir Bedeković. “Horizontal gene transfer-emerging multidrug resistance in hospital bacteria”. In: *Acta pharmacologica Sinica* 24.6 (2003), pp. 519–526.

## References (4)





---

-  Adi Stern et al. “Self-targeting by CRISPR: gene regulation or autoimmunity?” In: *Trends in Genetics* 26.8 (2010), pp. 335–340. ISSN: 0168-9525.
-  Bridget N. J. Watson, Raymond H. J. Staals, and Peter C. Fineran. “CRISPR-Cas-Mediated Phage Resistance Enhances Horizontal Gene Transfer by Transduction”. In: *mBio* 9.1 (2018). Ed. by Joseph Bondy-Denomy and Michael S. Gilmore.
-  James S. Godde and Amanda Bickerton. “The Repetitive DNA Elements Called CRISPRs and Their Associated Genes: Evidence of Horizontal Transfer Among Prokaryotes”. In: *Journal of Molecular Evolution* 62.6 (June 2006), pp. 718–729. ISSN: 1432-1432.



## References (5)

---

-  U. Gophna et al. “No evidence of inhibition of horizontal gene transfer by CRISPR-Cas on evolutionary timescales”. In: *ISME J* 9.9 (2015), pp. 2021–2027.
-  J. P. Onnela et al. “Intensity and coherence of motifs in weighted complex networks”. In: *Phys Rev E Stat Nonlin Soft Matter Phys* 71.6 Pt 2 (2005), p. 065103.
-  M. E. Newman. “Assortative mixing in networks”. In: *Phys. Rev. Lett.* 89.20 (2002), p. 208701.
-  M. E. Newman. “Analysis of weighted networks”. In: *Phys Rev E Stat Nonlin Soft Matter Phys* 70.5 Pt 2 (2004), p. 056131.