

CS 344: Design & Analysis of Algorithms

Lecture 4

Sep 12, 2019


Maximum subarray problem

Given an array, find a non-empty contiguous subarray that maximizes its sum:


	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
A	13	-3	-25	20	-3	-16	-23	18	20	-7	12	-5	-22	15	-4	7

maximum subarray


Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem




Maximum subarray problem



Maximum subarray problem



Maximum subarray problem



Master theorem

Divide and conquer solves a problem of size n by:

- splitting it into a subproblems of size n/b
- combining the answers in $O(n^d)$ time

where $a, b, d > 0$

Master theorem

If $T(n) = aT(n/b) + O(n^d)$ and $a > 0, b > 1, d \geq 0$, then

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log n) & \text{if } d = \log_b a \\ O(n^{\log_b a}) & \text{if } d < \log_b a \end{cases}$$

Maximum subarray problem

```
if  $high == low$   
    return ( $low, high, A[low]$ )  
else  $mid = \lfloor (low + high) / 2 \rfloor$   
    ( $left-low, left-high, left-sum$ ) =  
        FIND-MAXIMUM-SUBARRAY( $A, low, mid$ )  
    ( $right-low, right-high, right-sum$ ) =  
        FIND-MAXIMUM-SUBARRAY( $A, mid + 1, high$ )  
    ( $cross-low, cross-high, cross-sum$ ) =  
        FIND-MAX-CROSSING-SUBARRAY( $A, low, mid, high$ )  
    if  $left-sum \geq right-sum$  and  $left-sum \geq cross-sum$   
        return ( $left-low, left-high, left-sum$ )  
    elseif  $right-sum \geq left-sum$  and  $right-sum \geq cross-sum$   
        return ( $right-low, right-high, right-sum$ )  
    else return ( $cross-low, cross-high, cross-sum$ )
```

Maximum subarray problem

Step	Time
Solve the midpoint case	$O(n)$
Solve the left and right cases	$T(n/2)$ each
Find the max of those three	$O(1)$

Maximum subarray problem

We get the following recurrence:

$$T(n) = 2T(n/2) + O(n)$$

To match $T(n) = aT(n/b) + O(n^d)$:

- $a = 2$
- $b = 2$
- $d = 1$

Maximum subarray problem

If $T(n) = aT(n/b) + O(n^d)$ and $a > 0, b > 1, d \geq 0$, then

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log n) & \text{if } d = \log_b a \\ O(n^{\log_b a}) & \text{if } d < \log_b a \end{cases}$$

- $a = 2$
- $b = 2$
- $d = 1$

$$\log_b a = \log_2 2 = 1 = d$$

Use the second case.

Maximum subarray problem

If $T(n) = aT(n/b) + O(n^d)$ and $a > 0, b > 1, d \geq 0$, then

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log n) & \text{if } d = \log_b a \\ O(n^{\log_b a}) & \text{if } d < \log_b a \end{cases}$$

- $a = 2$
- $b = 2$
- $d = 1$

$$O(n^d \log n) = O(n^1 \log n) = O(n \log n)$$


Matrix multiplication

If A and B are $n \times n$ matrices, we can multiply them to produce a $n \times n$ matrix C :

$$A \cdot B = C$$

then c_{ij} is the dot product of row i of A and column j of B :

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$



Matrix multiplication

Naïve method:

```
for i from 1 to n:  
  for j from 1 to n:  
    c[i][j] = 0.0  
    for k from 1 to n:  
      c[i][j] += a[i][k] * b[k][j]
```

How long does this take?

Matrix multiplication

Can we view this as a divide and conquer algorithm?

Suppose we decompose each matrix into four $n/2 \times n/2$ blocks:

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}, \quad B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}, \quad C = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix}$$

Matrix multiplication

Then we can define multiplication as such:

$$\begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \cdot \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$$

Matrix multiplication

So we have these subproblems to compute:

$$C_{11} = A_{11} \cdot B_{11} + A_{12} \cdot B_{21}$$

$$C_{12} = A_{11} \cdot B_{12} + A_{12} \cdot B_{22}$$

$$C_{21} = A_{21} \cdot B_{11} + A_{22} \cdot B_{21}$$

$$C_{22} = A_{21} \cdot B_{12} + A_{22} \cdot B_{22}$$

Matrix multiplication

$n = A.rows$

let C be a new $n \times n$ matrix

if $n == 1$

$$c_{11} = a_{11} \cdot b_{11}$$

else partition A , B , and C as in equations (4.9)

$$C_{11} = \text{SQUARE-MATRIX-MULTIPLY-RECURSIVE}(A_{11}, B_{11}) \\ + \text{SQUARE-MATRIX-MULTIPLY-RECURSIVE}(A_{12}, B_{21})$$

$$C_{12} = \text{SQUARE-MATRIX-MULTIPLY-RECURSIVE}(A_{11}, B_{12}) \\ + \text{SQUARE-MATRIX-MULTIPLY-RECURSIVE}(A_{12}, B_{22})$$

$$C_{21} = \text{SQUARE-MATRIX-MULTIPLY-RECURSIVE}(A_{21}, B_{11}) \\ + \text{SQUARE-MATRIX-MULTIPLY-RECURSIVE}(A_{22}, B_{21})$$

$$C_{22} = \text{SQUARE-MATRIX-MULTIPLY-RECURSIVE}(A_{21}, B_{12}) \\ + \text{SQUARE-MATRIX-MULTIPLY-RECURSIVE}(A_{22}, B_{22})$$

return C

Matrix multiplication

- Divide into 8 blocks of size $n/2 \times n/2$
- Recursively multiply
- Add (some of) the resulting matrices

Matrix multiplication

Step	Time
Divide into 8 blocks of size $n/2 \times n/2$	$O(1)$
Recursively multiply	$T(n/2)$ each
Add resulting matrices	$O(n^2)$

Matrix multiplication

Then the time required has the form:

$$T(n) = 8T(n/2) + O(n^2)$$

Matrix multiplication

$$T(n) = 8T(n/2) + O(n^2)$$

Master theorem: $T(n) = aT(n/b) + O(n^d)$

- $a = 8$
- $b = 2$
- $d = 2$

Matrix multiplication

If $T(n) = aT(n/b) + O(n^d)$ and $a > 0, b > 1, d \geq 0$, then

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log n) & \text{if } d = \log_b a \\ O(n^{\log_b a}) & \text{if } d < \log_b a \end{cases}$$

- $a = 8$
- $b = 2$
- $d = 2$

$$\log_b a = \log_2 8 = 3 > 2 = d$$

Use case 3.

Matrix multiplication

If $T(n) = aT(n/b) + O(n^d)$ and $a > 0, b > 1, d \geq 0$, then

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log n) & \text{if } d = \log_b a \\ O(n^{\log_b a}) & \text{if } d < \log_b a \end{cases}$$

- $a = 8$
- $b = 2$
- $d = 2$

$$O(n^{\log_b a}) = O(n^3)$$

Matrix multiplication

So matrix multiplication is $O(n^3)$.

Is n^3 also a lower bound?

(Is it also $\Omega(n^3)$, and thus $\Theta(n^3)$?)

Actually, there is a clever rearrangement that can do better!

It was found in 1969 by Volker Strassen, and is known as Strassen's method.

“Strassen’s method is not at all obvious.”

– CLRS

Strassen's method

There are four steps to Strassen's method:

- Divide the matrices into $n/2 \times n/2$ matrices as before
- Add/subtract some of these to create 10 new temporary matrices S_1, \dots, S_{10}
- Compute 7 matrix products P_1, \dots, P_7
- Compute submatrices $C_{11}, C_{12}, C_{21}, C_{22}$

$$\begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \cdot \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$$

Strassen's method

First compute the S matrices:

$$S_1 = B_{12} - B_{22} ,$$

$$S_2 = A_{11} + A_{12} ,$$

$$S_3 = A_{21} + A_{22} ,$$

$$S_4 = B_{21} - B_{11} ,$$

$$S_5 = A_{11} + A_{22} ,$$

$$S_6 = B_{11} + B_{22} ,$$

$$S_7 = A_{12} - A_{22} ,$$

$$S_8 = B_{21} + B_{22} ,$$

$$S_9 = A_{11} - A_{21} ,$$

$$S_{10} = B_{11} + B_{12} .$$

Strassen's method

Then compute the P matrices:

$$P_1 = A_{11} \cdot S_1 = A_{11} \cdot B_{12} - A_{11} \cdot B_{22} ,$$

$$P_2 = S_2 \cdot B_{22} = A_{11} \cdot B_{22} + A_{12} \cdot B_{22} ,$$

$$P_3 = S_3 \cdot B_{11} = A_{21} \cdot B_{11} + A_{22} \cdot B_{11} ,$$

$$P_4 = A_{22} \cdot S_4 = A_{22} \cdot B_{21} - A_{22} \cdot B_{11} ,$$

$$P_5 = S_5 \cdot S_6 = A_{11} \cdot B_{11} + A_{11} \cdot B_{22} + A_{22} \cdot B_{11} + A_{22} \cdot B_{22} ,$$

$$P_6 = S_7 \cdot S_8 = A_{12} \cdot B_{21} + A_{12} \cdot B_{22} - A_{22} \cdot B_{21} - A_{22} \cdot B_{22} ,$$

$$P_7 = S_9 \cdot S_{10} = A_{11} \cdot B_{11} + A_{11} \cdot B_{12} - A_{21} \cdot B_{11} - A_{21} \cdot B_{12} .$$

Finally, combine these to get the C submatrices:

$$C_{11} = P_5 + P_4 - P_2 + P_6$$

$$C_{12} = P_1 + P_2$$

$$C_{21} = P_3 + P_4$$

$$C_{22} = P_1 + P_5 - P_3 - P_7$$

Strassen's method

Expanding C_{11} :

$$\begin{array}{r} A_{11} \cdot B_{11} + A_{11} \cdot B_{22} + A_{22} \cdot B_{11} + A_{22} \cdot B_{22} \\ \quad - A_{22} \cdot B_{11} \quad \quad \quad + A_{22} \cdot B_{21} \\ \quad - A_{11} \cdot B_{22} \quad \quad \quad - A_{12} \cdot B_{22} \\ \quad \quad \quad - A_{22} \cdot B_{22} - A_{22} \cdot B_{21} + A_{12} \cdot B_{22} + A_{12} \cdot B_{21} \\ \hline A_{11} \cdot B_{11} \quad \quad \quad + A_{12} \cdot B_{21} , \end{array}$$

Expanding C_{12} :

$$\begin{array}{r} A_{11} \cdot B_{12} - A_{11} \cdot B_{22} \\ + A_{11} \cdot B_{22} + A_{12} \cdot B_{22} \\ \hline A_{11} \cdot B_{12} \qquad + A_{12} \cdot B_{22} , \end{array}$$

Expanding C_{21} :

$$\begin{array}{r} A_{21} \cdot B_{11} + A_{22} \cdot B_{11} \\ - A_{22} \cdot B_{11} + A_{22} \cdot B_{21} \\ \hline A_{21} \cdot B_{11} \qquad \qquad + A_{22} \cdot B_{21} , \end{array}$$

Strassen's method

Expanding C_{22} :

$$\begin{array}{rcl}
 A_{11} \cdot B_{11} + A_{11} \cdot B_{22} + A_{22} \cdot B_{11} + A_{22} \cdot B_{22} & & \\
 \quad - A_{11} \cdot B_{22} & + A_{11} \cdot B_{12} & \\
 \quad \quad - A_{22} \cdot B_{11} & \quad - A_{21} \cdot B_{11} & \\
 - A_{11} \cdot B_{11} & - A_{11} \cdot B_{12} + A_{21} \cdot B_{11} + A_{21} \cdot B_{12} & \\
 \hline
 & A_{22} \cdot B_{22} & + A_{21} \cdot B_{12} ,
 \end{array}$$

Strassen's method

Step	Time
Divide the matrices	$O(1)$
Add/subtract to get S_1, \dots, S_{10}	$O(n^2)$
Multiply to get P_1, \dots, P_7	$T(n/2)$ each
Compute $C_{11}, C_{12}, C_{21}, C_{22}$	$O(n^2)$

So the overall time is

$$T(n) = 7T(n/2) + O(n^2)$$

Strassen's method

$$T(n) = 7T(n/2) + O(n^2)$$

- $a = 7$
- $b = 2$
- $d = 2$

$$\log_2 7 \approx 2.807 > d$$

So using case 3 of master theorem, $T(n) = O(n^{\log_2 7}) \approx O(n^{2.807})$

Matrix multiplication

So is matrix multiplication $\Theta(n^{2.807})$?

Let ω denote the exponent.

- $2 \leq \omega \leq 3$ (why?)
- $O(n^{2.807})$ (Strassen, 1969)
- $O(n^{2.376})$ algorithm found in 1990 (Coopersmith-Winograd)
- $O(n^{2.3729})$ in 2013 (Virginia Williams)
- $O(n^{2.3728639})$ in 2014 (François Le Gall)

$2 \leq \omega < 2.373$, but we still don't know if $\omega = 2$ is possible!

Another clever rearrangement

Suppose we have two complex numbers, $a + bi$, and $c + di$.

We can multiply them as such:

$$\begin{aligned}(a + bi)(c + di) &= ac + adi + bci + bdi^2 \\ &= ac + (ad + bc)i - bd \\ &= ac - bd + (ad + bc)i\end{aligned}$$

Another clever rearrangement

$$(a + bi)(c + di) = ac - bd + (ad + bc)i$$

So we have to do four multiplications.

Another clever rearrangement

$$(a + bi)(c + di) = ac - bd + (ad + bc)i$$

But Gauss noticed that

$$ad + bc = (a + b)(c + d) - ac - bd$$

Another clever rearrangement

$$(a + bi)(c + di) = ac - bd + (ad + bc)i$$

$$ad + bc = (a + b)(c + d) - ac - bd$$

So we can compute this with only three multiplications!

Another clever rearrangement

Both are $\Theta(1)$, so does this matter?

Let's consider multiplying n -bit binary numbers...

Multiplying n -bit numbers

Let's consider multiplying n -bit binary numbers x and y .

And let's assume n is a power of 2, for simplicity.

Can we divide and conquer this problem?

Multiplying n -bit numbers

Divide: split x and y into two pieces of length $n/2$

$$x = \boxed{x_L} \boxed{x_R} = 2^{n/2}x_L + x_R$$
$$y = \boxed{y_L} \boxed{y_R} = 2^{n/2}y_L + y_R.$$

E.g., if $x = 10110110_2$, this is also equal to $2^4 \cdot 1011_2 + 0110_2$.

Multiplying n -bit numbers

Divide: split x and y into two pieces of length $n/2$

$$\begin{aligned}(2^{n/2}x_L + x_R)(2^{n/2}y_L + y_R) &= 2^n x_L y_L + 2^{n/2} x_R y_L + 2^{n/2} x_L y_R + x_R y_R \\ &= 2^n x_L y_L + 2^{n/2} (x_R y_L + x_L y_R) + x_R y_R\end{aligned}$$

We need four recursive multiplications.

Multiplying n -bit numbers

This gives a running time of

$$T(n) = 4T(n/2) + O(n)$$

which is $O(n^2)$ by the master method.

Multiplying n -bit numbers

Let's try Gauss's trick:

$$(2^{n/2}x_L + x_R)(2^{n/2}y_L + y_R) = 2^n x_L y_L + 2^{n/2}(x_R y_L + x_L y_R) + x_R y_R$$
$$x_R y_L + x_L y_R = (x_L + x_R)(y_L + y_R) - x_L y_L - x_R y_R$$


Multiplying n -bit numbers

Now with only three recursive multiplications, this gives a running time of

$$T(n) = 3T(n/2) + O(n)$$

which is $O(n^{\log_2 3}) \approx O(n^{1.59})$.

Tower of Hanoi



The goal is to move all discs from peg A to C, but a larger disc can never go on top of a smaller one.

Image credit: Evanherk

https://commons.wikimedia.org/wiki/File:Tower_of_Hanoi.jpeg

Tower of Hanoi

The idea behind the recursive solution:

- If we could somehow get all $n - 1$ discs from A to B, then we could move the largest disc to C.
- Then we could move all $n - 1$ discs from B to C.

Tower of Hanoi

```
moveDiscs(src, dest, n):  
    if n == 1:  
        move top disc from src to dest  
    otherwise:  
        moveDiscs(src, otherPeg, n - 1)  
        move remaining disc from src to dest  
        moveDiscs(otherPeg, dest, n - 1)
```

Tower of Hanoi

Step	Time
Move from src to other	$T(n - 1)$
Move one disc	$O(1)$
Move from other to dest	$T(n - 1)$

Tower of Hanoi

So we get a time of


$$T(n) = 2T(n-1) + O(1)$$

But it's not in the form of the master theorem. Now what?

We can solve this using another method, called recursion trees.


$$T(n)$$

We can solve this using another method, called recursion trees.




Tower of Hanoi

Expanding one level:



Continue until we reach the leaves.

Tower of Hanoi



- How many levels are there?
- How much work is done at each level?
- What is the sum of all levels?

Tower of Hanoi

- How much work is done at each level?
 - 2^k nodes, each with c work, so $c2^k$
- What is the sum of all levels?

$$\sum_{k=0}^{n-1} c2^k = c(2^n - 1)$$

This implies that the recursive algorithm takes $O(2^n)$ steps.

Tower of Hanoi

Another way to see this:

n	#steps
1	1
2	3
3	7
4	15

Tower of Hanoi

We can guess that this is $2^n - 1 = O(2^n)$, but we have to prove it.

Use induction!

Tower of Hanoi

Claim: The recursive algorithm takes $2^n - 1$ steps.

- Base case: $n = 1$, easy to check.
- Inductive case:
 - Assume $T(n - 1) = 2^{n-1} - 1$
 - Use the inductive hypothesis to prove $T(n) = 2^n - 1$:

$$\begin{aligned}T(n) &= 2T(n - 1) + 1 \\&= 2(2^{n-1} - 1) + 1 \\&= 2^n - 2 + 1 \\&= 2^n - 1\end{aligned}$$

Summary

- Divide and conquer is a powerful tool for designing algorithms
- Often get a recurrence of the form $T(n) = aT(n/b) + O(n^d)$
- If so, use the master theorem
- If not, draw the recursion tree and sum the work at each node
- Sometimes you'll get a general answer with some coefficients
- Use induction to solve and verify