START HERE · read the maths. $\int_a^b f(x)\,dx$: only x changes, from a to b; every other letter is temporarily constant. Integrate to $F(x)$, then calculate $[F(x)]_a^b = F(b) - F(a)$ (**upper minus lower**). In $\int_c^d[\int_{a(y)}^{b(y)} f(x,y)\,dx]dy$, do the bracketed **inner x-integral first**, substitute its limits, then do the **outer y-integral**. $\sum_{n=1}^{N}$ = add the N terms · $\prod$ = multiply them · $\bar{x} = \frac{1}{N}\sum x_n$ · $\propto$ = "equal apart from a constant".

Symbols. θ = unknown parameter · $\hat{\theta}$ = estimate · $x[n]$ or $x_1,\dots,x_N$ = data · $E[\cdot]$ = expected/long-run average · $\mathrm{Var}(x) = E[x^2] - E[x]^2$ · $r[k]$ = correlation k steps apart · $P(\theta)$ = power spectral density (PSD) · σ^2 = noise variance · $Q(t) = P(Z > t)$ for standard Gaussian Z.

PART 1 · Probability (1.5 p)


1A · Joint PDF: region, integrals, marginals

IF YOU SEE $p(x,y)$ with a region, or "find c", "probability", "marginal", "conditional", "independent", "domain"

Meaning. $p(x,y)$ describes how likely the pair (x,y) is. The **allowed region** (also called support/domain) is where the given formula applies; outside it, $p = 0$. Integrating p over the WHOLE allowed region must give 1—that is the reason "find c" works. **If asked for the probability of a condition:** shade only the part of the allowed region where that condition is also true, then integrate p over the shaded part using the same slice method below. This answer is a probability; **do not set it equal to 1**.

LIMITS — copy this procedure; do not guess. For $dx\,dy$, x is INNER and y is OUTER. **(1)** Draw the allowed region and label each boundary equation. **(2)** Fix outer y at one allowed value; draw a horizontal slice. **(3)** Where the slice enters/leaves the region gives the lower/upper x-limits; solve the boundary for x if needed. These limits may contain y. **(4)** Sweep the slice across the whole region: the smallest/largest possible y are numerical outer limits. For $dy\,dx$, swap every word $x \leftrightarrow y$ and use a vertical slice. Rectangle → constant limits; non-rectangle → at least one limit depends on the outer letter.

FULL EXAMPLE (Apr25), no skipped steps. $p(x,y) = cx+1$ on $x \ge 0$, $y \ge 0$, $x+y<1$. The boundaries meet at $(0,0),(1,0),(0,1)$; the sloped boundary is $y = 1-x$, equivalently $x = 1-y$. For $dx\,dy$, a horizontal slice has $0 \le x \le 1-y$, and slices cover $0 \le y \le 1$.



A · Find c (normalise). Write the whole double integral before calculating: $1 = \int_{y=0}^{1}\left[\int_{x=0}^{1-y}(cx+1)\,dx\right]dy$. **Inner, x changes; y, c are constants:** $\int_0^{1-y}(cx+1)dx = \left[\frac{cx^2}{2}+x\right]_0^{1-y} = \frac{c}{2}(1-y)^2 + (1-y) - 0$. **Outer, only y remains.** Expand $(1-y)^2 = 1-2y+y^2$: $1 = \int_0^1\left[\frac{c}{2}(1-2y+y^2)+1-y\right]dy = \left[\frac{c}{2}\left(y-y^2+\frac{y^3}{3}\right)+y-\frac{y^2}{2}\right]_0^1$. Substitute $y=1$, then subtract the value at $y=0$ (zero): $1 = \frac{c}{2}(1-1+\frac{1}{3})+1-\frac{1}{2} = \frac{c}{6}+\frac{1}{2}$. Thus $\frac{c}{6} = \frac{1}{2} \Rightarrow \boxed{c=3}$. **Check:** $c=3$ makes $p = 3x+1 \ge 0$ throughout the triangle.

B · Marginal = integrate away the OTHER variable. For $p_X(x)$, use a vertical slice: fixed x, $0 \le y \le 1-x$; the region's x-shadow is $0 \le x \le 1$: $p_X(x) = \int_0^{1-x}(3x+1)dy = [3xy+y]_0^{1-x} = (3x+1)(1-x) = 1+2x-3x^2$, and $p_X(x) = 0$ otherwise. **Check:** $\int_0^1(1+2x-3x^2)dx = [x+x^2-x^3]_0^1 = 1$. Similarly $p_Y(y) = \int_0^{1-y}(3x+1)dx = \frac{3}{2}(1-y)^2 + (1-y) = \frac{5}{2} - 4y + \frac{3}{2}y^2$ for $0 \le y \le 1$, 0 otherwise.

C · Conditional = joint ÷ marginal of the variable GIVEN. $p(x \mid y) = p(x,y)/p_Y(y)$. Here $p(x \mid y) = \frac{3x+1}{\frac{5}{2}-4y+\frac{3}{2}y^2}$ for $0 \le x \le 1-y$ (with $0 \le y < 1$); outside, 0. For $p(y \mid x)$, divide by $p_X(x)$ instead. **Do not reuse the original two-variable range without saying which variable is fixed.**

D · Independent? Both must hold: **(1)** support is a rectangle (possibly infinite), **(2)** $p(x,y) = p_X(x)p_Y(y)$ everywhere. Here the triangle fails test 1 → **not independent**. Square example: $p = 6x^2y$ on $[0,1]^2$ gives $p_X = 3x^2$, $p_Y = 2y$, whose product is $6x^2y \to$ independent.

Fast checks: every pdf/marginal ≥ 0 and integrates to 1 on its stated domain · every probability lies in $[0,1]$ · conditional integrates to 1 over the remaining variable's slice · polynomial rule $\int x^n dx = x^{n+1}/(n+1)$, $n \ne -1$.

1B · WSS / ergodic (multiple choice)

IF YOU SEE "WSS"/"wide-sense stationary"/"stationary", "ergodic", dice, random phase, random amplitude

Definitions. WSS requires BOTH: mean $m(t) = E[x(t)]$ is constant, and $r(t_1,t_2) = E[x(t_1)x(t_2)]$ depends only on the lag $\tau = t_1 - t_2$. **Ergodic in the mean** = the time average of EACH realisation equals the ensemble mean $E[x(t)]$. Ensemble average = average across all possible signals at fixed t; time average = average along one recorded signal.

1. **Mean test:** calculate $E[x(t)]$; deterministic functions of t stay, random letters become their means. If t remains → not WSS; stop.
2. **Correlation test:** multiply $x(t_1)x(t_2)$, expand, then take E. Use $\cos a\cos b = \frac{1}{2}[\cos(a+b)+\cos(a-b)]$ and $E[\cos(k+U)] = 0$ for uniform phase U. If only $t_1 - t_2$ remains → WSS. If t_1+t_2 or a separate t remains → not WSS.
3. **Ergodic test — actually compare averages; do not use "random letter frozen" as a shortcut.** Compute $\lim_{T\to\infty}\frac{1}{2T}\int_{-T}^{T}x(t)dt$ (or the discrete sum) for one realisation and compare with step 1. For ergodicity in variance, repeat with x^2 and compare $E[x^2]$.

Random phase, $x(t) = \cos(\omega t + U)$, U **uniform on** $[0, 2\pi)$**:** $E[x(t)] = 0$, so mean passes. $r = E[\cos(\omega t_1 + U)\cos(\omega t_2 + U)] = \frac{1}{2}\cos(\omega(t_1-t_2)) + \frac{1}{2}E[\cos(\omega(t_1+t_2)+2U)] = \frac{1}{2}\cos(\omega(t_1-t_2)) \to$ **WSS**. One realisation is still a cosine, whose long-time average is $0 \to$ **ergodic in the mean**, even though its phase is fixed.

Random amplitude, $x(t) = A$ **for all** t**:** $E[x] = E[A]$ and $r = E[A^2]$ are constants → **WSS**. But one realisation has time average A, not generally $E[A] \to$ **NOT ergodic in the mean**. Fresh independent dice/samples at every time → time average sees the full distribution → ergodic under the exam's usual assumptions; streamed values uniform in $[-A, A]$ have mean 0.

Vocabulary: random process = set of possible signals; realisation = one recorded signal; random variable = one numerical outcome. Strict stationarity concerns whole distributions; WSS checks only mean and correlation.

1C · Spectral factorisation

IF YOU SEE $P(\theta) = \dfrac{\text{number}}{A + B\cos\theta}$ + "spectral factorisation"/"factorization", "innovation filter $L(z)$", "σ_i^2"

1. Read A and B off the denominator $A + B\cos\theta$ **with their signs** ($25 - 24\cos\theta$: $A = 25$, $B = -24$). Two candidate roots $r = \dfrac{A \pm \sqrt{A^2 - B^2}}{B}$: keep the one between -1 and 1, call it a.
2. $C = \dfrac{B}{2a}$ (just a helper number).
3. $\sigma_i^2 = \dfrac{\text{number on top}}{C}$, $L(z) = \dfrac{1}{1 + a\,z^{-1}}$.
4. Check $|a| < 1$ and $\sigma_i^2 > 0$ — a negative σ_i^2 means you kept the wrong root. Sanity: $\dfrac{\sigma_i^2}{1 + a^2 + 2a\cos\theta}$ must give back the $P(\theta)$ you started from.

Apr24, $P = \frac{7}{25 - 24\cos\theta}$*:*
Step 1 — root: $A = 25$, $B = -24$, so $\sqrt{A^2 - B^2} = \sqrt{625 - 576} = 7$. Candidates: $\frac{25+7}{-24} = -\frac{4}{3}$ and $\frac{25-7}{-24} = -\frac{3}{4}$. Keep $a = -\frac{3}{4}$ because it is between -1 and 1.
Step 2 — helper C**:** $C = \frac{B}{2a} = \frac{-24}{2(-3/4)} = \frac{-24}{-3/2} = 16$.
Step 3 — answers: $\sigma_i^2 = \frac{\text{top}}{C} = \frac{7}{16}$ and $L(z) = \frac{1}{1+az^{-1}} = \boxed{\frac{1}{1 - \frac{3}{4}z^{-1}}}$.
Multiple choice "what must the innovation filter be?" → **minimum phase** (poles and zeros inside the unit circle — the label is the whole answer; Jun26 a).

1D · Expected value and the MC words

IF YOU SEE "expected value", "discrete or continuous", "exactly k of n", "covariance", "contour plot"

- **Expected value = probability-weighted average.** Discrete: $E[x] = \sum_x x\,p(x)$. Continuous: $E[x] = \int x\,p(x)dx$ with the pdf's ordinary domain limits. For $E[x^2]$, put $x^2p(x)$ inside instead. With two variables, use the explicit slice limits from 1A and integrate $x\,p(x,y)$ (or $y\,p(x,y)$ for $E[y]$). The pdf alone sums/integrates to 1, **not** to the mean. **Mode** = most likely value. **Variance** $\mathrm{Var}(x) = E[x^2] - (E[x])^2$; standard deviation $= \sqrt{\mathrm{Var}}$.
- **Discrete** = the values can be listed: "number of ...", counts, dice, digits 0–255. **Continuous** = a measurement: time, distance, weight, money.
- **Rules:** $E[c\cdot z] = cE[z]$ · $E[c] = c\cdot E[z_1+z_2] = E[z_1] + E[z_2]$ — a constant factor comes out. Uniform $[a,b]$ → mean $\frac{a+b}{2}$ ($[-A,A] \to 0$).
- **"Exactly k of n tries"** (binomial): $C(n,k)\,p^k(1-p)^{n-k}$, p = chance of one try, $C(n,k) = \frac{n(n-1)\cdots(n-k+1)}{k!}$. *e.g.* $n = 4$, $k = 2$, $p = 0.3$: $C(4,2) = \frac{4\cdot3}{2\cdot1} = 6 \to 6(0.3)^2(0.7)^2 = 0.2646$.
- **Covariance:** $\mathrm{Cov}(x,y) = E[xy] - E[x]E[y]$ · sign = whether one goes up when the other does · the number changes with units · correlation coefficient $= \mathrm{Cov}/(\sigma_x\sigma_y)$, always in $[-1,1]$. Independent → covariance 0; covariance 0 does **not always** prove independence. *Example:* $p(0,0) = p(1,1) = \frac{1}{2}$: $E[x] = E[y] = E[xy] = \frac{1}{2} \to \mathrm{Cov} = \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$.
- **Contour plot, bi-variate Gaussian:** covariance = the two corner (off-diagonal) entries of the covariance matrix: $0 \to$ ellipses aligned with the axes · **tilted** ellipses → covariance $\ne 0$.
- **Definition MC / T/F:** $F(t) = P(x \le t)$ (the cumulative), so $F(-\infty) = 0$, $F(\infty) = 1$ · independent = $P(A \mid B) = P(A)$, the same as $P(A \cap B) = P(A)P(B)$ ($\cap$ = "both happen") · **disjoint** = they can never both happen, $P(A \cap B) = 0$ (the opposite!) · Bayes = conditional probability rearranged. A T/F list → the swap test of box 2D, any topic.

PART 2 · Estimation (3–3.5 p)

2A · Maximum likelihood (MLE)

IF YOU SEE a pdf $p(x \mid \theta)$, samples $x_1..x_N$, "maximum likelihood", "derive the MLE", "find an estimator", "estimate of ..."

For N **iid** samples (independent, same pdf), $L(\theta) = \prod_{n=1}^{N} p(x_n \mid \theta)$: multiply one pdf by itself N times, using $x_1,\dots,x_N$. Maximise $\ln L$ instead; ln turns multiplication into addition and does not move the maximum.

1. **Log:** $\boxed{\ell(\theta) = \ln L = \sum_n \ln p(x_n \mid \theta)}$. Keep every term containing θ. A constant term repeated for every sample becomes N times that term.
2. **Differentiate with respect to θ:** treat every x_n as a known number; terms without θ vanish. Write $\ell'(\theta)$.
3. **Solve** $\ell'(\theta) = 0$, respecting the parameter's allowed range. Multiply through by denominators first if useful. The solution $\hat{\theta}$ is the MLE; if no stationary point exists, check allowed endpoints. A quick maximum check is $\ell''(\hat{\theta}) < 0$ when applicable.

How to take ln **of a pdf:** write the pdf as $\frac{\text{above}}{\text{below}} \cdot e^u$ (it is pieces multiplied, some in a fraction). Then each piece becomes one term: a piece a **above the line** → $+\ln a$ · a piece b **below the line** → $-\ln b$ · $e^u \to$ just u (the e disappears). Two pieces below the line split: $-\ln(2\theta) = -\ln 2 - \ln\theta$. A power θ^k below the line → $-k\ln\theta$. Derivatives: $\ln\theta \to \frac{1}{\theta}\cdot\theta^k \to k\theta^{k-1}\cdot\frac{1}{\theta} \to -\frac{1}{\theta^2}\cdot e^{c\theta} \to ce^{c\theta}\cdot\cos\theta \to -\sin\theta\cdot\sin\theta \to \cos\theta$. Solving $= 0$: multiply the equation by the denominators (θ^2, θ, ...) first — that does not change the solution. **Trap:** if the pdf only exists for x up to something containing θ (uniform on $[0,\theta]$), these 3 steps are useless — $\hat{\theta} = \max_n x_n$, and **the CRLB does not exist. Only if Fisher information / CRLB is asked:** CRLB is the lowest possible variance of an unbiased estimator; reaching it = efficient. **(1)** Take $\ln p$ for ONE sample. **(2)** Differentiate with respect to the unknown twice. **(3)** Put a minus in front and take the expectation: $i(\theta) = -E[\text{second derivative of } \ln p]$. If x remains, replace it using the known $E[x]$. **(4)** For N iid samples, $\mathcal{I} = N\,i(\theta)$ and $\boxed{CRLB = 1/(N\,i(\theta))}$. *Exponential example,* $p = \lambda e^{-\lambda x}$*:* $\ln p = \ln\lambda - \lambda x \to$ first derivative $= 1/\lambda - x \to$ second derivative $= -1/\lambda^2 \to i(\lambda) = -E[-1/\lambda^2] = 1/\lambda^2 \to CRLB = \lambda^2/N$. **Shortcut values of i:** Gaussian mean, known variance σ^2: $1/\sigma^2$ · exponential rate λ: $1/\lambda^2$ · Bernoulli θ: $1/[\theta(1-\theta)]$ · Rayleigh $p = \frac{x}{\theta^2}e^{-x^2/2\theta^2}$: $4/\theta^2$, hence CRLB $= \theta^2/(4N)$.
Check your $\hat{\theta}$: uniform $[0,\theta] \to \max_n x_n$ (the trap above) · exponential rate $\lambda \to 1/\bar{x}$ · Poisson $\lambda \to \bar{x}$ · Bernoulli $\to k/N$ (k successes in N).
Apr24, follow the three steps. $p(x \mid \theta) = \frac{x}{2\theta}e^{-x^2/(2\theta)}$. One sample: $\ln p = \ln x - \ln 2 - \ln\theta - \frac{x^2}{2\theta}$. Therefore $\ell = \sum \ln x_n - N\ln 2 - N\ln\theta - \frac{\sum x^2}{2\theta}$. Differentiate: $\ell' = -\frac{N}{\theta} + \frac{\sum x^2}{2\theta^2}$. Set to 0 and multiply by $2\theta^2$: $-2N\theta + \sum x_n^2 = 0 \to \boxed{\hat{\theta} = \frac{1}{2N}\sum x_n^2}$.
Jun26 Gamma $\frac{x^{k-1}e^{-x/\theta}}{\Gamma(k)\theta^k}$, n patients: $\ln p = (k-1)\ln x - \frac{x}{\theta} - k\ln\theta - \ln\Gamma(k)$. Add all n logs: $\ell = (k-1)\sum\ln x_i - \frac{\sum x_i}{\theta} - nk\ln\theta - n\ln\Gamma(k)$. Differentiate and drop terms without θ: $\ell' = \frac{\sum x_i}{\theta^2} - \frac{nk}{\theta} = 0$. Multiply by θ^2: $\sum x_i - nk\theta = 0 \to \boxed{\hat{\theta} = \frac{\sum x_i}{nk} = \frac{\bar{x}}{k}}$.
Sigmoid trick. IF YOU SEE $\lambda = \frac{1}{1+e^{-\lambda}}$: treat λ as the unknown, find $\hat{\lambda}$, then convert with $\hat{\theta} = -\ln(\frac{1}{\hat{\lambda}} - 1)$.
Apr25, one observation: $p = 3\lambda e^{-4\lambda x/7} \to \ln p = \ln 3 + \ln\lambda - \frac{4\lambda x}{7} \to$ derivative $\frac{1}{\lambda} - \frac{4x}{7} = 0 \to \frac{1}{\lambda} = \frac{4x}{7} \to \hat{\lambda} = \frac{7}{4x}$. Convert: $\hat{\theta} = -\ln(1/\hat{\lambda} - 1) = -\ln(\frac{4x}{7} - 1)$, valid when $x > \frac{7}{4}$.

2B · MAP with a prior (Beta / Gamma)

IF YOU SEE "prior", "posterior", "MAP", "maximum a posteriori"/"a-posteriori", "MMSE"/"minimum mean square error", "Bayesian estimator". Beta, Gamma

Central rule: posterior ∝ likelihood × prior. **MAP** = parameter value at the posterior's maximum; **MMSE** = posterior mean. Multiplicative constants not containing the unknown do not change MAP.

1. **Likelihood:** write each observation's probability, keep only factors containing the unknown, multiply them; equal bases → add exponents.
2. **Multiply by the prior**, adding exponents again. Call the result $q(\theta)$. It may be unnormalised (its integral need not be 1).
3. Match q to a row below: A, B are the FINAL exponents after including every observation and the prior.

Result looks like	Posterior	MAP	MMSE (mean)
$\theta^A(1-\theta)^B$	Beta$(A+1, B+1)$	$\frac{A}{A+B}$	$\frac{A+1}{A+B+2}$
$\lambda^A e^{-B\lambda}$	Gamma$(A+1, B)$	$\frac{A}{B}$	$\frac{A+1}{B}$

Prior Beta$(a,b) = \theta^{a-1}(1-\theta)^{b-1}$ (its mean $\frac{a}{a+b}$). Prior Gamma$(a,b) = \lambda^{a-1}e^{-b\lambda}$ (mean $\frac{a}{b}$). The prior pulls the estimate towards that mean. One observation x gives: Poisson count → $\lambda^x e^{-\lambda}$ · "k-th success on trial x" → $\theta^k(1-\theta)^{x-k}$ · "k successes in n tries" → $\theta^k(1-\theta)^{n-k}$ · exponential → $\lambda e^{-\lambda x}$.
If your result is not in the table: MAP = the allowed θ where q is largest; usually solve $\frac{d}{d\theta}\ln q = 0$. MMSE = $\frac{\int\theta q(\theta)d\theta}{\int q(\theta)d\theta}$ using the allowed parameter limits. The denominator is required: it makes the posterior's total probability 1. If the best point is outside the prior's allowed interval, choose the nearest endpoint for MAP.
Jun26-2c (Beta): each observation contributes $\theta^2(1-\theta)^{x-2}$.
Step 1 — multiply the five likelihood factors: θ powers add: $2+2+2+2+2 = 10$. The $(1-\theta)$ powers add: $(3-2)+(5-2)+(4-2)+(6-2)+(4-2) = 12$. Likelihood $= \theta^{10}(1-\theta)^{12}$.
Step 2 — multiply the prior: Beta(2,3) contributes $\theta^{2-1}(1-\theta)^{3-1} = \theta^1(1-\theta)^2$. Final powers: $q = \theta^{11}(1-\theta)^{14}$.
Step 3 — read the table with $A = 11$, $B = 14$**:** MAP $= \frac{11}{11+14} = \frac{11}{25} = 0.44$; MMSE $= \frac{12}{11+14+2} = \frac{12}{27} = 0.444$.
Jun26-2d (Gamma), Poisson observations 2, 4, 3, 5, 2, 4*:*
Step 1 — likelihood: λ powers add: $2+4+3+5+2+4 = 20$. Six observations give six factors $e^{-\lambda}$, so likelihood $= \lambda^{20}e^{-6\lambda}$.
Step 2 — multiply prior Gamma(3,2): prior $= \lambda^{3-1}e^{-2\lambda} = \lambda^2 e^{-2\lambda}$. Add powers and exponential coefficients: $q = \lambda^{22}e^{-8\lambda}$.
Step 3 — read the table with $A = 22$, $B = 8$**:** MAP $= \frac{22}{8} = 2.75$; MMSE $= \frac{23}{8} = 2.875$. Without the prior, MLE $= \frac{20}{6} = 3.33$; the prior pulls toward its mean $\frac{3}{2}$.
Nov22: unnormalised $q = \lambda e^{-(x+a)\lambda} \to$ MAP $= 1/(x+a)$; MMSE uses the normalising ratio $= 2/(x+a)$ (fully integrated in Calculations). For a sigmoid parameter, use the conversion at the end of 2A.

2C · Least-squares line

IF YOU SEE points (x,y) + "least squares"/"least-squares"/"LSE", "linear function", "fit a line"

1. Means $\bar{x}$ and $\bar{y}$ (bar = average).
2. Slope: $a = \frac{\sum(x-\bar{x})(y-\bar{y})}{\sum(x-\bar{x})^2}$.
3. Intercept: $b = \bar{y} - a\bar{x}$. Answer: $y = ax + b$. Check: $x = \bar{x}$ must give $y = \bar{y}$.

Apr25, points $(0,3),(2,4),(3,3)$*:*
Step 1 — averages: $\bar{x} = \frac{0+2+3}{3} = \frac{5}{3}$ and $\bar{y} = \frac{3+4+3}{3} = \frac{10}{3}$.
Step 2 — subtract the average from EACH point: $x - \bar{x} = (-\frac{5}{3}, \frac{1}{3}, \frac{4}{3})$ and $y - \bar{y} = (-\frac{1}{3}, \frac{2}{3}, -\frac{1}{3})$.
Step 3 — form the two sums: numerator $= (-\frac{5}{3})(-\frac{1}{3}) + (\frac{1}{3})(\frac{2}{3}) + (\frac{4}{3})(-\frac{1}{3}) = \frac{5+2-4}{9} = \frac{3}{9} = \frac{1}{3}$. Denominator $= (-\frac{5}{3})^2 + (\frac{1}{3})^2 + (\frac{4}{3})^2 = \frac{25+1+16}{9} = \frac{14}{9}$.
Step 4 — slope, then intercept: $a = \frac{1/3}{14/9} = \frac{1}{14}$, $b = \bar{y} - a\bar{x} = \frac{10}{3} - \frac{1}{14}\frac{5}{3} = \frac{45}{14}$. Therefore $\boxed{y = \frac{1}{14}x + \frac{45}{14}}$. Check: putting $x = \bar{x}$ gives $y = \bar{y}$.
Also: $x[n] = \theta + \text{noise} \to \hat{\theta} = \bar{x}$. $x[n] = \ln\theta + \text{noise} \to \hat{\theta} = e^{\bar{x}}$ (noise = random wobble; fitting a constant by least squares is just its average).
Beyond a straight line. $J(\theta) = \sum_n (x[n] - s[n;\theta])^2$, $\hat{\theta} = \arg\min J$ (the θ making J smallest) — write this when they ask "what kind of estimator". One amplitude $s[n] = Ah[n]$: $\hat{A} = \frac{\sum x[n]h[n]}{\sum h[n]^2}$ (*e.g.* $x = 3,5$, $h = 1,2$: $\hat{A} = \frac{3+10}{1+4} = 2.6$).
Linear model $x = \mathbf{H}\theta + w$: $\hat{\theta} = (\mathbf{H}^T\mathbf{H})^{-1}\mathbf{H}^T x$; the **error** $x - \mathbf{H}\hat{\theta}$ is orthogonal (perpendicular) to the columns of $\mathbf{H}$ (it is the error, **not** $\hat{\theta}$); weighted LS with $\mathbf{W} = \mathbf{C}_x^{-1}$ = **BLUE** (best linear unbiased estimator). Nonlinear: transform it (the $e^{\bar{x}}$ line), or iterate **Gauss–Newton** $\theta_{k+1} = \theta_k + (\mathbf{G}^T\mathbf{G})^{-1}\mathbf{G}^T(x - s(\theta_k))$, $\mathbf{G}[:,i] = \frac{\partial s}{\partial\theta_i}$ — Newton–Raphson works too, "only Gauss–Newton" is **false**.

2D · Part 2 true / false

IF YOU SEE "which of the following statements" (in Part 2) about MLE, MAP, MMSE, Fisher, CRLB

T/F method: translate each option into an equation before deciding. A common false option is a true sentence with ONE word swapped: hunt *always · regardless · strictly · maximise/minimise · mean/maximum*. This is a warning sign, **not proof**; "always" is killed by one counterexample, and a true sentence needs no swap word.

The fact (true)	The lie they print (false)
MMSE = **mean** of the posterior; MAP = **maximum**	the two swapped
MMSE = MAP for a symmetric one-peak posterior	"**always** different" / "always equal"
MAP = likelihood × prior, needs a prior (Jun26 c)	"MAP→MLE **regardless of** the prior" · "Bayesians skip priors"
Bayes estimate **minimises** the Bayes risk (Nov22 b)	"maximises"
Quadratic cost → MMSE; uniform cost → MAP	"uniform cost → MMSE"
MLE = argmax of the likelihood **over** θ	"over x" · "only linear models"
MLE is **asymptotically** unbiased and efficient	"**always** reaches the CRLB"
Invariance: $\alpha = \psi(\theta)$ 1-to-1 → $\hat{\theta} = \psi^{-1}(\hat{\alpha})$ (Jan22 c)	—
Fisher of N iid (independent, same pdf) $= N\,i(\theta)$, $\mathcal{I} \ge 0$; $\mathrm{Var}(\hat{\theta}) \ge \frac{1}{\mathcal{I}(\theta)}$	"$\mathcal{I}$ strictly negative" · "$\ge \mathcal{I}(\theta)$"
CRLB $= \frac{1}{Ni(\theta)}$ shrinks $\sim \frac{1}{N}$ (Jan22 b)	"**always** derivable" (no, if range of x depends on θ)
efficient = variance **equal to** the CRLB	"**below** the CRLB"
Fisher info for "x = signal(θ) + Gaussian noise σ^2"**:** $\mathcal{I} = \frac{1}{\sigma^2}\sum\left(\frac{d\,\text{signal}}{d\theta}\right)^2$.	

Nov22 $A[\cos\theta, \sin\theta]$**: phase MLE** $= \arctan\frac{x_2}{x_1}$ (point the model vector at the data), $\mathcal{I} = \frac{A^2}{\sigma^2}$.

PART 3 · Spectral estimation (3 p)

$r[k]$ = autocorrelation at lag k (the given numbers; lag k = k steps apart). Rewrite cosines: $\cos 2\theta = 2\cos^2\theta - 1$, $\cos 3\theta = 4\cos^3\theta - 3\cos\theta$, $\cos\alpha\cos\beta = \frac{\cos(\alpha+\beta)+\cos(\alpha-\beta)}{2}$ (to expand $|L|^2$ of box 1C).
Draw it — the question says "determine **and plot**", only the numbers are graded: $r[k] \to$ vertical **stems** (one stick per lag) at the lags. $P(\theta) \to$ compute the value at $\theta = 0$ and at $\theta = \pi$, draw **one smooth curve** through those two heights and write them on the axis. Peak at 0 = low-pass (low frequencies), peak at π = high-pass, peak in between = band-pass.

3A · Correlogram

IF YOU SEE $r[0], r[1], r[2]\ldots$ + "correlogram", "only these lags", "PSD as a function of $\cos\theta$", correlation values given

$P(\theta) = r[0] + 2r[1]\cos\theta + 2r[2]\cos(2\theta) + 2r[3]\cos(3\theta) + \cdots$. Stop at the largest given lag. **Why the 2?** Correlation at lag $+k$ equals correlation at lag $-k$; their two equal contributions combine into $2r[k]\cos(k\theta)$. The zero lag occurs only once, so **do not double** $r[0]$. From a data list, $\hat{r}[k] = \frac{1}{N}$(sum of products of values k positions apart): use only pairs that both exist, but still divide by the full data length N. Reverse question (P given): $r[0]$ = constant term; $r[k]$ = HALF the coefficient of $\cos(k\theta)$. Check your expression at $\theta = 0$ (every cosine = 1) and $\theta = \pi$ (cosines alternate between -1 and 1).
Jun26 3, 2, 1: $3 + 4\cos\theta + 2\cos 2\theta = 4\cos^2\theta + 4\cos\theta + 1 = (2\cos\theta + 1)^2$. *From data* $x = [1,2,3]$: $\hat{r}[0] = \frac{1+4+9}{3} = \frac{14}{3}$, $\hat{r}[1] = \frac{2+6}{3} = \frac{8}{3}$ (only the 2 overlapping pairs), $\hat{r}[2] = \frac{3}{3} = 1$.

3B · AR(1)

IF YOU SEE "AR(1)", "first-order autoregressive"

Use the LEFT-side convention $x[n] + a\,x[n-1] = i[n]$ (equivalently $x[n] = -a\,x[n-1] + i[n]$), with white-noise variance σ^2. The sign of a must match this equation.

1. $a = -\dfrac{r[1]}{r[0]}$
2. $\sigma^2 = r[0] + a\,r[1]$
3. $P = \dfrac{\sigma^2}{1 + a^2 + 2a\cos\theta}$ (check $|a| < 1$ and $\sigma^2 > 0$, else a sign went wrong)

Apr25 $r[0] = 2$, $r[1] = -\frac{1}{4}$: $a = -\frac{-1/4}{2} = \frac{1}{8}$ · $\sigma^2 = 2 + \frac{1}{8}(-\frac{1}{4}) = \frac{63}{32}$ → $P = \frac{63/32}{1 + \frac{1}{64} + \frac{1}{4}\cos\theta}$ · $\frac{64}{64} = \frac{126}{65 + 16\cos\theta}$.

3C · AR(2) / Yule–Walker

IF YOU SEE "AR(2)", "Yule–Walker", $r[0], r[1], r[2]$

Convention: $x[n] + a_1 x[n-1] + a_2 x[n-2] = i[n]$. Substitute the three given correlations in order:

1. $D = r[0]^2 - r[1]^2$ (shared denominator)
2. $a_1 = \dfrac{-r[1]\,(r[0] - r[2])}{D}$, $a_2 = \dfrac{r[1]^2 - r[0]\,r[2]}{D}$
3. $\sigma^2 = r[0] + a_1 r[1] + a_2 r[2]$
4. $P = \dfrac{\sigma^2}{1 + a_1^2 + a_2^2 + 2a_1(1 + a_2)\cos\theta + 2a_2\cos 2\theta}$

If they want the matrix (Yule–Walker = steps 1–2 in matrix form): $\begin{bmatrix} r[0] & r[1] \\ r[1] & r[0] \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} = -\begin{bmatrix} r[1] \\ r[2] \end{bmatrix}$.

Jun26 $r = 3, 2, 1$: $D = 3^2 - 2^2 = 5$ · $a_1 = \frac{-2(3-1)}{5} = -\frac{4}{5}$ · $a_2 = \frac{2^2 - 3\cdot 1}{5} = \frac{1}{5}$ · $\sigma^2 = 3 + (-\frac{4}{5})2 + \frac{1}{5}(1) = \frac{8}{5}$ → step 4, then $\cos 2\theta = 2\cos^2\theta - 1$ → $P = \frac{10}{5\cos^2\theta - 12\cos\theta + 8}$.

3D · MA(1)

IF YOU SEE "MA(1)", "moving average"

Model: $x[n] = i[n] + b\,i[n-1]$. PSD: $P = r[0] + 2r[1]\cos\theta$ — it uses $r[0], r[1]$ only: ignore $r[2]$ and beyond. If they ask for b and σ^2: solve $b^2 - \frac{r[0]}{r[1]}b + 1 = 0$, keep the root between -1 and 1; then $\sigma^2 = \frac{r[1]}{b}$.

Apr25 $r = 2, -\frac{1}{4}$: $P = 2 - \frac{1}{2}\cos\theta$. b, σ^2 asked too: $b^2 - \frac{r[0]}{r[1]}b + 1 = b^2 + 8b + 1 = 0 \to b = -4 + \sqrt{15} = -0.127$ (the root with $|b| < 1$), $\sigma^2 = \frac{r[1]}{b} = \frac{4 + \sqrt{15}}{4} = 1.97$; check $r[0] = \sigma^2(1 + b^2) = 2$ ✓

3E · Model from an equation

IF YOU SEE $x[n] = \ldots$ **+ "which kind of process", "calculate the autocorrelation of** x**"**

1 · Name the model by looking only at the right-hand side. i means input noise. Only noise terms → **MA**; the largest noise delay is q. Noise AND earlier x terms → **ARMA**; the largest earlier-x delay is p. Earlier x terms with only $i[n]$ → **AR**: use 3B/3C. If asked from a PSD shape: sharp peaks → AR, deep nulls → MA, both → ARMA.

2 · MA autocorrelation: copy coefficients, then multiply pairs. For $x[n] = i[n] + b_1 i[n-1] + b_2 i[n-2] + \cdots$, write the coefficient list $(1, b_1, b_2, \ldots)$ and noise variance σ^2. $r[0] = \sigma^2$(sum of EVERY coefficient squared). $r[1] = \sigma^2$(sum of products of neighbouring coefficients). $r[2] = \sigma^2$(sum of products two positions apart), and so on. After the largest delay, $r[k] = 0$.

Apr24, every pair shown: coefficients $(1, -\frac{1}{2}, \frac{1}{4}, -\frac{1}{2})$, $\sigma^2 = 1$.
$r[0] = 1^2 + (-\frac{1}{2})^2 + (\frac{1}{4})^2 + (-\frac{1}{2})^2 = \frac{25}{16}$.
$r[1] = (1)(-\frac{1}{2}) + (-\frac{1}{2})(\frac{1}{4}) + (\frac{1}{4})(-\frac{1}{2}) = -\frac{3}{4}$.
$r[2] = (1)(\frac{1}{4}) + (-\frac{1}{2})(-\frac{1}{2}) = \frac{1}{4} \cdot r[3] = (1)(-\frac{1}{2}) = -\frac{1}{2}$.

Backwards (coefficients asked from $r[k]$**):** for MA(2), start at the largest lag: $b_2 = r[2]/\sigma^2$; then $r[1] = \sigma^2 b_1(1 + b_2)$, so $b_1 = r[1]/[\sigma^2(1 + b_2)]$. Finally $r[0] = \sigma^2(1 + b_1^2 + b_2^2)$.

3 · ARMA(1,1): move earlier x **to the LEFT.** Match $x[n] + \alpha x[n-1] = i[n] + \beta i[n-1]$; the signs come from this left-side form. Then $P = \sigma^2\dfrac{1 + \beta^2 + 2\beta\cos\theta}{1 + \alpha^2 + 2\alpha\cos\theta}$. If correlations are asked: $r[0] = \sigma^2\dfrac{1 + \beta^2 - 2\alpha\beta}{1 - \alpha^2}$, $r[1] = -\alpha r[0] + \beta\sigma^2$, and each following $r[k]$ is the previous one times $-\alpha$.

Nov22: $x[n] = i[n] - i[n-1] + \frac{3}{4}x[n-1]$ becomes $x[n] - \frac{3}{4}x[n-1] = i[n] - i[n-1]$, so $\alpha = -\frac{3}{4}$, $\beta = -1$. With $\sigma^2 = \frac{7}{8}$: numerator $= \frac{7}{8}(1 + 1 - 2\cos\theta)$, denominator $= 1 + \frac{9}{16} - \frac{3}{2}\cos\theta \to P = \dfrac{28(1 - \cos\theta)}{25 - 24\cos\theta}$.

3F · Pick the plot

IF YOU SEE "which plot", "which figure", "plots A–D", "match the plot"

Work out each PSD at $\theta = 0$ (all cosines $= 1$) and at $\theta = \pi$ ($\cos\theta = -1$, $\cos 2\theta = 1$, $\cos 3\theta = -1$). Pick the plot with those heights. Only a correlogram can go below 0.

Apr25: correlogram 1.13 → 2.13, MA 1.5 → 2.5, AR 1.56 → 2.57 (heights at $0 \to \pi$) → plot D. **Second type** (windowed cosine, DTFT vs DFT — the DTFT is the whole curve, the DFT is the dots on it): infinite cosine = **2 spikes** · windowed = **main lobe + side lobes** (one big bump + small ones) · DFT = **samples of that same curve** · zero padding = **more dots**, not more resolution.

3G · Welch / Bartlett count

IF YOU SEE "Welch"/"Bartlett"/"WOSA", N **samples, segment length** L**, overlap %, "how many segments"**

Convert overlap to a fraction first (50 % → 0.5). Step/start spacing $D = L(1 - \text{overlap})$. Segment starts are $0, D, 2D, \ldots$; include a start only if its whole length-L segment fits. Hence $K = \left\lfloor \frac{N - L}{D} \right\rfloor + 1$. Bartlett = 0 overlap, so $K = \lfloor N/L \rfloor$. Ideal independent-segment variance ratio $= 1/K$; real overlapping Welch segments are correlated, so reduction is less.

Jun26 $N = 1000$, $L = 200$, 50 %: $D = 200(1 - 0.5) = 100$, $K = \frac{1000 - 200}{100} + 1 = 9$ segments, ratio $\frac{1}{9}$.

3H · Energy or power signal

IF YOU SEE "energy or power signal", "total energy", "average power"

- Dies out or has finite length → **energy signal**: $0 < E = \sum_n |x[n]|^2 < \infty$, average power $= 0$.
- Repeats/never dies (sin, cos, nonzero constant, $\cos t\,u(t)$) → **power signal**: $E = \infty$, finite nonzero P = time average of $|x|^2$. A sinusoid $A\sin(\omega t)$ or $A\cos(\omega t)$ has $P = A^2/2$.
- Continuous time: $E = \int_{-\infty}^{\infty} |x(t)|^2 dt$; for a periodic signal, $P = \frac{1}{T}\int_{t_0}^{t_0+T} |x(t)|^2 dt$ over ONE period T. Discrete periodic: $P = \frac{1}{N}\sum_{n=0}^{N-1} |x[n]|^2$.

Apr25: $A\sin\omega t$ → power $\frac{A^2}{2}$. e^{-n} for $n \geq 0$ → energy $\sum e^{-2n} = \frac{1}{1 - e^{-2}} \approx 1.16$ (use $\sum_{n\geq 0} q^n = \frac{1}{1-q}$). *Continuous* e^{-t}, $t \geq 0$: dies out → energy $\int_0^\infty e^{-2t} dt = \frac{1}{2}$ (the $c = 2$ line of the Calculations strip).

3I · Name the method

IF YOU SEE a table of descriptions to fill with method names, "name the method", bias/variance to match

The description says…	Method
$\lvert\text{DFT}\rvert^2$ divided by N, whole signal	Periodogram
DFT of the estimated autocorrelation	Correlogram
DFT of the estimated autocorrelation, then truncate/window it at maximum lag $M < N$; "consistent"	Blackman–Tukey
NON-overlapping segments, average the periodograms	Bartlett
OVERLAPPING segments, average the periodograms	Welch (WOSA)
$\sigma^2\lvert B\rvert^2/\lvert A\rvert^2$, "linear random signal model"	Parametric (AR / MA / ARMA)
"averages periodograms"	Bartlett and Welch

How: hunt ONE trigger word per description — even reworded, the trigger stays.

3J · Part 3 true / false

IF YOU SEE "which of the following statements" (in Part 3) about PSD, windows, periodogram, Welch

Same swap test as 2D — hunt: *improves/reduces · exactly · more/less · both/only · resolution↔leakage*.

The fact (true)	The lie they print (false)
Periodogram is **not** consistent: bias → 0 but $\text{Var}\{\hat{P}\} \approx P^2$ stays (Jun26 b)	"it is consistent"
Bias affects **both** deterministic and stochastic signals (Apr25 a)	"only stochastic"
Bartlett: variance $\propto \frac{1}{K}$, bias **up** (Nov22 a)	"variance drops by zero-padding" (it splits the data)
Welch (overlapping): variance drops **less** than $\frac{1}{K}$	"exactly $\frac{1}{K}$"
Main-lobe width → resolution · side-lobe height → leakage (Nov22 b)	the two swapped
Windows (Hamming…) cut **leakage**; rectangular has best **resolution**	"a window improves resolution" · "triangular beats rectangular"
Zero-padding: more points (step $\frac{2\pi}{L}$), **no** better resolution	"improves resolution"
Window type does **not** change the periodogram's variance; leakage follows the window, not the record length	"longer data/window reduces leakage"
Periodogram and correlogram have the **same** bias	"periodogram has better bias"

Consistent = variance → 0 as $N \to \infty$. The periodogram's bias → 0 but $\text{Var}\{\hat{P}\} \approx P^2$ stays put.

AR order (Jun26-3e). Bigger p → residual error always goes down → risk: **overfitting** (fake peaks). Fix: **AIC** $= N\ln\hat{\sigma}^2 + 2p$ (also FPE, MDL): the error term falls, the penalty $2p$ grows → choose the p with the smallest AIC. Any PSD can be approximated by an AR model of **high enough order**.

PART 4 · Detection (2–2.5 p)

H_0 = nothing there, H_1 = something there. P_{FA} = decide H_1 when H_0 is true (false alarm). P_D = decide H_1 when H_1 is true. $P_M = 1 - P_D$ (miss).

4A · Likelihood-ratio test

IF YOU SEE two pdfs (one for H_0**, one for** H_1**) + "binary hypothesis"/"hypothesis testing", "likelihood ratio test", "when accept** H_1**", threshold** λ, γ, η**, "**P_{FA}**"/"**P_D**"**

1. Put H_1 ON TOP: $L(\mathbf{x}) = p(\mathbf{x} \mid H_1)/p(\mathbf{x} \mid H_0)$. For independent samples, multiply the per-sample ratios; equal exponentials combine by adding exponents. Cancel common factors.
2. Start with $L(\mathbf{x}) > \eta \Rightarrow$ decide H_1. Take ln on both sides (allowed because ln is increasing): $\ln L > \ln\eta$. Expand logs BEFORE isolating the statistic.
3. Isolate x, $\sum x_n$, $\sum |x_n|$, or $\sum x_n^2$. Multiplying/dividing by a negative flips $>$ to $<$; square roots need the stated domain.
4. **Check the decision side numerically:** choose one simple value on each side, evaluate L, and mark where $L > \eta$. Then compute error areas on those sides. Equality can be assigned either way for continuous pdfs (probability 0).

Apr25 $\frac{6x^{11}}{3x^5} = 2x^6 > 0.2 \to x^6 > 0.1 \to x > 0.1^{1/6} = 0.68$.

Jun26 Rayleigh: $p(x \mid H_j) = \frac{x}{\sigma_j^2} e^{-x^2/(2\sigma_j^2)}$, $x \geq 0$, $\sigma_1 > \sigma_0$.

Step 1 — divide p_1 **by** p_0**:** x cancels, leaving $L = \frac{\sigma_0^2}{\sigma_1^2}\exp[\frac{x^2}{2}(\frac{1}{\sigma_0^2} - \frac{1}{\sigma_1^2})]$.

Step 2 — use $L > \eta$ **and take ln:** $\frac{x^2}{2}(\frac{1}{\sigma_0^2} - \frac{1}{\sigma_1^2}) > \ln(\eta\frac{\sigma_0^2}{\sigma_1^2})$.

Step 3 — isolate x**:** the bracket is positive, so no inequality flip. Multiply by its reciprocal and take the positive square root: $\boxed{x > \sqrt{\dfrac{2\sigma_0^2\sigma_1^2}{\sigma_1^2 - \sigma_0^2}\ln(\eta\dfrac{\sigma_1^2}{\sigma_0^2})}}$. Larger separation between the pdfs → better detection.

Apr24 Gaussian $\frac{1}{\sqrt{2\pi}}e^{-y^2/2}$ vs exponential $\frac{1}{4}e^{-y/2}$, $\gamma = 1$: the constant $\ln\frac{1/4}{1/\sqrt{2\pi}} = \ln\frac{4}{2.51} = -0.467 \to -0.467 + \frac{y^2}{2} - \frac{y}{2} > 0$ → $y^2 - y - 0.934 > 0$ → root $y = \frac{1 + \sqrt{1 + 3.74}}{2} = 1.59$ → accept H_1 if $y > 1.59$. $P_{FA} = Q(1.59) = 0.056$, $P_D = \frac{1}{2}e^{-0.795} = 0.23$ (printed answer — that pdf is unnormalised, keep the $\frac{1}{2}$).

Nov22 $\frac{1}{2}e^{-|x|}$ vs $e^{-2|x|}$, N samples: $L = 2^N e^{-\sum|x_n|}$ → accept H_1 if $\sum|x_n| < N\ln 2 - \ln\lambda$.

Probabilities. P_{FA} = area under the H_0 pdf where you decide H_1. P_D = area under the H_1 pdf in the same place. Tail formulas: Gaussian $X \sim N(\mu, \sigma^2)$: $P(X > t) = Q((t - \mu)/\sigma)$; standard Gaussian $\mu = 0, \sigma = 1 \to Q(t)$ · exponential ce^{-cx}, $x \geq 0$: $P(X > t) = e^{-ct}$ and $P(X < t) = 1 - e^{-ct}$ · Rayleigh: $P(X > t) = e^{-t^2/(2\sigma^2)}$ · symmetric two tails $|X| > t \to$ twice one tail. **Always use the distribution under the hypothesis named by the error:** P_{FA} uses H_0; P_D uses H_1.

"Threshold for $P_{FA} = 0.1$": set the H_0 area $= 0.1$, solve for t. **Read** Q **backwards:** with a Gaussian H_0 the t with $Q(t) = 0.1$ is $t = 1.28$ (Numbers strip); then P_D = area under the H_1 pdf from that same t on.

Nov22: $1 - e^{-t} = 0.1 \to t = -\ln 0.9$; then $P_D = 1 - e^{-2t} = 0.19$.

"Plot / indicate P_{FA} and P_M": draw both pdfs and a vertical line at the threshold. Shade the H_0 curve on the "decide H_1" side → P_{FA}. Shade the H_1 curve on the "decide H_0" side → P_M. (Nov22: H_1 when $y < 1$, so $P_{FA} = H_0$ area left of 1, $P_M = H_1$ area right of 1.)

Gaussian with any σ**.** $p(x) = \dfrac{1}{\sigma\sqrt{2\pi}}e^{-x^2/2\sigma^2}$ (mean 0). Two zero-mean Gaussians, σ_1 vs σ_0 (pe21-4b): $\ln L = N\ln\frac{\sigma_0}{\sigma_1} + \frac{1}{2}(\frac{1}{\sigma_0^2} - \frac{1}{\sigma_1^2})\sum_n x^2[n] \to$ decide H_1 when $\sum_n x^2[n]$ is **big** = **energy detector** (bigger variance = more energy).

Left tail (printed with the question — pe21-4a). The printed formula $\frac{1}{2}[1 + \text{erf}\frac{t-\mu}{\sigma\sqrt{2}}]$ gives $P(x < t)$, so $P(x > t) = 1 - $ that. First compute $z = \frac{t-\mu}{\sigma}$, then read $\text{erf}\frac{z}{\sqrt{2}}$ at the column $x = z$ (x runs -3 to 3; the value at $-z$ is minus the value at z).

Matched filter (jan22-4a): correlate the data with the known signal $s[n]$ (FIR filter $h[n] =$ **the-reversed** s) → biggest output SNR; "only for uncorrelated noise" is **false**.

4B · Priors and costs (Bayes)

IF YOU SEE priors $P(H_0), P(H_1)$**, costs** C_{ij}**, "posterior", "Bayesian hypothesis test", "**P_M**" / "miss"**

C_{ij} = cost of deciding i when j is true. C_{10} = false alarm cost, C_{01} = miss cost. "Minimises the Bayes risk" = makes the total cost $C_{00}P(H_0)(1 - P_{FA}) + C_{10}P(H_0)P_{FA} + C_{11}P(H_1)P_D + C_{01}P(H_1)P_M$ as small as possible.

- Threshold $\eta = \dfrac{P(H_0)\,(C_{10} - C_{00})}{P(H_1)\,(C_{01} - C_{11})}$. Equal error costs and 0 for correct → $\eta = \frac{P(H_0)}{P(H_1)}$. Then do 4A with this η.
  Jun26-4c: $\eta = \frac{0.9}{0.1} = 9$ → put $\eta = 9$ in the Rayleigh answer of 4A.
- **Posteriors given** → accept H_1 if $\dfrac{P(H_1 \mid x)}{P(H_0 \mid x)} > \dfrac{C_{10} - C_{00}}{C_{01} - C_{11}}$.
  Apr25-4c: $C_{01} = 100\,C_{10}$ → right side $= 0.01$. Posterior ratio $= 5 > 0.01$ → accept H_1 → the student fails: **yes**.
- **"Where** $p_1 P_1 = p_0 P_0$**"** (p = the two pdfs, P = the two priors) → write the equality and solve.
  Nov22: $0.6 \cdot 2e^{-2y} = 0.4\,e^{-y} \to e^{-y} = \frac{1}{3} \to y = \ln 3$; decide H_1 for $y < \ln 3$. $P_{FA} = 1 - e^{-\ln 3} = \frac{2}{3}$, $P_M = e^{-2\ln 3} = \frac{1}{9}$.

4C · Part 4 true / false

IF YOU SEE "which of the following statements" (in Part 4) about hypothesis testing, P_{FA}**, errors**

Same swap test as 2D — the verb is the question: risk **min** · P_D **max** · P_e **min** (P_e = total error $= P(H_0)P_{FA} + P(H_1)P_M$, equal costs) · η goes **up** with $P(H_0)$, **down** with $P(H_1)$.

The fact (true)	The lie they print (false)
Bayes detection **minimises** the Bayes risk	"maximises"
$C_{01} = C_{10}$, $C_{00} = C_{11} = 0$ → minimises total error P_e	*trap:* scaling all costs leaves η unchanged
Equal priors + equal costs → maximum-likelihood rule	—
Higher $P(H_0)$ raises η → harder to pick H_1	"higher $P(H_1)$ raises η" (lowers — Jun26 c)
Neyman–Pearson: fix P_{FA}, **maximise** P_D	"minimise P_D" · "fix P_D, max P_{FA} "
NP's solution is the LRT; the LRT itself needs **no** priors	"the LRT needs P_0, P_1"
ln of the ratio never changes the decisions	"changes the decisions"
Errors are unavoidable: the pdfs overlap	"a threshold kills both errors"
ROC = P_D plotted against P_{FA} as the threshold moves: up and right · diagonal = guessing · better = top-left	"below diagonal = wrong σ" (flipped)
miss = false negative = type II · false alarm = false positive = type I (Nov22)	labels swapped

Matched filter & energy-detector sentences → the last blocks of 4A: time-reversed s · "only for uncorrelated noise" **false** · $\sum x^2$ **big** · a narrow H_1 pdf → the sum comes out **small**.

Calculations · the algebra under every answer

Integrals. These formulas are **printed in the question itself** (q3-23-2h, nov22-2f) — you copy them: $\int e^{cx}dx = \frac{1}{c}e^{cx}$ · $\int xe^{cx}dx = \frac{cx-1}{c^2}e^{cx}$ · $\int x^2e^{cx}dx = (\frac{x^2}{c} - \frac{2x}{c^2} + \frac{2}{c^3})e^{cx}$ · $\int x^2e^{-cx}dx = -e^{-x}(x^2 + 2x + 2)$ (by parts twice) · $\int \cos u\,du = \sin u$ · $\int \sin u\,du = -\cos u$. Evaluate $[F(x)]_a^b = F(b) - F(a)$: at $x = \infty$ only e^{cx} with $c < 0$ becomes 0, at $x = 0$ every x is 0 so only the plain number is left → $\int_0^\infty e^{-cx}dx = \frac{1}{c}$ · $\int_0^\infty xe^{-cx}dx = \frac{1}{c^2}$ · $\int_0^\infty x^2e^{-cx}dx = \frac{2}{c^3}$.

Nov22-2f: MMSE $= \dfrac{\int_0^\infty \theta\cdot\theta e^{-(x+a)\theta}d\theta}{\int_0^\infty \theta e^{-(x+a)\theta}d\theta} = \dfrac{2/(x+a)^3}{1/(x+a)^2} = \dfrac{2}{x+a}$.

$\ln$ **and** e**.** $\ln(ab) = \ln a + \ln b \cdot \ln\frac{a}{b} = \ln a - \ln b \cdot \ln a^k = k\ln a \cdot \ln e^a = a \cdot e^{\ln a} = a \cdot e^a e^b = e^{a+b}$.

Get the letter alone: $e^{ky} = A \to y = \frac{\ln A}{k} \cdot x^k = a \to x = a^{1/k} \cdot y = k\ln x \to x = e^{y/k}$.

Derivatives. $(c)' = 0 \cdot (\theta^k)' = k\theta^{k-1} \cdot (1/\theta)' = -1/\theta^2 \cdot (\ln\theta)' = 1/\theta \cdot (e^{u(\theta)})' = e^{u(\theta)}u'(\theta) \cdot [\ln f(\theta)]' = f'(\theta)/f(\theta) \cdot (uv)' = u'v + uv' \cdot \cos' = -\sin$, $\sin' = \cos$. Differentiating with respect to θ means all x_n are constants.

Sums. $\sum_{n\geq 0} q^n = \frac{1}{1-q}$ for $|q| < 1 \cdot \sum_n x_n = N\bar{x} \cdot \sum_n c = Nc \cdot \sum_n(ax_n + b) = a\sum_n x_n + Nb$.

Fractions & equations. $\frac{a}{b} \pm \frac{c}{d} = \frac{ad \pm bc}{bd} \cdot \frac{a}{b} \div \frac{c}{d} = \frac{ad}{bc} \cdot \frac{a}{b} = \frac{c}{d} \to ad = bc \cdot ax^2 + bx + c = 0 \to x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ · multiplying by a negative flips $>$ into $<$ · $|x| > t = $ two cases.

Numbers / Gaussian tail. $\ln 2 = 0.693 \cdot \ln 3 = 1.099 \cdot \ln 10 = 2.303 \cdot e^{-2} = 0.135 \cdot \sqrt{2\pi} = 2.5066 \cdot Q(-t) = 1 - Q(t) \cdot Q(0) = 0.5 \cdot Q(1) = 0.1587 \cdot Q(1.28) = 0.10 \cdot Q(1.645) = 0.05 \cdot Q(1.96) = 0.025 \cdot Q(2.33) = 0.01$. For a two-sided interval $\pm t$, central area $= 1 - 2Q(t)$.

Calculation rescue · when an algebra step stops you

1 · Exponential integral: antiderivative, THEN limits. To calculate $\int_0^\infty \theta e^{-b\theta}d\theta$ with $b > 0$, use the xe^{cx} formula above with $x = \theta$, $c = -b$: $\int \theta e^{-b\theta}d\theta = -e^{-b\theta}(\theta/b + 1/b^2)$. At ∞, $e^{-b\theta} = 0$, so the upper value is 0. At 0, the value is $-1/b^2$. Upper minus lower gives $\boxed{0 - (-1/b^2) = 1/b^2}$. Do not forget to subtract the lower value.

2 · Where the Beta MAP formula comes from. If $q = \theta^A(1 - \theta)^B$, then $\ln q = A\ln\theta + B\ln(1 - \theta)$. Differentiate: $A/\theta - B/(1 - \theta)$. Set equal to 0: $A/\theta = B/(1 - \theta) \to A(1 - \theta) = B\theta \to A = (A + B)\theta \to \boxed{MAP = A/(A + B)}$. This is why you only need the final exponents in 2B.

3 · Where the Gamma MAP formula comes from. If $q = \lambda^A e^{-B\lambda}$, then $\ln q = A\ln\lambda - B\lambda$. Differentiate: $A/\lambda - B$. Set to $0 \to A/\lambda = B \to A = B\lambda \to \boxed{MAP = A/B}$. MMSE is NOT the peak; use the table's posterior mean $(A + 1)/B$.

4 · A negative power in an MLE derivative. $\frac{d}{d\theta}[-\frac{S}{2\theta}] = \frac{d}{d\theta}[-\frac{S}{2}\theta^{-1}] = -\frac{S}{2}(-1)\theta^{-2} = \boxed{+\dfrac{S}{2\theta^2}}$. Here $S = \sum x_n^2$ is constant while differentiating with respect to θ. The two minus signs cancel.

5 · Solving an exponential threshold. $e^{-ct} = a \to$ take ln: $-ct = \ln a \to \boxed{t = -\ln(a)/c}$. Example: $e^{-2t} = 0.1 \to -2t = \ln 0.1 = -2.303 \to t = 1.151$. If the equation has e^{-ct} on the OTHER side, first isolate it by division.

6 · Quadratic formula and choosing the MA root. For $b^2 + 8b + 1 = 0$, use $a = 1, b_{quad} = 8, c = 1$: $b = \frac{-8 \pm \sqrt{64 - 4}}{2} = -4 \pm \sqrt{15}$. The two values are -0.127 and -7.873; MA(1) keeps $\boxed{-4 + \sqrt{15} = -0.127}$ because $|b| < 1$. Do not confuse the coefficient b in the quadratic formula with the MA parameter.

7 · AR(2): the jump from coefficients to the final PSD. Jun26 has $a_1 = -\frac{4}{5}$, $a_2 = \frac{1}{5}$, $\sigma^2 = \frac{8}{5}$. Compute the denominator pieces separately: $1 + a_1^2 + a_2^2 = 1 + \frac{16}{25} + \frac{1}{25} = \frac{42}{25}$ · $2a_1(1 + a_2) = 2(-\frac{4}{5})(\frac{6}{5}) = -\frac{48}{25}$ · $2a_2 = \frac{2}{5}$. Therefore $P = \dfrac{8/5}{42/25 - (48/25)\cos\theta + (2/5)\cos 2\theta}$. Now replace $\cos 2\theta = 2\cos^2\theta - 1$: constant $= \frac{42}{25} - \frac{2}{5} = \frac{32}{25}$ and $\cos^2$ coefficient $= \frac{4}{5}$. Multiplying top and bottom by 25 gives $\dfrac{40}{32 - 48\cos\theta + 20\cos^2\theta} = \boxed{\dfrac{10}{8 - 12\cos\theta + 5\cos^2\theta}}$.

8 · Correlogram: the jump to a perfect square. Let $c = \cos\theta$. For Jun26: $3 + 4c + 2\cos 2\theta = 3 + 4c + 2(2c^2 - 1) = 3 + 4c + 4c^2 - 2 = \boxed{4c^2 + 4c + 1 = (2c + 1)^2}$. A squared PSD is nonnegative, which is a useful check.

9 · Sums with actual numbers. For $x = [1, 2, 3]$, $N = 3$ and $\bar{x} = (1 + 2 + 3)/3 = 2$. $\sum x_n^2 = 1^2 + 2^2 + 3^2 = 14$ · $\sum(x_n - \bar{x})^2 = (-1)^2 + 0^2 + 1^2 = 2$ · $\hat{r}[0] = 14/3$ · $\hat{r}[1] = (1 \cdot 2 + 2 \cdot 3)/3 = 8/3$ · $\hat{r}[2] = (1 \cdot 3)/3 = 1$. The sums have fewer products as the lag grows, but the denominator stays $N = 3$.

10 · P_{FA} **and** P_D**: same decision region, different pdf.** Suppose you decide H_1 when $x > t$, with $p(x \mid H_0) = e^{-x}$ and $p(x \mid H_1) = 2e^{-2x}$, $x \geq 0$. Then $P_{FA} = \int_t^\infty e^{-x}dx = [-e^{-x}]_t^\infty = 0 - (-e^{-t}) = e^{-t}$. Use the SAME limits under H_1: $P_D = \int_t^\infty 2e^{-2x}dx = [-e^{-2x}]_t^\infty = e^{-2t}$. If $P_{FA} = 0.1$, then $e^{-t} = 0.1 \to t = -\ln 0.1 = 2.303$ and $P_D = e^{-2(2.303)} = 0.01$. Do not integrate H_1 for P_{FA}.

Final 10-second rescue. Integral → integrate first, substitute upper and lower second · multiplication inside ln → addition outside · division inside ln → subtraction · derivative → letters other than the unknown are constants · multiplying an inequality by a negative → flip it · finish with a range/sign/probability check.