

Value the concept of evidence-based practice as integral to determining best clinical practice.

Appreciate the strengths and weakness of evidence for using lidocaine gel for pain reduction during urethral catheterization in children.

References

- Boots BK, Edmundson EE. A controlled, randomised trial comparing single to multiple application lidocaine analgesia in paediatric patients undergoing urethral catheterisation procedures. *J Clin Nurs*. 2010;19(5–6):744–748.
- Gerard LL, Cooper CS, Duethman KS, et al. Effectiveness of lidocaine lubricant for discomfort during pediatric urethral catheterization. *J Urol*. 2003;170(2 Pt 1):564–567.
- Gray M. Atraumatic urethral catheterization of children. *Pediatr Nurs*. 1996;22(4):306–310.
- Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336(7650):924–926.
- Mularoni PP, Cohen LL, DeGuzman M, et al. A randomized clinical trial of lidocaine gel for reducing infant distress during urethral catheterization. *Pediatr Emerg Care*. 2009;25(7):439–443.
- Vaughn H, Paton EA, Bush A, et al. Does lidocaine gel alleviate the pain of bladder catheterization in young children? A randomized, controlled trial. *Pediatrics*. 2005;116(4):917–920.

*Adapted from the Quality and Safety Education for Nurses website at <http://www.qsen.org>.

In male patients, grasp the penis with the nondominant hand and retract the foreskin. In uncircumcised newborns and infants, the foreskin may be adhered to the shaft; use care when retracting. If the penis is pendulous, place a sterile drape under the penis. Using