

Base individualized care plan on patient values, clinical expertise, and evidence.

Integrate evidence into practice by carefully assessing pain and other symptoms in children at the end of life.

Attitudes

Value the concept of evidence-based practice as integral to determining best clinical practice.

Appreciate strengths and weakness of evidence for symptom assessment and management at the end of life.

References

- Andresen EM, Seecharan GA, Toce SS. Provider perceptions of child deaths. *Arch Pediatr Adolesc Med*. 2004;158(5):430–435.
- Bradshaw G, Hinds PS, Lensing S, et al. Cancer-related deaths in children and adolescents. *J Palliat Med*. 2005;8(1):86–95.
- Carter BS, Howenstein BS, Gilmer MJ, et al. Circumstances surrounding the deaths of hospitalized children: opportunities for pediatric palliative care. *Pediatrics*. 2004;114(3):361–366.
- Contro N, Larson J, Scofield S, et al. Family perspectives on the quality of pediatric palliative care. *Arch Pediatr Adolesc Med*. 2002;156(1):14–19.
- Drake R, Frost J, Collins JJ. The symptoms of dying children. *J Pain Symptom Manage*. 2003;26(1):594–603.
- Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336(7650):924–926.
- Harris MB. Palliative care in children with cancer: which child and when? *J Natl Cancer Inst Monogr*. 2004;32:144–149.
- Hilden JM, Emanuel EJ, Fairclough DL, et al. Attitudes and practices among pediatric oncologists regarding end-of-life care: results of the 1998 American Society of Clinical Oncology Survey. *J Clin Oncol*. 2001;19(1):205–212.