

child.

- d. Reposition the child and repeat the procedure. Measure at least twice (ideally three times). Average the measurements for the final value.
- e. Demonstrate competency when measuring the growth of infants, children, and adolescents. Refresher sessions should be taken when a lack of standardization occurs.

## Correct Answers

- 1. b, c, d;
- 2. c;
- 3. a, b;
- 4. a;
- 5. a, d, e

## References

- American Academy of Pediatrics. *Culturally effective care toolkit*. <http://www.aap.org/en-us/professional-resources/practice-support/Patient-Management/Pages/Culturally-Effective-Care-Toolkit.aspx>; 2011.
- American Academy of Pediatrics Section on Ophthalmology and Committee on Practice and Ambulatory Medicine. Instrument-based pediatric vision screening policy statement. *Pediatrics*. 2012;130(5):983–986.
- Ball JW, Dains JE, Flynn JA, et al. *Seidel's guide to physical examination*. ed 8. Elsevier: St Louis; 2014.
- Batra P, Saha A, Faridi MM. Thermometry in children. *J Emerg Trauma Shock*. 2012;5(3):246–249.
- Blank L, Coster J, O'Cathain A, et al. The appropriateness of, and compliance with, telephone triage decisions: a systematic review and narrative synthesis. *J Adv Nurs*. 2012;68(12):2610–2621.
- Broner N, Embry VV, Gremminger MG, et al. *Mandatory*