

the professionals understand and care.

Many schools have instituted suicide prevention programs. These programs include services such as drop-in counseling and a peer counseling telephone line. Information can also be obtained from the American Association of Suicidology.*

NCLEX Review Questions

1. Which of the following should the nurse discuss with a 14-year-old adolescent female taking isotretinoin (Accutane) for the treatment of acne? Select all that apply.
 - a. Birth control methods
 - b. Feelings of depression
 - c. Sudden thoughts about hurting oneself
 - d. Blurred vision
 - e. Mood swings
2. Joanna, a 19-year-old pregnant female, is at the clinic for her first prenatal care visit. She is in her first trimester. Joanna has been recently diagnosed with HIV and is concerned about the health of her unborn fetus. The nurse counsels Joanna that treatments are available to prevent or minimize perinatal transmission of HIV. The current evidence-based recommendations to prevent perinatal transmission of HIV are to:
 - a. Administer intravenous zidovudine 24 hours before delivery
 - b. Start a combination of antiviral drugs as soon as possible
 - c. Avoid giving antiretroviral drugs until the 28th week of gestation
 - d. Wait until the infant is born because perinatal transmission before delivery is very rare
3. Which of the following conditions may lead to the development of obesity in children and adolescents? Select all that apply.
 - a. Hereditary low metabolism
 - b. Physical inactivity
 - c. Socioeconomic status
 - d. Use of food as a positive reinforcer of desired behaviors