

- [Guidance/Guidance/Manuals/downloads/som107ap_a_hosp](#)
2013.
- Child Health Corporation of America Nursing Falls Study Task Force. Pediatric falls: state of the science. *Pediatr Nurs*. 2009;35(4):227–231.
- Collins AS. Preventing health care–associated infections. Hughes RG. *Patient safety and quality: an evidence-based handbook for nurses*. Agency for Healthcare Research and Quality: Rockville, MD; 2008.
- Considine J, Brennan D. Effect of an evidence-based education programme on ED discharge advice for febrile children. *J Clin Nurs*. 2007;16(9):1687–1694.
- Cook IF, Murtagh J. Comparative reactogenicity and parental acceptability of pertussis vaccines administered into the ventrogluteal area and anterolateral thigh in children aged 2, 4, 6, and 18 months. *Vaccine*. 2003;21(23):3330–3334.
- Cook IF, Murtagh J. Ventrogluteal area—a suitable site for intramuscular vaccination of infants and toddlers. *Vaccine*. 2006;24(13):2403–2408.
- Curley MAQ, Moloney-Harmon PA. *Critical care nursing of infants and children*. ed 2. Saunders/Elsevier: Philadelphia; 2001.
- Curley MA, Quigley SM, Lin M. Pressure ulcers in pediatric intensive care: incidence and associated factors. *Pediatr Crit Care Med*. 2003;4(3):284–290.
- Curley MA, Razmus IS, Roberts KE, et al. Predicting pressure ulcer risk in pediatric patients: the Braden Q Scale. *Nurs Res*. 2003;52(1):22–33.
- Ellett ML, Beckstrand J. Examination of gavage tube placement in children. *J Soc Pediatr Nurs*. 1999;4(2):51–60.
- Emergency Nurses Association. *Family presence at the bedside during invasive procedures and resuscitation*.
<https://www.ena.org/practice-research/research/CPG/Documents/FamilyPresenceCPG.pdf>
2005.
- Essink-Tebbes CM, Wuis EW, Liem KD, et al. Safety of lidocaine-prilocaine cream application four times a day in premature neonates: a pilot study. *Eur J Pediatr*. 1999;158(5):421–423.