

- Semin Fetal Neonatal Med.* 2010;15(3):136–140.
- Watson R. Hyperbilirubinemia. *Crit Care Nurs Clin North Am.* 2009;21(1):97–120.
- Weiner SM, Finnegan LP. Drug withdrawal in the neonate. Gardner SL, Carter BS, Enzman-Hines M, et al. *Merenstein and Gardner's handbook of neonatal intensive care.* ed 7. Mosby/Elsevier: St Louis; 2011.
- Wilson KL, Zelig CM, Harvey JP, et al. Persistent pulmonary hypertension of the newborn is associated with mode of delivery and not with maternal use of selective serotonin reuptake inhibitors. *Am J Perinatol.* 2011;28(1):19–24.
- Witt C. Skin assessment. Tappero EP, Honeyfield ME. *Physical assessment of the newborn.* ed 5. NICU Ink: Santa Rosa, CA; 2015.
- Woythaler MA, McCormick MC, Smith VC. Late preterm infants have worse 24-month neurodevelopmental outcomes than term infants. *Pediatrics.* 2011;127(3):e622–e629.
- Yang LJS. Neonatal brachial plexus palsy — management and prognostic factors. *Semin Perinatol.* 2014;38(4):222–234.
- Zahorodny W, Rom C, Whitney W, et al. The neonatal withdrawal inventory: a simplified score of newborn withdrawal. *J Dev Behav Pediatr.* 1998;19(2):89–93.
- Zaichkin J. *Newborn intensive care: what every parent needs to know.* ed 3. American Academy of Pediatrics: Elk Grove, IL; 2010.
- Zheng JW, Zhang L, Zhou Q, et al. A practical guide to treatment of infantile hemangiomas of the head and neck. *Int J Clin Exp Med.* 2013;6(10):851–860.
- Zlatunich CO, Packman S. Galactosaemia: early treatment with an elemental formula. *J Inherit Metab Dis.* 2005;28(2):163–168.

---

\*Information is available from Vascular Birthmarks Foundation, <http://www.birthmark.org>.

\*<http://www.hmbana.org>.

\*Information is available from the National Institute of Child Health and Human Development's Safe to Sleep Public Education Campaign, <http://www.nichd.nih.gov/sts>.

\*<http://www.safercar.gov/parents/CarSeats/Right-Seat-Age-And-Size->