

- a. Shallow or normal respirations, hypertension, and tachycardia
- b. Fruity breath odor and decreasing level of consciousness
- c. Headache, hunger, and excessive irritability
- d. Normal urine output with specific gravity less than 1.020 and a trace of ketones

Correct Answers

1. b, c, e; 2. b; 3. b; 4. c; 5. b

References

- Alatzoglou KS, Dattani MT. Genetic causes and treatment of isolated growth hormone deficiency — an update. *Nat Rev Endocrinol*. 2010;6(10):562–576.
- American Academy of Pediatrics, Rose SR, Section on Endocrinology and Committee on Genetics of the American Thyroid Association, et al. Update of newborn screening and therapy for congenital hypothyroidism. *Pediatrics*. 2006;7(6):2290–2303.
- American Diabetes Association. Report of the Expert Committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2001;24(Suppl 1):S5–S20.
- Amin N, Mushtaq T, Alvi S. Fifteen-minute consultation: the child with short stature. *Arch Dis Child Educ Pract Ed*. 2015;100(4):180–184.
- Batista DL, Riar J, Keil M, et al. Diagnostic tests for children who are referred for the investigation of Cushing syndrome. *Pediatrics*. 2007;120(3):e575–e586.
- Biro FM, Huang B, Crawford PB, et al. Pubertal correlates in black and white girls. *J Pediatr*. 2006;148(2):234–240.
- Bryant J, Baxter L, Cave CB, et al. Recombinant growth hormone for idiopathic short stature in children and adolescents. *Cochrane Database Syst Rev*. 2007;(3) [CD004440].
- Carel JC, Léger J. Clinical practice: precocious puberty. *N Engl J Med*. 2008;358(22):2366–2377.
- Centers for Disease Control and Prevention. *Children and*