

1. d;
2. b;
3. b;
4. b, c, d;
5. b, d, e

## References

- Ambuel B, Hamlett KW, Marx CM, et al. Assessing distress in pediatric intensive care environments: the COMFORT scale. *J Pediatr Psychol*. 1992;17(1):95–109.
- Amstein P, Broglio K, Wuhrman E, et al. Use of placebos in pain management. *Pain Manag Nurs*. 2011;12(4):225–229.
- Anand KJ, Hickey PR. Halothane-morphine compared with high-dose sufentanil for anesthesia and postoperative analgesia in neonatal cardiac surgery. *N Engl J Med*. 1992;326(1):1–9.
- Azize PM, Humphreys A, Cattani A. The impact of language on the expression and assessment of pain in children. *Intensive Crit Care Nurs*. 2011;27(5):235–243.
- Babl FE, Crellin D, Cheng J, et al. The use of faces, legs, activity, cry and consolability scale to assess procedural pain and distress in young children. *Pediatr Emerg Care*. 2012;28(12):1281–1296.
- Boerlage AA, Ista E, Duivenvoorden HJ, et al. The COMFORT behavior scale detects clinically meaningful effects of analgesic and sedative treatment. *Eur J Pain*. 2015;19(4):473–479.
- Breau LM, McGrath PJ, Camfield CS, et al. Psychometric properties of the non-communicating children's pain checklist—revised. *Pain*. 2002;99(1-2):349–357.
- Cignacco EL, Sellam G, Stoffel L, et al. Oral sucrose and “facilitated tucking” for repeated pain relief in preterms: a randomized controlled trial. *Pediatrics*. 2012;129(2):299–308.
- Claar RL, Walker LS. Functional assessment of pediatric pain patients: psychometric properties of the functional disability inventory. *Pain*. 2006;121(1-2):77–84.