

- in neonate, [200](#), [206t-209t](#)
 - of muscles, [109](#)
 - of neck, in neonates, [200](#), [206t-209t](#)
 - of neurologic system, [109-110](#), [109b](#), [110f-112f](#), [111t-112t](#)
 - in neonate, [202](#)
 - of nose, [97-98](#), [97f](#), [111t-112t](#)
 - in neonates, [200](#), [206t-209t](#)
 - physiologic measurements in, [81-88](#), [82b](#)
 - preparation of child for, [75-77](#), [75b](#), [76t](#), [77f](#)
 - sequence of examination in, [72-75](#)
 - of skin, [88-89](#), [89f](#), [89t](#)
 - in neonate, [198](#), [206t-209t](#)
 - of spine, [108](#)
 - in neonates, [201](#)
- Physical fitness, in adolescence, [459-460](#), [461f](#)
- Physical inactivity, obesity and, [485](#)
- Physical neglect, [419](#)
- Physical stressors, in intensive care unit, [572b](#)
- Physical therapy, chest, [627](#)
- Physiologic anorexia, in toddler, [365](#)
- Physiologic craniotabes, [198](#)
- Physiologic splitting, [103](#)
- Piaget cognitive theory, [44t](#), [45-46](#)
 - in school-age child, [431](#)
- PICCs. *See* [Peripherally inserted central catheters \(PICCs\)](#)
- Pigeon toe, [109](#)
- Pincer grasp, [307-308](#), [307f](#)
- Pinna