

- a. Place the infant to sleep in the prone position.
- b. Place the infant in a prone position when awake (approximately 15 minutes).
- c. Alternate the infant's head position (side of head) when asleep.
- d. Have the infant wear a soft helmet for 23 to 24 hours a day.
- e. Place the infant to sleep in an infant seat twice a day.

## Correct Answers

1. d; 2. a, d, e; 3. b, c; 4. b, c, d; 5. d; 6. b, c

## References

- Adams SM, Good MW, Defranco GM. Sudden infant death syndrome. *Am Fam Physician*. 2009;79(10):870–874.
- Akhnikh S, Engelberts AC, van Sleuwen BE, et al. The excessively crying infant: etiology and treatment. *Pediatr Ann*. 2014;43(4):e69–e75.
- Alanne S, Nermes M, Soderlund R, et al. Quality of life in infants with atopic dermatitis and healthy infants: a follow-up from birth to 24 months. *Acta Pediatr*. 2011;100(8):e65–e70.
- American Academy of Pediatrics. Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. *Pediatrics*. 2008;122(5):1142–1148.
- American Academy of Pediatrics. *Pediatric nutrition handbook*. ed 7. American Academy of Pediatrics: Elk Grove Village, IL; 2014.
- American Academy of Pediatrics, Task Force on Sudden Infant Death Syndrome. SIDS and other sleep-related infant deaths: expansion of recommendations for a safe infant sleeping environment. *Pediatrics*. 2011;128(5):1030–1038.
- Amthor RE, Cole SM, Manary MJ. The use of home-based therapy with ready-to-use therapeutic food to treat malnutrition in a rural area during a food crisis. *J Am Diet Assoc*. 2009;109(3):464–467.
- Ashworth A. Nutrition, food security, and health. Kliegman RM, Stanton BF, St. Geme JW, et al. *Nelson textbook of*