flush. *Pediatr Nurs*. 1992;18(2):111–113.
Goldberg M, Sankaran R, Givelichian L, et al. Maintaining patency of peripheral intermittent infusion devices with heparinized saline and saline: a randomized double blind controlled trial in neonatal intensive care and a review of literature. *Neonat Intensive Care*. 1999;12(1):18–22.
Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336(7650):924–926.
Gyr P, Burroughs T, Smith K, et al. Double blind comparison of heparin and saline flush solutions in maintenance of peripheral infusion devices. *Pediatr Nurs*. 1995;21(4):383–389.
Hanrahan KS, Kleiber C, Berends S. Saline for peripheral intravenous locks in neonates: Evaluating a change in practice. *Neonatal Netw*. 2000;19(2):19–24.
Hanrahan KS, Kleiber C, Fagan C. Evaluation of saline for IV locks in children. *Pediatr Nurs*. 1994;20(6):549–552.
Heilskov J, Kleiber C, Johnson K, et al. A randomized trial of heparin and saline for maintaining intravenous locks in neonates. *J Soc Pediatr Nurs*. 1998;3(3):111–116.
Infusion Nurses Society. Infusion nursing standards of practice. *J Infus Nurs*. 2011;34(1S):S63–S64.
Klenner AF, Fusch C, Rakow A, et al. Benefit and risk of heparin for maintaining peripheral venous catheters in neonates: a placebo-controlled trial. *J Pediatr*. 2003;143(6):741–745.
Kotter RW. Heparin vs. saline for intermittent intravenous device maintenance in neonates. *Neonat Netw*. 1996;15(6):43–47.
Le Duc K. Efficacy of normal saline solution versus heparin solution for maintaining patency of peripheral intravenous catheters in children. *J Emerg Nurs*. 1997;23(4):306–309.
McMullen A, Fioravanti ID, Pollack D, et al. Heparinized saline or normal saline as a flush solution in intermittent intravenous lines in infants and children. *MCN Am J Matern Child Nurs*. 1993;18(2):78–85.
Mok E, Kwong TK, Chan ME. A randomized controlled trial for maintaining peripheral intravenous lock in children. *Int*