

- randomised controlled trial. *BMJ*. 2000;321(7266):931–933.
- Dilli D, Küçük IG, Dallar Y. Interventions to reduce pain during vaccination in infancy. *J Pediatr*. 2009;154(3):385–390.
- Efe E, Ozer ZC. The use of breast-feeding for pain relief during neonatal immunization injections. *Appl Nurs Res*. 2007;20(1):10–16.
- Gaskell S, Binns F, Heyhoe MB, et al. Taking the sting out of needles: education for staff in primary care. *Paediatr Nurs*. 2005;17(4):24–28.
- Hatfield LA, Gusic ME, Dyer AM, et al. Analgesic properties of oral sucrose during routine immunizations at 2 and 4 months of age. *Pediatrics*. 2008;121(2):e327–e334.
- Hogan ME, Probst J, Wong K, et al. A randomized-controlled trial of parent-led tactile stimulation to reduce pain during infant immunization injections. *Clin J Pain*. 2014;30(3):259–265.
- Ipp M, Cohen E, Goldbach M, et al. Effect of choice of measles-mumps-rubella vaccine on immediate vaccination pain in infants. *Arch Pediatr Adolesc Med*. 2004;158(4):323–326.
- Ipp MM, Gol R, Goldbach M, et al. Adverse reactions to diphtheria, tetanus, pertussis-polio vaccination at 18 months of age: effect of injection site and needle length. *Pediatrics*. 1989;83(5):679–682.
- Ipp M, Parkin PC, Lear N, et al. Order of vaccine injection and infant pain response. *Arch Pediatr Adolesc Med*. 2009;163(5):469–472.
- Ipp M, Taddio A, Goldbach M, et al. Effects of age, gender and holding on pain response during infant immunization. *Can J Clin Pharmacol*. 2004;11(1):e2–e7.
- Ipp M, Taddio A, Sam J, et al. Vaccine-related pain: randomised controlled trial of two injection techniques. *Arch Dis Child*. 2007;92(12):1105–1108.
- Kassab M, Sheehy A, King M, et al. A double-blind randomised controlled trial 25% oral glucose for pain relief in 2-month old infants undergoing immunisation. *Int J Nurs Stud*. 2012;49(3):249–256.
- Kostandy R, Anderson GC, Good M. Skin-to-skin contact diminishes pain from hepatitis B vaccine injection in