

- reporting and keeping youth safe*. Administration on Children, Youth and Families, Family and Youth Services Bureau: Washington, DC; 2013.
- Clark JA, Kieh-Lai MW, Sarnaik A, et al. Discrepancies between direct and indirect blood pressure measurements using various recommendations for arm cuff selection. *Pediatrics*. 2002;110(5):920–923.
- Dosman C, Andrews D. Anticipatory guidance for cognitive and social-emotional development: birth to five years. *Paediatr Child Health*. 2012;17(2):75–80.
- Foote JM, Brady LH, Burke AL, et al. *Evidence-based clinical practice guideline on linear growth measurement of children*. https://www.pedsendo.org/assets/education_training/PENS 2014.
- Harlor AD Jr, Bower C, Committee on Practice and Ambulatory Medicine, Section on Otolaryngology—Head and Neck Surgery. Hearing assessment in infants and children: recommendations beyond neonatal screening. *Pediatrics*. 2009;124(4):1252–1263.
- Mathers M, Keyes M, Wright M. A review of the evidence on the effectiveness of children's vision screening. *Child Care Health Dev*. 2010;36(6):754–780.
- Mehra S, Eavey RD, Keamy DG. The epidemiology of hearing impairment in the United States: newborns, children, and adolescents. *Otolaryngol Head Neck Surg*. 2009;140(4):461–472.
- National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics*. 2004;114(2 Suppl 4th Report):555–576.
- Park MK, Menard SW, Schoolfield J. Oscillometric blood pressure standards for children. *Pediatr Cardiol*. 2005;26(5):601–607.
- Purc-Stephenson RJ, Thrasher C. Patient compliance with telephone triage recommendations: a meta-analytic review. *Patient Educ Couns*. 2012;87(2):135–142.
- Purc-Stephenson RJ, Thrasher C. Nurses' experiences with