

- Clark L. Pain management in the pediatric population. *Crit Care Nurs Clin North Am.* 2011;23(2):291–301.
- Cline ME, Herman J, Shaw ER, et al. Standardization of the visual analogue scale. *Nurs Res.* 1992;41(6):378–380.
- Crosta QR, Ward TM, Walker AJ, et al. A review of pain measures for hospitalized children with cognitive impairment. *J Spec Pediatr Nurs.* 2014;19(2):109–118.
- De Jong AE, Tuinebreijer WE, Bremer M, et al. Construct validity of two pain behavior observation measurement instruments for young children with burns by Rasch analysis. *Pain.* 2012;153(11):2260–2266.
- Fernandes AM, De Campos C, Batalha L, et al. Pain assessment using the adolescent pediatric pain tool: a systematic review. *Pain Res Manag.* 2014;19(4):212–218.
- Fortier MA, Wahi A, Bruce C, et al. Pain management at home in children with cancer: a daily diary study. *Pediatr Blood Cancer.* 2014;61(6):1029–1033.
- Franck LS, Harris SK, Soetenga DJ, et al. The Withdrawal Assessment Tool–1 (WAT-1): an assessment instrument for monitoring opioid and benzodiazepine withdrawal symptoms in pediatric patients. *Pediatr Crit Care Med.* 2008;9(6):573–580.
- Franck LS, Ridout D, Howard R, et al. A comparison of pain measures in newborn infants after cardiac surgery. *Pain.* 2011;152(8):1758–1765.
- Franck L, Vilardi J. Assessment and management of opioid withdrawal in ill neonates. *Neonatal Netw.* 1995;14(2):39–48.
- Gaina A, Sekine M, Chen X, et al. Validity of child sleep diary questionnaire among junior high school children. *J Epidemiol.* 2004;14(1):1–4.
- Garra F, Singer AJ, Domingo A, et al. The Wong-Baker pain FACES scale measures pain, not fear. *Pediatr Emerg Care.* 2013;29(1):17–20.
- Gold JI, Mahrer NE, Yee J, et al. Pain, fatigue and health-related quality of life in children and adolescents with chronic pain. *Clin J Pain.* 2009;25(5):407–412.
- Habich M, Wilson D, Thielk D, et al. Evaluating the effectiveness of pediatric pain management guidelines. *J Pediatr Nurs.* 2012;27(4):336–345.