

Describe methods for using NS or HS flush solution in pediatric IV lines.

Skills

Base individualized care plan on patient values, clinical expertise, and evidence.

Integrate evidence into practice on NS or HS flush solution in pediatric IV lines.

Attitudes

Value the concept of evidence-based practice as integral to determining best clinical practice.

Appreciate the strengths and weakness of evidence for NS or HS flush solution in pediatric IV lines.

References

American Society of Hospital Pharmacists Commission on Therapeutics. ASHP therapeutic position statement on the institutional use of 0.9% sodium chloride injection to maintain patency of peripheral indwelling intermittent infusion devices. *Am J Health Syst Pharm*. 2006;63(13):1273–1275.

Arnts IJ, Heijnen JA, Wilbers HT, et al. Effectiveness of heparin solution versus normal saline in maintaining patency of intravenous locks in neonates: a double blind randomized controlled study. *J Adv Nurs*. 2011;67(12):2677–2685.

Beecroft PC, Bossert E, Chung K, et al. Intravenous lock patency in children: dilute heparin versus saline. *J Pediatr Pharm Practice*. 1997;2(4):211–223.

Crews BE, Gnann KK, Rice MH, et al. Effects of varying intervals between heparin flushes on pediatric catheter longevity. *Pediatr Nurs*. 1997;23(1):87–91.

Danek GD, Noris EM. Pediatric IV catheters: efficacy of saline