

- Ann.* 2014;43(7):e169–e175.
- Twycross A. Managing pain in children: where to from here? *J Clin Nurs.* 2010;19(15-16):2090–2099.
- Uman LS, Birnie KA, Noel M, et al. Psychological interventions for needle-related procedural pain and distress in children and adolescents. *Cochrane Database Syst Rev.* 2013;(10) [CD005179].
- Valrie CR, Bromberg MH, Palermo T, et al. A systematic review of sleep in pediatric pain populations. *J Dev Behav Pediatr.* 2013;34(2):120–128.
- Van Dijk A, McGrath PA, Pickett W, et al. Pain prevalence in 9- to 13-year-old schoolchildren. *Pain Res Manage.* 2006;11(4):234–240.
- Varni JW, Seid M, Rode CA. The PedsQL: measurement model for the pediatric quality of life inventory. *Med Care.* 1999;37(2):126–139.
- Villarruel AM, Denyes MJ. Pain assessment in children: theoretical and empirical validity. *Adv Nurs Sci.* 1991;14(2):32–41.
- Voepel-Lewis T, Malviya S, Tait AR, et al. A comparison of the clinical utility of pain assessment tools for children with cognitive impairment. *Anesth Analg.* 2008;106(1):72–78.
- von Baeyer CL, Lin V, Seidman LC, et al. Pain charts (body maps or manikins) in assessment of the location of pediatric pain. *Pain Manag.* 2011;1(1):61–68.
- Walker LS, Greene JW. The functional disability inventory: measuring a neglected dimension of child health status. *J Pediatr Psychol.* 1991;16(1):39–58.
- Wong DL, Baker CM. Pain in children: comparison of assessment scales. *Pediatr Nurs.* 1988;14(1):9–17.
- World Health Organization. *WHO guidelines on the pharmacological treatment of persisting pain in children with medical illnesses.* World Health Organization: Geneva; 2012.
- Yellon RF, Kenna MA, Cladis FP, et al. What is the best non-codeine post adenotonsilectomy pain management for children? *Laryngoscope.* 2014;124(8):1737–1738.
- Yilmaz G, Cavlan N, Oguz M, et al. Oral sucrose administration to reduce pain response during immunization in 16-19-month infants: a randomized,