

and rough activities until allowed by the surgeon.”

5. What is the 24-hour fluid requirement for a child weighing 32 kg?
- a. 1920 ml/day
 - b. 1740 ml/day
 - c. 1840 ml/day
 - d. 1620 ml/day

Correct Answers

1. a, c, e; 2. a, b, d; 3. a; 4. a, c, e; 5. b

References

- American Academy of Pediatrics Subcommittee on Urinary Tract Infection, Steering Committee on Quality Improvement and Management, Roberts KB. Urinary tract infection: clinical practice guideline for the diagnosis and management of the initial UTI in febrile infants and children 2 to 24 months. *Pediatrics*. 2011;128(3):595–610.
- American Academy of Pediatrics Task Force on Circumcision. Circumcision policy statement. *Pediatrics*. 2012;130(3):585–586.
- Balat A, Karakok M, Guler E, et al. Local defense systems in the prepuce. *Scand J Urol Nephrol*. 2008;42(1):63–65.
- Blatt AE, Liebman SE. Drug induced acute kidney injury. *Hosp Med Clin*. 2013;2(4):e525–e541 <http://www.sciencedirect.com/science/article/pii/S22115>
- Bukowski TP, Zeman PA. Hypospadias: of concern but correctable. *Contemp Pediatr*. 2001;18(2):89–109.
- Carrero JJ, Stenvinkel P, Cuppari L, et al. Etiology of the protein-energy wasting syndrome in chronic kidney disease: a consensus statement from the International Society of Renal Nutrition and Metabolism (ISRNM). *J Ren Nutr*. 2013;23(2):77–90.
- Centers for Disease Control and Prevention. *Recommended immunization schedules for persons aged 0 through 18 years—United States*. <http://www.cdc.gov/vaccines/schedules>;