

References

- Abdel Razek A, Az El-Dein N. Effect of breast-feeding on pain relief during infant immunization injections. *Int J Nurs Pract.* 2009;15(2):99–104.
- Balshem H, Helfand M, Schunemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol.* 2011;64(4):401–406.
- Blount RL, Devine KA, Cheng PS, et al. The impact of adult behaviors and vocalizations on infant distress during immunizations. *J Pediatr Psychol.* 2008;33(10):1163–1174.
- Campbell L, Pillai Riddell R, Garfield H, et al. A cross-sectional examination of the relationship between caregiver proximal soothing and infant pain over the first year of life. *Pain.* 2013;154(6):813–823.
- Chermont AG, Falcão LF, de Souza Silva EH. Skin-to-skin contact and/or oral 25% dextrose for procedural pain relief for term newborn infants. *Pediatrics.* 2009;124(6):e1101–e1107.
- Cohen LL, Bernard RS, McClellan C, et al. Topical anesthesia versus distraction for infants' immunization distress: evaluation with a 6-month follow-up. *Children's Health Care.* 2006;35(2):103–121.
- Cohen LL, MacLaren JE, Fortson BL, et al. Randomized clinical trial of distraction for infant immunization pain. *Pain.* 2006;125(1-2):165–171.
- Cramer-Berness LJ, Friedman AG. Behavioral interventions for infant immunizations. *Children's Health Care.* 2005;34(2):95–111.
- Curry DM, Brown C, Wrona S. Effectiveness of oral sucrose for pain management in infants during immunizations. *Pain Manag Nurs.* 2012;13(3):139–149.
- Davenport JM. A systematic review ascertain whether the standard needle is more effective than a longer or wider needle in reducing the incidence of local reaction in children receiving primary immunization. *J Adv Nurs.* 2004;46(1):66–77.
- Diggle L, Deeks J. Effect of needle length on incidence of local reactions to routine immunisation in infants aged 4 months: