

- 2012;17(5):153–154.
- Shah V, Ohlsson A. Venepuncture versus heel lance for blood sampling in term neonates. *Cochrane Database Syst Rev*. 2007; (4) [CD001452].
- Shen V, Li X, Murdock M, et al. Recombinant tissue plasminogen activator (alteplase) for restoration of function to occluded central venous catheters in pediatric patients. *J Pediatr Hematol Oncol*. 2003;25(1):38–45.
- Shepherd AJ, Glenesk A, Niven CA, et al. A Scottish study of heel-prick blood sampling in newborn babies. *Midwifery*. 2005;22(2):158–168.
- Sobhani P, Christopherson J, Ambrose PJ, et al. Accuracy of oral liquid measuring devices: comparison of dosing cup and oral dosing syringe. *Ann Pharmacother*. 2008;42(1):46–52.
- Stevens B, Johnston C, Taddio A, et al. Management of pain from heel lance with lidocaine-prilocaine (EMLA) cream: is it safe and efficacious in preterm infants? *J Dev Behav Pediatr*. 1999;20(4):216–221.
- Taylor C, Subaiya L, Corsino D. Pediatric cuffed endotracheal tubes: an evolution of care. *Ochsner J*. 2011;11(1):52–56.
- Tillett J. Adolescents and informed consent: ethical and legal issues. *J Perinat Neonatal Nurs*. 2005;19(2):112–121.
- Tobias JD, Ross AK. Intraosseous infusion: a review for the anesthesiologist with a focus on pediatric use. *Anesth Analg*. 2010;110(2):391–401.
- Turner TW. Intravesical catheter knotting: an uncommon complication of urinary catheterization. *Pediatr Emerg Care*. 2004;20(2):115–117.
- Uman LS, Chambers CT, McGrath PJ, et al. Psychological interventions for needle-related procedural pain and distress in children and adolescents. *Cochrane Database Syst Rev*. 2006;(4) [CD005179].
- Vertanen H, Fellman V, Brommels M, et al. An automatic incision device for obtaining blood samples from the heels of the preterm infants causes less damage than a conventional manual lancet. *Arch Dis Child Fetal Neonatal Ed*. 2001;84(1):F53–F55.
- Walsh A, Edwards H. Management of childhood fever by parents: literature review. *J Adv Nurs*. 2006;54(2):217–222.