

Trained nighttime crying, [344t](#)

Transcatheter device closure, in cardiac catheterization, [740t](#)

Transcranial Doppler (TCD), for sickle cell anemia, [792-793](#)

Transcutaneous bilirubinometry (TcB), [259](#)

Transcutaneous monitoring (TCM), [626](#)

Transdermal analgesia, [132](#)

Transesophageal recording, [772](#)

Transfusion

- exchange, for hemolytic disease of the newborn, [266](#), [266f](#)
- intrauterine, for hemolytic disease of the newborn, [266](#)

Transgender family, [19](#)

Transient cyanosis, [190](#)

Transilluminator, [614](#), [614f](#)

Transitional objects, in social development, [358-359](#), [359f](#)

Transmission-based precautions, in infection control, [150](#), [151b](#), [591](#), [592b](#)

Transmucosal analgesia, [132](#)

Transplantation

- blood/marrow, [818-819](#)
 - for cancer, [818-819](#), [825f](#)
 - for leukemia, [828](#)
- heart, [774-775](#)
- for renal failure, [870](#)

Transportation, of infants and children, [592-593](#), [593f](#)

Transposition, of great vessels, [750b-752b](#)

Trauma, in visual impairment, [542-544](#)

Traumatic injury, [947-958](#)

- amputation in, [957-958](#)