

- b. Discontinue breastfeeding and administer Pedialyte for 24 hours.
- c. Observe infant for labored breathing or apnea (cessation of breathing).
- d. Instill normal saline drops in both nares and suction thoroughly before feeding and before placing to sleep.
- e. Place infant to sleep on his side with the head of bed slightly elevated to facilitate breathing.
- f. Keep the infant out of daycare or nursery.

## Correct Answers

1. c; 2. a, b, d, f; 3. a; 4. c; 5. a, c, d, f

## References

- Abrahams SW, Labbok MH. Breastfeeding and otitis media: a review of recent evidence. *Curr Allergy Asthma Rep*. 2011;11(6):508–512.
- Akinbami LJ, Moorman JE, Liu X. Asthma prevalence, health care use, and mortality: United States, 2005–2009. *Natl Health Stat Report*. 2011;32:1–14.
- American Academy of Otolaryngology—Head and Neck Surgery. *Clinical indicators: adenoidectomy*. <http://www.entnet.org/sites/default/files/Adenoidectomy-CI%20Updated%208-7-14.pdf>; 2012.
- American Academy of Pediatrics. Sports drinks and energy drinks for children and adolescents: are they appropriate? *Pediatrics*. 2011;127(6):1182–1189.
- American Academy of Pediatrics. *Withdrawal of cold medicines: addressing parent concerns*. <https://www.aap.org/en-us/professional-resources/practice-support/pages/Withdrawal-of-Cold-Medicines-Addressing-Parent-Concerns.aspx>; 2015.
- American Academy of Pediatrics Committee on Infectious Diseases, American Academy of Pediatrics Bronchiolitis Guidelines Committee. Updated guidance for palivizumab prophylaxis among infants and young children at increased