Although the risk of vitamin A toxicity from these doses (they are 100 to 200 times the recommended dietary allowance) is relatively low, nurses should instruct parents on safe storage of the drug. Ideally, vitamin A should be dispensed in the age-appropriate unit dose to prevent excessive administration and possible toxicity.

Provide Comfort

Many communicable diseases cause skin manifestations that are bothersome to the child. The chief discomfort from most rashes is itching, and measures such as cool baths (usually without soap) and lotions (e.g., calamine) are helpful.

Nursing Alert

When lotions with active ingredients such as diphenhydramine in Caladryl are used, they are applied sparingly, especially over open lesions, where excessive absorption can lead to drug toxicity. Use these lotions with caution in children who are simultaneously receiving an oral antihistamine. Cooling the lotion in the refrigerator beforehand often makes it more soothing on the skin than at room temperature.

To avoid overheating, which increases itching, children should wear lightweight, loose, nonirritating clothing and keep out of the sun. If the child persists in scratching, keep the nails short and smooth or use mittens and clothes with long sleeves or legs. For severe itching, antipruritic medication, such as diphenhydramine (Benadryl) or hydroxyzine (Atarax), may be required, especially when the child has trouble sleeping because of itching. Loratadine, cetirizine, and fexofenadine do not cause drowsiness and may be preferred for urticaria during the day.

An elevated temperature is common, and both antipyretic medicine (acetaminophen or ibuprofen) and environmental manipulation are implemented (see Controlling Elevated Temperatures in Chapter 20). Acetaminophen is effective in lowering the fever but does not significantly reduce the symptoms of itching, anorexia, abdominal pain, fussiness, or vomiting.

A sore throat, another frequent symptom, is managed with lozenges, saline rinses (if the child is old enough to cooperate), and