

advance of the procedure to allow for processing of information.

- d. Provide privacy; describe how the body will be covered and what will be exposed.
- e. Allowing adolescents to talk with other adolescents who have had the same procedure may increase their level of anxiety and is not recommended.

## Correct Answers

1. a; 2. a, b, c, d; 3. b, c, d; 4. b; 5. a, d

## References

- Abo A, Chen L, Johnston P, et al. Positioning for lumbar puncture in children evaluated by bedside ultrasound. *Pediatrics*. 2010;125(5):e1149–e1153.
- American Academy of Pediatrics, Committee on Pediatric Emergency Medicine and Committee on Bioethics. Consent for emergency medical services for children and adolescents. *Pediatrics*. 2011;128(2):427–433.
- American Academy of Pediatrics, Subcommittee on Urinary Tract Infections, Steering Committee on Quality Improvement and Management, Roberts KB. Urinary tract infection: clinical practice guideline for the diagnosis and management of the initial UTI in febrile infants and children 2 to 24 months. *Pediatrics*. 2011;128(3):595–610.
- American Academy of Pediatrics, Task Force on Sudden Infant Death Syndrome. SIDS and other sleep-related infant deaths: expansion of recommendations for a safe infant sleeping environment. *Pediatrics*. 2011;128(5):1030–1039.
- American Association of Critical Care Nurses. *Family presence during CPR and invasive procedures*. <http://ajcc.aacnjournals.org/content/16/3/283.full>; 2006.
- American Heart Association. 2010 American Heart Association guidelines for CPR and ECC. *Circulation*. 2010;122(Suppl 2):S276–S282.
- Amlung SR, Miller WL, Bosley LM. The 1999 National