

- and scoring methods. *Pain*. 1975;1(3):277–299.
- Meredith JR, O'Keefe KP, Galwankar S. Pediatric procedural sedation and analgesia. *J Emerg Trauma Shock*. 2008;1(2):88–96.
- Merkel SI, Voepel-Lewis T, Shayevitz JR, et al. The FLACC: a behavioral scale for scoring postoperative pain in young children. *Pediatr Nurs*. 1997;23(3):293–297.
- Michelet D, Andreu-Gallien J, Bensalah T, et al. A meta-analysis of the use of nonsteroidal antiinflammatory drugs for pediatric postoperative pain. *Anesth Analg*. 2012;114(2):393–406.
- Mudd S. Intranasal fentanyl for pain management in children: a systematic review of the literature. *J Pediatr Health Care*. 2011;25(5):316–322.
- Myers C, Stuber ML, Bonamer-Rheingans JL, et al. Complementary therapies and childhood cancer. *Cancer Control*. 2005;12(3):172–180.
- Oakes LL. *Infant and child pain management*. Springer Publishing: New York; 2011.
- Pasero C, McCaffrey M. *Pain assessment and pharmacologic management*. Elsevier: St Louis; 2011.
- Pillai Riddell R, Racine N, Turcotte K, et al. Nonpharmacological management of procedural pain in infants and young children: an abridged Cochrane review. *Pain Res Manag*. 2011;16(5):321–330.
- Puchalski M, Hummel P. The reality of neonatal pain. *Adv Neonatal Care*. 2002;2(5):233–244.
- Quinn BL, Sheldon LK, Cooley ME. Pediatric pain assessment by drawn faces scales: a review. *Pain Manag Nurs*. 2014;15(4):909–918.
- Racoosin JA, Roberson DW, Pacanowski MA, et al. New evidence about an old drug—risk with codeine after adenotonsillectomy. *N Engl J Med*. 2013;368(23):2155–2157.
- Rastogi S, Campbell F. Drugs for neuropathic pain. McGrath PC, Stevens BJ, Walker SM, et al. *Oxford textbook of paediatric pain*. Oxford University Press: Oxford UK; 2014.
- Saps M, Youssef N, Miranda A, et al. Multicenter, randomized, placebo-controlled trial of amitriptyline in children with functional gastrointestinal disorders.