

Pediatr Nurs. 2009;14(4):239–244.

Jean-Mary MB, Dicanzio J, Shaw J, et al. Limited accuracy and reliability of infrared axillary and aural thermometers in a pediatric outpatient population. *J Pediatr.* 2002;141(5):671–676.

Martin SA, Kline AM. Can there be a standard for temperature measurement in the pediatric intensive care unit? *AACN Clin Issues.* 2004;15(2):254–266.

Schuh S, Komar L, Stephens D, et al. Comparison of the temporal artery and rectal thermometry in children in the emergency department. *Pediatr Emerg Care.* 2004;20(11):736–741.

Siberry GK, Diener-West M, Schappell E, et al. Comparison of temple temperatures with rectal temperatures in children under 2 years of age. *Clin Pediatr.* 2002;41(6):405–414.

Stine CA, Flook DM, Vincze DL. Rectal versus axillary temperatures: is there a significant difference in infants less than 1 years of age? *J Pediatr Nurs.* 2012;3:265–270.

Titus MO, Hulsey T, Heckman J, et al. Temporal artery thermometry utilization in pediatric emergency care. *Clin Pediatr.* 2009;48(2):190–193.

*Adapted from the Quality and Safety Education for Nurses (QSEN) Institute.

Box 4-9

Recommended Temperature Screening Routes in Infants and Children

Birth to 2 Years Old

Axillary

Rectal—if definitive temperature reading is needed for infants older than 1 month of age

2 to 5 Years Old

Axillary