

high-risk behaviors.

- d. In such infants, antiretroviral medication should be started within 12 hours of birth.

Correct Answers

1. a, b, d, e;
2. b;
3. a, b, d, e, f;
4. c;
5. b

References

- Ackerman JP, Riggins T, Black MM. A review of the effects of prenatal cocaine exposure among school-aged children. *Pediatrics*. 2010;125(3):554–565.
- Adamkin DH, American Academy of Pediatrics, Committee on Fetus and Newborn. Postnatal glucose homeostasis in late-preterm and term infants. *Pediatrics*. 2011;127(3):575–579.
- Ahmed AH, Sands LP. Effects of pre- and post-discharge interventions on breastfeeding outcomes and weight gain among premature infants. *J Obstet Gynecol Neonatal Nurs*. 2010;39(1):53–63.
- Alfaleh K, Anabrees J, Bassler D, et al. Probiotics for prevention of necrotizing enterocolitis in preterm infants. *Cochrane Database Syst Rev*. 2011;2011(3) [CD005496].
- Altimier L. Thermoregulation: what's new, what's not. *Newborn Infant Nurs Rev*. 2012;12(1):51–63.
- Altimier L, White RD. The neonatal intensive care unit (NICU) environment. Kenner C, Lott J. *Comprehensive neonatal care: an interdisciplinary approach*. ed 5. Springer: New York; 2014.
- American Academy of Pediatrics, American College of Obstetricians and Gynecologists. *Guidelines for perinatal care*.