

- transplantation: a review. *Transplant Res Risk Manage*. 2013;5:21–31.
- Sijstermans K, Hack WW, Meijer RW, et al. The frequency of undescended testis from birth to adulthood: a review. *Int J Androl*. 2008;31(1):1–11.
- Suominen JS, Santtila P, Taskinen S. Sexual function in patients operated for bladder exstrophy and epispadias. *J Urol*. 2015;194(1):195–199.
- Susantitaphong P, Cruz DN, Cerda J, et al. World incidence of AKI: a meta-analysis. *Clin J Am Soc Nephrol*. 2013;8(9):1482–1493.
- Thumfart J, Pommer W, Querfeld U, et al. Intensified hemodialysis in adults, and in children and adolescents. *Dtsch Arztebl Int*. 2014;111(14):237–243.
- Vasudeva P, Madersbacher H. Factors implicated in pathogenesis of urinary tract infections in neurogenic bladders: some revered, few forgotten, others ignored. *Neurourol Urodyn*. 2014;33(1):95–100.
- Wesseling-Perry K, Salusky IB. Phosphate binders, vitamin D and calcimimetics in the management of chronic kidney disease-mineral bone disorders (CKD-MBD) in children. *Pediatr Nephrol*. 2013;28(4):617–625.

*The definition of oliguria varies extensively in the literature, from 1.8 to 4 dl/m² every 24 hours.

*30 E. 33rd St., New York, NY 10016; 212-889-2210, 800-622-9010;
<http://www.kidney.org>. In Canada: Kidney Foundation of Canada, 300-5165
 Sherbrooke St. West, Montreal, QC H4A 1T6; 514-369-4806, 800-361-7494;
<http://www.kidney.ca>.