

- healthy full-term neonates. *Neonatal Netw.* 2013;32(4):274–280.
- McGowan A, Cottrell S, Roberts R. Minimizing pain response during routine infant immunisation. *Community Practitioner.* 2013 <http://www.readperiodicals.com/201306/3006230571.htm>
- O'Brien L, Taddio A, Ipp M, et al. Topical 4% amethocaine gel reduces the pain of subcutaneous measles-mumps-rubella vaccination. *Pediatrics.* 2004;114(6):e720–e724.
- Racine NM, Pillai Riddell RR, Flora D, et al. A longitudinal examination of verbal reassurance during infant immunization: occurrence and examination of emotional availability as a potential moderator. *J Pediatr Psychol.* 2012;37(8):935–944.
- Saeidi R, Asnaashari Z, Amirnejad M, et al. Use of “kangaroo care” to alleviate the intensity of vaccination pain in newborns. *Iran J Pediatr.* 2011;21(1):99–102.
- Shah PS, Aliwalas L, Shah V. Breastfeeding or breastmilk to alleviate procedural pain in neonates: a systematic review. *Breastfeed Med.* 2007;2(2):74–84.
- Shah VS, Taddio A, Hancock R, et al. Topical amethocaine gel 4% for intramuscular injection in term neonates: a double-blind, placebo-controlled, randomized trial. *Clin Ther.* 2008;30(1):166–174.
- Sparks LA, Setlik J, Luhman J. Parental holding and positioning to decrease IV distress in young children: a randomized controlled trial. *J Pediatr Nurs.* 2007;22(6):440–447.
- Taddio A, Chambers CT, Halperin SA, et al. Inadequate pain management during routine childhood immunizations: the nerve of it. *Clin Ther.* 2009;31(Suppl 2):S152–S167.
- Taddio A, Ilersich AL, Ipp M, et al. Physical interventions and injection techniques for reducing injection pain during routine childhood immunizations: systematic review of randomized controlled trials and quasi-randomized controlled trials. *Clin Ther.* 2009;31(Suppl 2):S48–S76.
- Thyr M, Sundholm A, Teeland L, et al. Oral glucose as an analgesic to reduce infant distress following immunization at the age of 3, 5 and 12 months. *Acta Paediatr.* 2007;96(2):233–236.