

FACTORS FOR UNIT CONVERSIONS

Quantity	Equivalent Values
Mass	1 kg = 1000 g = 0.001 metric ton (tonne) = 2.20462 lb _m = 35.27392 oz 1 lb _m = 16 oz = 5 × 10 ⁻⁴ ton = 453.593 g = 0.453593 kg
Length	1 m = 100 cm = 1000 mm = 10 ⁶ microns (μm) = 10 ¹⁰ angstroms (Å) = 39.37 in = 3.2808 ft = 1.0936 yd = 0.0006214 mile 1 ft = 12 in = 1/3 yd = 0.3048 m = 30.48 cm
Volume	1 m ³ = 1000 L = 10 ⁶ cm ³ = 10 ⁶ mL = 35.3145 ft ³ = 219.97 imperial gallons = 264.17 gal = 1056.68 qt 1 ft ³ = 1728 in ³ = 7.4805 gal = 29.922 qt = 0.028317 m ³ = 28.317 L
Density	1 g/cm ³ = 1000 kg/m ³ = 62.43 lb _m /ft ³ = density of liquid water at 4°C (reference for specific gravities)
Force	1 N = 1 kg·m/s ² = 10 ⁵ dynes = 10 ⁵ g·cm/s ² = 0.22481 lb _f 1 lb _f = 32.174 lb _m ·ft/s ² = 4.4482 N = 4.4482 × 10 ⁵ dynes
Pressure	1 atm = 1.01325 × 10 ⁵ N/m ² (Pa) = 101.325 kPa = 1.01325 bar = 1.01325 × 10 ⁶ dynes/cm ² = 14.696 lb _f /in ² (psi) = 760 mm Hg at 0°C (torr) = 10.333 m H ₂ O(l) at 4°C = 29.921 inches Hg at 0°C = 406.8 inches H ₂ O(l) at 4°C
Energy	1 J = 1 N·m = 10 ⁷ ergs = 10 ⁷ dyne·cm = 1 kg·m ² /s ² = 2.778 × 10 ⁻⁷ kW·h = 0.23901 cal = 0.23901 × 10 ⁻³ kcal (food calorie) = 0.7376 ft·lb _f = 9.486 × 10 ⁻⁴ Btu
Power	1 W = 1 J/s = 1 N·m/s = 0.23901 cal/s = 0.7376 ft·lb _f /s = 9.486 × 10 ⁻⁴ Btu/s = 1.341 × 10 ⁻³ hp

$$s_r(h_r) = \frac{p_i}{p_i^*(T)} \times 100\%$$

$$s_m(h_m) = \frac{p_i}{P-p_i} = \frac{\text{moles of vapor}}{\text{moles of vapor-free (dry) gas}}$$

$$s_a(h_a) = \frac{p_i M_i}{(P-p_i) M_{\text{dry}}} = \frac{\text{mass of vapor}}{\text{mass of dry gas}}$$

$$s_p(h_p) = \frac{s_m}{s_m^*} \times 100\% = \frac{p_i/(P-p_i)}{p_i^*/(P-p_i^*)} \times 100\%$$

Example: The factor to convert grams to lb_m is $\left(\frac{2.20462 \text{ lb}_m}{1000 \text{ g}}\right)$ or $\left(\frac{1 \text{ lb}_m}{453.593 \text{ g}}\right)$.

$$\begin{aligned}\rho_{\text{H}_2\text{O}(l)}(4^\circ\text{C}) &= 1.000 \text{ g/cm}^3 \\ &= 1000. \text{ kg/m}^3 \\ &= 62.43 \text{ lb}_m/\text{ft}^3\end{aligned}$$

Table 5.2-1 Standard Conditions for Gases

System	T_s	P_s	V_s	n_s
SI	273 K	1 atm	0.022415 m ³	1 mol
CGS	273 K	1 atm	22.415 L	1 mol
U.S. customary	492°R	1 atm	359.05 ft ³	1 lb-mole

$$\%XS = \frac{n_i - n_{\text{stoich}}}{n_{\text{stoich}}}$$

$$\frac{1}{\bar{\rho}} = \sum (x_i / \rho_i)$$

$$\bar{\rho} = \sum x_i \rho_i$$

$$\Delta T_b = T_{bs} - T_{b0} = \frac{RT_{b0}^2}{\Delta \bar{H}_v} x$$

$$\Delta T_m = T_{m0} - T_{ms} = \frac{RT_{m0}^2}{\Delta \bar{H}_m} x$$

Table 5.3-1 Pitzer Acentric Factors

(Values with * were in the 4th edition of the reference in Footnote 1.)

Compound	Acentric Factor, ω
Ammonia	0.257
Argon	-0.002
Carbon dioxide	0.225
Carbon monoxide	0.045
Chlorine*	0.073
Ethane	0.099
Ethylene	0.087
Hydrogen sulfide	0.090
Methane	0.011
Methanol	0.565
Nitrogen	0.037
Oxygen*	0.021
Propane	0.152
Sulfur dioxide*	0.251
Water	0.344

ATOMIC WEIGHTS AND NUMBERS

Atomic weights apply to naturally occurring isotopic compositions and are based on an atomic mass of $^{12}\text{C} = 12$

Element	Symbol	Atomic Number	Atomic Weight	Element	Symbol	Atomic Number	Atomic Weight
Actinium	Ac	89	—	Iridium	Ir	77	192.2
Aluminum	Al	13	26.9815	Iron	Fe	26	55.847
Americium	Am	95	—	Krypton	Kr	36	83.80
Antimony	Sb	51	121.75	Lanthanum	La	57	138.91
Argon	Ar	18	39.948	Lawrencium	Lr	103	—
Arsenic	As	33	74.9216	Lead	Pb	82	207.19
Astatine	At	85	—	Lithium	Li	3	6.939
Barium	Ba	56	137.34	Lutetium	Lu	71	174.97
Berkelium	Bk	97	—	Magnesium	Mg	12	24.312
Beryllium	Be	4	9.0122	Manganese	Mn	25	54.9380
Bismuth	Bi	83	208.980	Mendelevium	Md	101	—
Boron	B	5	10.811	Mercury	Hg	80	200.59
Bromine	Br	35	79.904	Molybdenum	Mo	42	95.94
Cadmium	Cd	48	112.40	Neodymium	Nd	60	144.24
Calcium	Ca	20	40.08	Neon	Ne	10	20.183
Californium	Cf	98	—	Neptunium	Np	93	—
Carbon	C	6	12.01115	Nickel	Ni	28	58.71
Cerium	Ce	58	140.12	Niobium	Nb	41	92.906
Cesium	Cs	55	132.905	Nitrogen	N	7	14.0067
Chlorine	Cl	17	35.453	Nobelium	No	102	—
Chromium	Cr	24	51.996	Osmium	Os	75	190.2
Cobalt	Co	27	58.9332	Oxygen	O	8	15.9994
Copper	Cu	29	63.546	Palladium	Pd	46	106.4
Curium	Cm	96	—	Phosphorus	P	15	30.9738
Dysprosium	Dy	66	162.50	Platinum	Pt	78	195.09
Einsteinium	Es	99	—	Plutonium	Pu	94	—
Erbium	Er	68	167.26	Polonium	Po	84	—
Europium	Eu	63	151.96	Potassium	K	19	39.102
Fermium	Fm	100	—	Praseodymium	Pr	59	140.907
Fluorine	F	9	18.9984	Promethium	Pm	61	—
Francium	Fr	87	—	Protactinium	Pa	91	—
Gadolinium	Gd	64	157.25	Radium	Ra	88	—
Gallium	Ga	31	69.72	Radon	Rn	86	—
Germanium	Ge	32	72.59	Rhenium	Re	75	186.2
Gold	Au	79	196.967	Rhodium	Rh	45	102.905
Hafnium	Hf	72	178.49	Rubidium	Rb	37	84.57
Helium	He	2	4.0026	Ruthenium	Ru	44	101.07
Holmium	Ho	67	164.930	Samarium	Sm	62	150.35
Hydrogen	H	1	1.00797	Scandium	Sc	21	44.956
Indium	In	49	114.82	Selenium	Se	34	78.96
Iodine	I	53	126.9044	Silicon	Si	14	28.086

Atomic weights apply to naturally occurring isotopic compositions and are based on an atomic mass of $^{12}\text{C} = 12$

Element	Symbol	Atomic Number	Atomic Weight	Element	Symbol	Atomic Number	Atomic Weight
Silver	Ag	47	107.868	Tin	Sn	50	118.69
Sodium	Na	11	22.9898	Titanium	Ti	22	47.90
Strontium	Sr	38	87.62	Tungsten	W	74	183.85
Sulfur	S	16	32.064	Uranium	U	92	238.03
Tantalum	Ta	73	180.948	Vanadium	V	23	50.942
Technetium	Tc	43	—	Xenon	Xe	54	131.30
Tellurium	Te	52	127.60	Ytterbium	Yb	70	173.04
Terbium	Tb	65	158.924	Yttrium	Y	39	88.905
Thallium	Tl	81	204.37	Zinc	Zn	30	65.37
Thorium	Th	90	232.038	Zirconium	Zr	40	91.22
Thulium	Tm	69	168.934				

THE GAS CONSTANT (R)

$$8.314 \text{ m}^3 \cdot \text{Pa}/(\text{mol} \cdot \text{K})$$

$$0.08314 \text{ L} \cdot \text{bar}/(\text{mol} \cdot \text{K})$$

$$0.08206 \text{ L} \cdot \text{atm}/(\text{mol} \cdot \text{K})$$

$$62.36 \text{ L} \cdot \text{mm Hg}/(\text{mol} \cdot \text{K})$$

$$0.7302 \text{ ft}^3 \cdot \text{atm}/(\text{lb-mole} \cdot ^\circ\text{R})$$

$$10.73 \text{ ft}^3 \cdot \text{psia}/(\text{lb-mole} \cdot ^\circ\text{R})$$

$$8.314 \text{ J}/(\text{mol} \cdot \text{K})$$

$$1.987 \text{ cal}/(\text{mol} \cdot \text{K})$$

$$1.987 \text{ Btu}/(\text{lb-mole} \cdot ^\circ\text{R})$$

$$|\varepsilon| < 1\% \text{ if } \hat{V}_{\text{ideal}} = \frac{RT}{P} > 5 \text{ L/mol (80 ft}^3/\text{lb-mole) (diatomic gases)}$$

$$> 20 \text{ L/mol (320 ft}^3/\text{lb-mole) (other gases)}$$

$$W = mg$$

$$\bar{M} = \sum y_i M_i$$

$$\frac{1}{\bar{M}} = \sum \left(\frac{x_i}{M_i} \right)$$

$$P = P_0 + \rho gh$$

$$P_{\text{abs}} = P_g + P_{\text{atm}}$$

$$P_1 + \rho_1 g d_1 = P_2 + \rho_2 g d_2 + \rho_f g h$$

$$T(\text{K}) = T(^{\circ}\text{C}) + 273.15$$

$$T(^{\circ}\text{R}) = T(^{\circ}\text{F}) + 459.67$$

$$T(^{\circ}\text{R}) = 1.8T(\text{K})$$

$$T(^{\circ}\text{F}) = 1.8T(^{\circ}\text{C}) + 32$$

$$\frac{P\hat{V}}{RT} = 1 + \frac{B}{\hat{V}}$$

$$B_0 = 0.083 - \frac{0.422}{T_r^{1.6}} \quad B_1 = 0.139 - \frac{0.172}{T_r^{4.2}} \quad B = \frac{RT_c}{P_c} (B_0 + \omega B_1)$$

$$T'_c = \sum y_i T_{ci} \quad P'_c = \sum y_i P_{ci}$$