

Techniques: (with Set Cover as an example)

- Solve LP and round solution (Sec. 1.3 + 1.7)
- Primal-dual alg.: combinatorial alg.  
based on LP formulation (Sec. 1.4 + 1.5)
- Greedy alg. (Sec. 1.6)

## Section 1.2: Set Cover as an LP

### Set Cover

Input:

$$E = \{e_1, e_2, \dots, e_n\}$$


$$\mathcal{S} = \{S_1, S_2, \dots, S_m\}, \text{ where}$$

$S_j \subseteq E$  has weight  $w_j$ .

**Objective:** Find a cheapest possible subset of  $\mathcal{S}$  covering all elements

**OPT:** value (total weight) of optimum solution

Ex:



$$w_1 = 1$$

$$w_2 = 2$$

$$w_3 = 3$$

$\{S_1, S_2\}$  is a sol. of total weight 3.

This is optimal, so  $\text{OPT} = 3$  for this instance of Set Cover.

To cover  $e_1$ , we need  $S_1$

—— " ——  $e_2$  —— " ——  $S_1$  or  $S_2$

—— " ——  $e_3$  —— " ——  $S_1, S_2$  or  $S_3$

—— " ——  $e_4$  —— " ——  $S_2$  or  $S_3$

IP-formulation:

$$\min \quad x_1 w_1 + x_2 w_2 + x_3 w_3$$

$$\text{s.t.} \quad x_1 \geq 1$$

$$x_1 + x_2 \geq 1$$

$$x_1 + x_2 + x_3 \geq 1$$

$$x_2 + x_3 \geq 1$$

$$x_1, x_2, x_3 \in \{0, 1\}$$

More generally:

### IP for Set Cover

$$\begin{aligned} \min \quad & \sum_{j=1}^m x_j w_j \\ \text{s.t.} \quad & \sum_{j: e_i \in S_j} x_j \geq 1, \quad i = 1, 2, \dots, n \\ & x_j \in \{0, 1\}, \quad j = 1, 2, \dots, m \end{aligned}$$

$Z_{IP}^*$ : optimum solution value, i.e.,  $Z_{IP}^* = \text{OPT}$

### LP-relaxation

$$\begin{aligned} \min \quad & \sum_{j=1}^m x_j w_j \\ \text{s.t.} \quad & \sum_{j: e_i \in S_j} x_j \geq 1, \quad i = 1, 2, \dots, n \\ & 0 \leq x_j \leq 1, \quad j = 1, 2, \dots, m \end{aligned}$$

redundant

$Z_{LP}^*$ : Optimum solution value

Note that

$$Z_{LP}^* \leq Z_{IP}^* = \text{OPT}$$

## Section 1.3: A deterministic rounding algo.

The frequency of an element  $e$  is the #sets containing  $e$ :

$$f_e = |\{S \in \mathcal{S} \mid e \in S\}|$$

The frequency of an instance of Set Cover:

$$f = \max_{e \in E} \{f_e\}$$

Alg. 1 for Set Cover: LP-rounding

Solve LP

$$I \leftarrow \{j \mid x_j \geq \frac{1}{f}\}$$

We prove that Alg. 1 produces a set cover (Lemma 1.5)  
of total weight  $\leq f \cdot \text{OPT}$  (Thm 1.6)

### Lemma 1.5

$\{S_j \mid j \in I\}$  is a set cover

Proof:

For each  $e_i \in E$ ,  $\sum_{j: e_i \in S_j} x_j \geq 1$ .

Since  $\sum_{j: e_i \in S_j} x_j$  has at most  $f$  terms, at least one of the terms is at least  $\frac{1}{f}$ .

Thus, there is a set  $S_j$  s.t.

$e_i \in S_j$  and  $x_j \geq \frac{1}{f}$ .

This  $j$  is included in  $I$  □

### Thm 1.6

Alg. 1 is an  $f$ -approx. algo. for Set Cover.

Proof:

Correct by Lemma 1.5

Poly, since LP-solving is poly.

Approx. factor  $f$ :

Each  $x_j$  is rounded up to 1, only if it is already at least  $\frac{1}{f}$ .

Thus, each  $x_j$  is multiplied by at most  $f$ , i.e.,

$$\sum_{j \in I} w_j \leq \sum_{j \in I} f \cdot x_j \cdot w_j \leq \sum_{j=1}^m f \cdot x_j \cdot w_j = f \cdot Z_{LP}^* \leq f \cdot \text{OPT}$$

□

The Vertex Cover problem is a special case of Set Cover:

### Vertex Cover


Input:

$$G = (V, E)$$

Objective:

Find a min. card. vertex set  $C \subseteq V$   
s.t. each edge  $e \in E$  has at least one  
endpoint in  $C$ .

Ex:



With  $\mathcal{I} = V$  and  $E = E$ ,

Alg. 1 is a 2-approx. alg. for Vertex Cover.

One of the exercises for Tuesday:  
Write down LP for Vertex Cover.

# Section 1.4: The dual LP

What is a dual?

P

Ex:

$$\min 7x_1 + x_2 + 5x_3$$

$$\text{s.t. } x_1 - x_2 + 3x_3 \geq 10$$

$$5x_1 + 2x_2 - x_3 \geq 6$$

$$x_1, x_2, x_3 \geq 0$$

Primal

$$7x_1 + x_2 + 5x_3 \geq x_1 - x_2 + 3x_3 \geq 10$$

$$\begin{aligned} 7x_1 + x_2 + 5x_3 &\geq x_1 - x_2 + 3x_3 + 5x_1 + 2x_2 - x_3 \\ &\geq 10 + 6 = 16 \end{aligned}$$

$$\begin{aligned} 7x_1 + x_2 + 5x_3 &\geq 2(x_1 - x_2 + 3x_3) + 5x_1 + 2x_2 - x_3 \\ &\geq 2 \cdot 10 + 6 = 26 \end{aligned}$$

To find a largest possible lower bound on  $7x_1 + x_2 + 5x_3$ , we should determine  $y_1$  and  $y_2$  maximizing  $10y_1 + 6y_2$ , under the constraints that

$$\begin{aligned} 7x_1 + x_2 + 5x_3 &\stackrel{(*)}{\geq} \overbrace{y_1(x_1 - x_2 + 3x_3)}^{\geq 10} + \overbrace{y_2(5x_1 + 2x_2 - x_3)}^{\geq 6} \\ &= \underbrace{(y_1 + 5y_2)}_{\leq 7} x_1 + \underbrace{(-y_1 + 2y_2)}_{\leq 1} x_2 + \underbrace{(3y_1 - y_2)}_{\leq 5} x_3 \end{aligned}$$

and  $y_1, y_2, y_3 \geq 0$

otherwise  
" $\geq$ " becomes " $\leq$ "

necessary to satisfy (\*)



Thus, we arrive at the following problem:

D

$$\max \quad 10y_1 + 6y_2$$

$$\text{s.t.} \quad y_1 + 5y_2 \leq 7$$

$$-y_1 + 2y_2 \leq 1$$

$$3y_1 - y_2 \leq 5$$

$$y_1, y_2 \geq 0$$

Dual

In general:

Primal:

$$\min \quad c_1 x_1 + c_2 x_2 + \dots + c_n x_n$$

$$\text{s.t.} \quad a_{i1}x_1 + a_{i2}x_2 + \dots + a_{in}x_n \geq b_i, \quad i = 1, 2, \dots, m$$

$$x_j \geq 0, \quad j = 1, 2, \dots, n$$

Dual:

$$\max \quad b_1 y_1 + b_2 y_2 + \dots + b_m y_m$$


$$\text{s.t.} \quad a_{1j}y_1 + a_{2j}y_2 + \dots + a_{mj}y_m \leq c_j, \quad j = 1, 2, \dots, n$$

$$y_i \geq 0, \quad i = 1, 2, \dots, m$$

Returning to the example above:

The constraints of  $D$  ensure that the value of any sol. to  $D$  is a lower bound on the value of any sol. to  $P$ , i.e., for any pair  $x, y$  of sol. to  $P$  and  $D$  resp.,

$$10y_1 + 6y_2 \leq 7x_1 + x_2 + 5x_3 \quad \text{Weak duality}$$



Consider again the inequality leading to the constraints of the dual:

$$\begin{aligned}
 & \begin{array}{l} \Downarrow \\ y_1 = 0 \vee x_1 - x_2 + 3x_3 = 0 \\ \Downarrow \\ = 10y_1 \end{array} \quad \begin{array}{l} \Downarrow \\ y_2 = 0 \vee 5x_1 + 2x_2 - x_3 = 6 \\ \Downarrow \\ = 6y_2 \end{array} \\
 & \text{(*)} \quad 7x_1 + x_2 + 5x_3 \geq \underbrace{y_1(x_1 - x_2 + 3x_3)}_{\geq 10} + \underbrace{y_2(5x_1 + 2x_2 - x_3)}_{\geq 6} \\
 & = \underbrace{(y_1 + 5y_2)x_1}_{\leq 7} + \underbrace{(-y_1 + 2y_2)x_2}_{\leq 1} + \underbrace{(3y_1 - y_2)x_3}_{\leq 5} \\
 & \begin{array}{l} \Downarrow \\ y_1 + 5y_2 = 7 \\ \vee x_1 = 0 \end{array} \quad \begin{array}{l} \Downarrow \\ -y_1 + 2y_2 = 1 \\ \vee x_2 = 0 \end{array} \quad \begin{array}{l} \Downarrow \\ 3y_1 - y_2 = 5 \\ \vee x_3 = 0 \end{array} \\
 & \text{(*) becomes } =
 \end{aligned}$$

Thus,

$$\begin{array}{l} \Downarrow \\ \text{Complementary} \\ \text{Slackness} \\ \text{Conditions} \end{array} \left\{ \begin{array}{ll} 7x_1 + x_2 + 5x_3 = 10y_1 + 6y_2 \\ x_1 > 0 \Rightarrow y_1 + 5y_2 = 7 \\ x_2 > 0 \Rightarrow -y_1 + 2y_2 = 1 \\ x_3 > 0 \Rightarrow 3y_1 - y_2 = 5 \end{array} \right\} \text{ primal c.s.c.}$$
  
$$\left\{ \begin{array}{ll} y_1 > 0 \Rightarrow x_1 - x_2 + 3x_3 = 10 \\ y_2 > 0 \Rightarrow 5x_1 + 2x_2 - x_3 = 6 \end{array} \right\} \text{ dual c.s.c.}$$

By The **Strong Duality Theorem** (which we will not prove), there exist solutions fulfilling the c.s.c.

Moreover, if the c.s.c. are „close“ to being satisfied, the values of the primal and dual sol. are „close“ :

$$\begin{array}{l}
 \text{Relaxed} \\
 \text{Complementary} \\
 \text{Slackness} \\
 \text{Conditions}
 \end{array}
 \left\{
 \begin{array}{ll}
 x_1 > 0 & \Rightarrow y_1 + 5y_2 \geq 7/b \\
 x_2 > 0 & \Rightarrow -y_1 + 2y_2 \geq 1/b \\
 x_3 > 0 & \Rightarrow 3y_1 - y_2 \geq 5/b \\
 y_1 > 0 & \Rightarrow x_1 - x_2 + 3x_3 \leq 10c \\
 y_2 > 0 & \Rightarrow 5x_1 + 2x_2 - x_3 \leq 6c
 \end{array}
 \right.$$

$$\Downarrow$$

$$7x_1 + x_2 + 5x_3 \leq bc(10y_1 + 6y_2)$$

Proof:


$$\begin{aligned}
 & (y_1 + 5y_2)x_1 + (-y_1 + 2y_2)x_2 + (3y_1 - y_2)x_3 \\
 & \geq \frac{7}{b}x_1 + \frac{1}{b}x_2 + \frac{5}{b}x_3, \text{ by the Primal relaxed c.s.c.} \\
 & = \frac{1}{b}(7x_1 + x_2 + 5x_3)
 \end{aligned}$$

$\Downarrow$

$$\begin{aligned}
 7x_1 + x_2 + 5x_3 & \leq b((y_1 + 5y_2)x_1 + (-y_1 + 2y_2)x_2 + (3y_1 - y_2)x_3) \\
 & = b((x_1 - x_2 + 3x_3)y_1 + (5x_1 + 2x_2 - x_3)y_2) \\
 & \leq b(10cy_1 + 6cy_2), \text{ by the Dual r.c.s.c.} \\
 & = bc(10y_1 + 6y_2)
 \end{aligned}$$

What is the dual of the Set Cover LP?

Ex:



$$w_1 = 1$$

$$w_2 = 2$$

$$w_3 = 3$$

Primal:

$$\min \quad x_1 + 2x_2 + 3x_3$$

$$\text{s.t.} \quad x_1 \geq 1$$

$$x_1 + x_2 \geq 1$$

$$x_1 + x_3 \geq 1$$

$$x_2 + x_3 \geq 1$$

$$x_1, x_2, x_3 \geq 0$$

$$\text{OPT} = 3:$$

$$x_1 = x_2 = 1$$

Dual:

$$\max \quad y_1 + y_2 + y_3 + y_4$$

$$\text{s.t.} \quad y_1 + y_2 + y_3 \leq 1$$

$$y_2 + y_4 \leq 2$$

$$y_3 + y_4 \leq 3$$

$$y_1, y_2, y_3, y_4 \geq 0$$

$$\text{OPT} = 3:$$

$$y_1 = 1$$

$$y_4 = 2$$

or

$$y_3 = 1$$

$$y_4 = 2$$

### Set Cover Primal

$$\begin{aligned} \min \quad & \sum_{j=1}^m x_j w_j \\ \text{s.t.} \quad & \sum_{j: e_i \in S_j} x_j \geq 1, \quad i=1, 2, \dots, n \\ & x_j \geq 0, \quad j=1, 2, \dots, m \end{aligned}$$

Covering  
problem

### Set Cover Dual

$$\begin{aligned} \max \quad & \sum_{i=1}^n y_i \\ \text{s.t.} \quad & \sum_{e_i \in S_j} y_i \leq w_j, \quad j=1, 2, \dots, m \\ & y_i \geq 0, \quad i=1, 2, \dots, n \end{aligned}$$

Packing  
problem

Recall that the dual is constructed such that the value of any solution to the dual is a lower bound on the value of any solution to the primal:

$$Z_{\text{Primal}} \geq Z_{\text{Dual}} \quad (\text{weak duality property})$$

In fact,

$$Z_{\text{Primal}}^* = Z_{\text{Dual}}^* \quad (\text{strong duality property})$$