

DM545/DM871
Linear and Integer Programming

More on Duality

Marco Chiarandini

Department of Mathematics and Computer Science

October 5, 2026

Outline

Derivation
Dual Simplex
Sensitivity Analysis

1. Derivation

Lagrangian Duality

2. Dual Simplex

3. Sensitivity Analysis

- Derivation:
 1. economic interpretation
 2. bounding
 3. multipliers
 4. recipe
 5. Lagrangian
- Theory:
 - Symmetry
 - Weak duality theorem
 - Strong duality theorem
 - Complementary slackness theorem
- Economic interpretation
- Dual Simplex
- Sensitivity Analysis

Outline

Derivation
Dual Simplex
Sensitivity Analysis

1. Derivation

Lagrangian Duality

2. Dual Simplex

3. Sensitivity Analysis

Outline

Derivation
Dual Simplex
Sensitivity Analysis

1. Derivation

Lagrangian Duality

2. Dual Simplex

3. Sensitivity Analysis

Lagrangian Duality

Relaxation: if a problem is hard to solve then find an easier problem resembling the original one that provides information in terms of bounds. Then, search for the strongest bounds.

$$\begin{aligned} \min \quad & 13x_1 + 6x_2 + 4x_3 + 12x_4 \\ & 2x_1 + 3x_2 + 4x_3 + 5x_4 = 7 \\ & 3x_1 + \quad + 2x_3 + 4x_4 = 2 \\ & x_1, x_2, x_3, x_4 \geq 0 \end{aligned}$$

We wish to reduce to a problem easier to solve, ie:

$$\begin{aligned} \min \quad & c_1x_1 + c_2x_2 + \dots + c_nx_n \\ & x_1, x_2, \dots, x_n \geq 0 \end{aligned}$$

solvable by inspection: if $c_j < 0$ then $x_j = +\infty$, if $c_j \geq 0$ then $x_j = 0$.

Measure of violation of the constraints:

$$\begin{aligned} 7 - (2x_1 + 3x_2 + 4x_3 + 5x_4) \\ 2 - (3x_1 + \quad + 2x_3 + 4x_4) \end{aligned}$$

We relax these measures in obj. function with **Lagrangian multipliers** y_1, y_2 .

We obtain a family of problems:

$$PR(y_1, y_2) = \min_{x_1, x_2, x_3, x_4 \geq 0} \left\{ \begin{array}{l} 13x_1 + 6x_2 + 4x_3 + 12x_4 \\ + y_1(7 - 2x_1 - 3x_2 - 4x_3 - 5x_4) \\ + y_2(2 - 3x_1 - 2x_3 - 4x_4) \end{array} \right\}$$

1. for all $y_1, y_2 \in \mathbb{R} : \text{opt}(PR(y_1, y_2)) \leq \text{opt}(P)$
2. $\max_{y_1, y_2 \in \mathbb{R}} \{\text{opt}(PR(y_1, y_2))\} \leq \text{opt}(P)$

PR is easy to solve.

(It can be also seen as a proof of the weak duality theorem)

$$PR(y_1, y_2) = \min_{x_1, x_2, x_3, x_4 \geq 0} \left\{ \begin{array}{l} (13 - 2y_1 - 3y_2) x_1 \\ + (6 - 3y_1) x_2 \\ + (4 - 4y_1 - 2y_2) x_3 \\ + (12 - 5y_1 - 4y_2) x_4 \\ + 7y_1 + 2y_2 \end{array} \right\}$$

if coefficient of x is < 0 then bound is $-\infty$ then LB is useless

$$\begin{aligned} (13 - 2y_1 - 3y_2) &\geq 0 \\ (6 - 3y_1) &\geq 0 \\ (4 - 4y_1 - 2y_2) &\geq 0 \\ (12 - 5y_1 - 4y_2) &\geq 0 \end{aligned}$$

If they all hold then we are left with $7y_1 + 2y_2$ because all go to 0.

$$\begin{aligned} \max \quad & 7y_1 + 2y_2 \\ & 2y_1 + 3y_2 \leq 13 \\ & 3y_1 \leq 6 \\ & 4y_1 + 2y_2 \leq 4 \\ & 5y_1 + 4y_2 \leq 12 \end{aligned}$$

General Formulation

$$\begin{array}{ll}\min & z = \mathbf{c}^T \mathbf{x} & \mathbf{c} \in \mathbb{R}^n \\ & A\mathbf{x} = \mathbf{b} & A \in \mathbb{R}^{m \times n}, \mathbf{b} \in \mathbb{R}^m \\ & \mathbf{x} \geq \mathbf{0} & \mathbf{x} \in \mathbb{R}^n\end{array}$$

$$\max_{\mathbf{y} \in \mathbb{R}^m} \left\{ \min_{\mathbf{x} \in \mathbb{R}_+^n} \{ \mathbf{c}^T \mathbf{x} + \mathbf{y}^T (\mathbf{b} - A\mathbf{x}) \} \right\}$$

$$\max_{\mathbf{y} \in \mathbb{R}^m} \left\{ \min_{\mathbf{x} \in \mathbb{R}_+^n} \{ (\mathbf{c}^T - \mathbf{y}^T A) \mathbf{x} + \mathbf{y}^T \mathbf{b} \} \right\}$$

$$\begin{array}{ll}\max & \mathbf{b}^T \mathbf{y} \\ & A^T \mathbf{y} \leq \mathbf{c} \\ & \mathbf{y} \in \mathbb{R}^m\end{array}$$

Outline

Derivation
Dual Simplex
Sensitivity Analysis

1. Derivation

Lagrangian Duality

2. Dual Simplex

3. Sensitivity Analysis

- Dual simplex (Lemke, 1954): apply the simplex method to the dual problem and observe what happens in the primal tableau:

$$\begin{aligned}\max\{\mathbf{c}^T \mathbf{x} \mid A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\} &= \min\{\mathbf{b}^T \mathbf{y} \mid A^T \mathbf{y} \geq \mathbf{c}^T, \mathbf{y} \geq \mathbf{0}\} \\ &= -\max\{-\mathbf{b}^T \mathbf{y} \mid -A^T \mathbf{y} \leq -\mathbf{c}^T, \mathbf{y} \geq \mathbf{0}\}\end{aligned}$$

- We obtain a new algorithm for the primal problem: the **dual simplex**
It corresponds to the primal simplex applied to the dual
- We obtain also another way to handle infeasibility

Primal Simplex on Dual Problem

Example: Primal feasible start

Derivation
Dual Simplex
Sensitivity Analysis

Primal:

$$\begin{aligned} \max \quad & 4x_1 + x_2 + 3x_3 \\ & x_1 + 4x_2 \leq 1 \\ & 3x_1 + x_2 + x_3 \leq 3 \\ & x_1, x_2, x_3 \geq 0 \end{aligned}$$

- Initial tableau

x1	x2	x3	w1	w2	-z	b
1	4	0	1	0	0	1
3	1	1	0	1	0	3
4	1	3	0	0	1	0

- primal feasible start

Dual:

$$\begin{aligned} \min \quad & y_1 + 3y_2 \\ & y_1 + 3y_2 \geq 4 \\ & 4y_1 + y_2 \geq 1 \\ & y_2 \geq 3 \\ & y_1, y_2 \geq 0 \end{aligned}$$

- Initial tableau ($\min by \equiv -\max -by$)

y1	y2	z1	z2	z3	-p	b
-1	-3	1	0	0	0	-4
-4	-1	0	0	1	0	-1
0	-1	0	0	1	0	-3
-1	-3	0	0	0	1	0

- dual infeasible start

Primal Simplex on Dual Problem

Example: Primal infeasible start

Derivation
Dual Simplex
Sensitivity Analysis

Primal:

$$\begin{aligned} \max \quad & -x_1 - x_2 \\ & -2x_1 - x_2 \leq 4 \\ & -2x_1 + 4x_2 \leq -8 \\ & -x_1 + 3x_2 \leq -7 \\ & x_1, x_2 \geq 0 \end{aligned}$$

- Initial tableau

	x1	x2	w1	w2	w3	-z	b
----	----	----	----	----	----	----	----
	-2	-1	1	0	0	0	4
	-2	4	0	1	0	0	-8
	-1	3	0	0	1	0	-7
----	----	----	----	----	----	----	----
	-1	-1	0	0	0	1	0

infeasible start

- x_1 enters, w_2 leaves

Dual:

$$\begin{aligned} \min \quad & 4y_1 - 8y_2 - 7y_3 \\ & -2y_1 - 2y_2 - y_3 \geq -1 \\ & -y_1 + 4y_2 + 3y_3 \geq -1 \\ & y_1, y_2, y_3 \geq 0 \end{aligned}$$

- Initial tableau ($\min by \equiv -\max -by$)

	y1	y2	y3	z1	z2	-p	b
----	----	----	----	----	----	----	----
	2	2	1	1	0	0	1
	1	-4	-3	0	1	0	1
----	----	----	----	----	----	----	----
	-4	8	7	0	0	1	0

feasible start (thanks to $-x_1 - x_2$)

- y_2 enters, z_1 leaves

- x_1 enters, w_2 leaves

	x_1	x_2	w_1	w_2	w_3	$-z$	b
	0	-5	1	-1	0	0	12
	1	-2	0	-0.5	0	0	4
	0	1	0	-0.5	1	0	-3
	0	-3	0	-0.5	0	1	4

- w_2 enters, w_3 leaves

	x_1	x_2	w_1	w_2	w_3	$-z$	b
	0	-7	1	0	-2	0	18
	1	-3	0	0	-1	0	7
	0	-2	0	1	-2	0	6
	0	-4	0	0	-1	1	7

(note that we kept $c_j < 0$, ie, optimality)

- y_2 enters, z_1 leaves

	y_1	y_2	y_3	z_1	z_2	$-p$	b
	1	1	0.5	0.5	0	0	0.5
	5	0	-1	2	1	0	3
	-4	0	3	-12	0	1	-4

- y_3 enters, y_2 leaves

	y_1	y_2	y_3	z_1	z_2	$-p$	b
	2	2	1	1	0	0	1
	7	2	0	3	1	0	3
	-18	-6	0	-7	0	1	-7

Dual Simplex on Primal Problem

Primal simplex on primal problem:

1. pivot > 0
2. col c_j with wrong sign
3. row: $\min \left\{ \frac{b_i}{a_{ij}} : a_{ij} > 0, i = 1, \dots, m \right\}$

Dual simplex on primal problem:

1. pivot < 0
2. row $b_i < 0$
(condition of feasibility)
3. col: $\min \left\{ \left| \frac{c_j}{a_{ij}} \right| : a_{ij} < 0, j = 1, 2, \dots, n + m \right\}$
(least worsening solution)

- Primal works with feasible solutions towards optimality
- Dual works with optimal solutions towards feasibility

1. (primal) simplex on primal problem (the one studied so far)
2. Now: dual simplex on primal problem $\equiv$ primal simplex on dual problem
(implemented as dual simplex, understood as primal simplex on dual problem)

Uses of the Dual Simplex:

- Infeasible start
Dual based Phase I algorithm (**Dual-primal algorithm**)
- The dual simplex can work better than the primal in some cases.
Eg. since running time in practice between $2m$ and $3m$, then if $m = 99$ and $n = 9$ then better the dual

Dual based Phase I

Example:

$$\begin{array}{ll}\text{maximize} & z = x_1 - x_2 \\ \text{subject to} & x_1 + x_2 \leq 2 \\ & 2x_1 + 2x_2 \geq 2 \\ & x_1, x_2 \geq 0\end{array}$$

- Derivation:
 1. bounding
 2. multipliers
 3. recipe
 4. Lagrangian
- Theory:
 - Symmetry
 - Weak duality theorem
 - Strong duality theorem
 - Complementary slackness theorem
- Dual Simplex
- Sensitivity Analysis, Economic interpretation

Outline

Derivation
Dual Simplex
Sensitivity Analysis

1. Derivation

Lagrangian Duality

2. Dual Simplex

3. Sensitivity Analysis

Instead of solving each modified problem from scratch, exploit results obtained from solving the original problem.

$$\max\{\mathbf{c}^T \mathbf{x} \mid A\mathbf{x} = \mathbf{b}, \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}\} \quad (*)$$

(I) changes to coefficients of objective function: $\max\{\tilde{\mathbf{c}}^T \mathbf{x} \mid A\mathbf{x} = \mathbf{b}, \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}\}$ (primal)
 $\mathbf{x}^*$ of (*) remains feasible hence we can restart the simplex from $\mathbf{x}^*$

(II) changes to RHS terms: $\max\{\mathbf{c}^T \mathbf{x} \mid A\mathbf{x} = \tilde{\mathbf{b}}, \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}\}$ (dual)
 $\mathbf{x}^*$ optimal feasible solution of (*)

basic sol $\bar{\mathbf{x}}$ of (II): $\bar{\mathbf{x}}_N = \mathbf{x}_N^*, A_B \bar{\mathbf{x}}_B = \tilde{\mathbf{b}} - A_N \bar{\mathbf{x}}_N$

$\bar{\mathbf{x}}$ is dual feasible and we can start the dual simplex from there. If $\tilde{\mathbf{b}}$ differs from $\mathbf{b}$ only slightly it may be we are already optimal.

(III) introduce a new variable:

(primal)

$$\begin{aligned} \max \quad & \sum_{j=1}^6 c_j x_j \\ & \sum_{j=1}^6 a_{ij} x_j = b_i, \quad i = 1, \dots, 3 \\ & l_j \leq x_j \leq u_j, \quad j = 1, \dots, 6 \\ & [x_1^*, \dots, x_6^*] \text{ feasible} \end{aligned}$$

$$\begin{aligned} \max \quad & \sum_{j=1}^7 c_j x_j \\ & \sum_{j=1}^7 a_{ij} x_j = b_i, \quad i = 1, \dots, 3 \\ & l_j \leq x_j \leq u_j, \quad j = 1, \dots, 7 \\ & [x_1^*, \dots, x_6^*, 0] \text{ feasible} \end{aligned}$$

(IV) introduce a new constraint:

(dual)

$$\begin{aligned} & \sum_{j=1}^6 a_{4j} x_j = b_4 \\ & \sum_{j=1}^6 a_{5j} x_j = b_5 \\ & l_j \leq x_j \leq u_j \quad j = 7, 8 \end{aligned}$$

$$\begin{aligned} & [x_1^*, \dots, x_6^*] \text{ optimal} \\ & [x_1^*, \dots, x_6^*, x_7^*, x_8^*] \text{ dual feasible} \\ & x_7^* = b_4 - \sum_{j=1}^6 a_{4j} x_j^* \\ & x_8^* = b_5 - \sum_{j=1}^6 a_{5j} x_j^* \end{aligned}$$

Examples

(I) Variation of reduced costs:

$$\begin{aligned} \max \quad & 6x_1 + 8x_2 \\ & 5x_1 + 10x_2 \leq 60 \\ & 4x_1 + 4x_2 \leq 40 \\ & x_1, x_2 \geq 0 \end{aligned}$$

	x_1	x_2	x_3	x_4	$-z$	b
x_3	5	10	1	0	0	60
x_4	4	4	0	1	0	40
	6	8	0	0	1	0

The last tableau gives the possibility to estimate the effect of variations

	x_1	x_2	x_3	x_4	$-z$	b
x_2	0	1	$1/5$	$-1/4$	0	2
x_1	1	0	$-1/5$	$1/2$	0	8
	0	0	$-2/5$	-1	1	-64

- For a variable in basis the perturbation goes unchanged in the red. costs. Eg:

$$\max (6 + \delta)x_1 + 8x_2 \implies \bar{c}_1 = 1(6 + \delta) - \frac{2}{5} \cdot 5 - 1 \cdot 4 = \delta$$

then need to bring in canonical form and hence δ changes the obj value.

- For a variable not in basis, if it changes the sign of the reduced cost $\implies$ worth bringing in basis $\implies$ the δ term propagates to other columns

(II) Changes in RHS terms

	x_1	x_2	x_3	x_4	$-z$	b
x_3	5	10	1	0	0	$60 + \delta$
x_4	4	4	0	1	0	$40 + \epsilon$
	6	8	0	0	1	0

	x_1	x_2	x_3	x_4	$-z$	b
x_2	0	1	$1/5$	$-1/4$	0	$2 + 1/5\delta - 1/4\epsilon$
x_1	1	0	$-1/5$	$1/2$	0	$8 - 1/5\delta + 1/2\epsilon$
	0	0	$-2/5$	-1	1	$-64 - 2/5\delta - \epsilon$

(It would be more convenient to augment the second. But let's take $\epsilon = 0$.)

If $60 + \delta \implies$ all RHS terms change and we must check feasibility

Which are the multipliers for the first row? $k_1 = \frac{1}{5}, k_2 = -\frac{1}{4}, k_3 = 0$


I: $1/5(60 + \delta) - 1/4 \cdot 40 + 0 \cdot 0 = 12 + \delta/5 - 10 = 2 + \delta/5$

II: $-1/5(60 + \delta) + 1/2 \cdot 40 + 0 \cdot 0 = -60/5 + 20 - \delta/5 = 8 - 1/5\delta$

Risk that RHS becomes negative

Eg: if $\delta = -10 \implies$ tableau stays optimal but not feasible $\implies$ apply dual simplex

Graphical Representation



(III) Add a variable

$$\begin{aligned} \max \quad & 5x_0 + 6x_1 + 8x_2 \\ & 6x_0 + 5x_1 + 10x_2 \leq 60 \\ & 8x_0 + 4x_1 + 4x_2 \leq 40 \\ & x_0, x_1, x_2 \geq 0 \end{aligned}$$

Reduced cost of x_0 ? $c_j + \sum \pi_i a_{ij} = +1 \cdot 5 - \frac{2}{5} \cdot 6 + (-1)8 = -\frac{27}{5}$

To make worth entering in basis:

- increase its cost
- decrease the technological coefficient in constraint II: $5 - 2/5 \cdot 6 - a_{20} > 0$

(IV) Add a constraint

$$\begin{aligned}
 \max \quad & 6x_1 + 8x_2 \\
 & 5x_1 + 10x_2 \leq 60 \\
 & 4x_1 + 4x_2 \leq 40 \\
 & 5x_1 + 6x_2 \leq 50 \\
 & x_1, x_2 \geq 0
 \end{aligned}$$

Final tableau not in canonical form, need to iterate with dual simplex

	x_1	x_2	x_3	x_4	x_5	$-z$	b
x_2	0	1	$1/5$	$-1/4$		0	2
x_1	1	0	$-1/5$	$1/2$		0	8
	0	0	$-1/5$	-1	1	0	-2
	0	0	$-2/5$	-1	0	1	-64

(V) change in a technological coefficient:

	x_1	x_2	x_3	x_4	$-z$	b
x_3	5	$10 + \delta$	1	0	0	60
x_4	4	4	0	1	0	40
	6	8	0	0	1	0

- first effect on its column
- then look at c
- finally look at b

	x_1	x_2	x_3	x_4	$-z$	b
x_2	0	$(10 + \delta)1/5 + 4(-1/4)$	$1/5$	$-1/4$	0	2
x_1	1	$(10 + \delta)(-1/5) + 4(1/2)$	$-1/5$	$1/2$	0	8
	0	$-2/5\delta$	$-2/5$	-1	1	-6δ

Relevance of Sensitivity Analysis

- The dominant application of LP is mixed integer linear programming.
- In this context it is extremely important being able to begin with a model instantiated in one form followed by a sequence of problem modifications
 - row and column additions and deletions,
 - variable fixingsinterspersed with resolves

- Derivation:
 1. economic interpretation
 2. bounding
 3. multipliers
 4. recipe
 5. Lagrangian
- Theory:
 - Symmetry
 - Weak duality theorem
 - Strong duality theorem
 - Complementary slackness theorem
- Economic interpretation
- Dual Simplex
- Sensitivity Analysis