

DM545/DM871
Linear and Integer Programming

Duality

Marco Chiarandini

Department of Mathematics and Computer Science

September 28, 2026

1. Derivation and Motivation

2. Theory

1. Derivation and Motivation

2. Theory

Dual variables $\mathbf{y}$ in one-to-one correspondence with the constraints:

Primal problem:

$$\begin{aligned}\max \quad & z = \mathbf{c}^T \mathbf{x} \\ & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq \mathbf{0}\end{aligned}$$

Dual Problem:

$$\begin{aligned}\min \quad & w = \mathbf{b}^T \mathbf{y} \\ & A^T \mathbf{y} \geq \mathbf{c} \\ & \mathbf{y} \geq \mathbf{0}\end{aligned}$$

Bounding approach

$$\begin{aligned} z^* = \max \quad & 4x_1 + x_2 + 3x_3 \\ & x_1 + 4x_2 \leq 1 \\ & 3x_1 + x_2 + x_3 \leq 3 \\ & x_1, x_2, x_3 \geq 0 \end{aligned}$$

a feasible solution is a **lower bound** but how good?

By tentatives:

$$(x_1, x_2, x_3) = (1, 0, 0) \rightsquigarrow z^* \geq 4$$

$$(x_1, x_2, x_3) = (0, 0, 3) \rightsquigarrow z^* \geq 9$$

What about **upper bounds**?

$$\begin{array}{rcl} 2 \cdot (x_1 + 4x_2) & \leq & 2 \cdot 1 \\ + 3 \cdot (3x_1 + x_2 + x_3) & \leq & 3 \cdot 3 \\ \hline 4x_1 + x_2 + 3x_3 & \leq & 11x_1 + 11x_2 + 3x_3 \leq 11 \end{array}$$
$$\mathbf{c}^T \mathbf{x} \leq \mathbf{y}^T A \mathbf{x} \leq \mathbf{y}^T \mathbf{b}$$

Hence $z^* \leq 11$. Is this the best upper bound we can find?

multipliers $y_1, y_2 \geq 0$ that preserve sign of inequality

$$\begin{array}{rcl} y_1 \cdot (x_1 + 4x_2) & \leq & y_1(1) \\ y_2 \cdot (3x_1 + x_2 + x_3) & \leq & y_2(3) \\ \hline (y_1 + 3y_2)x_1 + (4y_1 + y_2)x_2 + y_2x_3 & \leq & y_1 + 3y_2 \end{array}$$

Coefficients

$$\begin{array}{rcl} y_1 + 3y_2 & \geq & 4 \\ 4y_1 + y_2 & \geq & 1 \\ y_2 & \geq & 3 \end{array}$$

$z = 4x_1 + x_2 + 3x_3 \leq (y_1 + 3y_2)x_1 + (4y_1 + y_2)x_2 + y_2x_3 \leq y_1 + 3y_2$
then to attain the best upper bound:

$$\begin{array}{rcl} \min & y_1 + 3y_2 \\ & y_1 + 3y_2 \geq 4 \\ & 4y_1 + y_2 \geq 1 \\ & y_2 \geq 3 \\ & y_1, y_2 \geq 0 \end{array}$$

$$\begin{array}{rcl}
 -z = & \pi_1 b_1 & + \pi_2 b_2 \dots + \pi_m b_m \\
 & \pi_1 a_{11} & + \pi_2 a_{21} \dots + \pi_m a_{m1} \leq -c_1 \\
 & \vdots & \ddots \qquad \qquad \qquad \vdots \\
 & \pi_1 a_{1n} & + \pi_2 a_{2n} \dots + \pi_m a_{mn} \leq -c_n \\
 & & \pi_1, \pi_2, \dots, \pi_m \leq 0
 \end{array}$$

$$\mathbf{y} = -\boldsymbol{\pi}$$

$$\begin{array}{rcl}
 -z = & (-y_1 b_1) & + (-y_2 b_2) \dots + (-y_m b_m) \\
 & (-y_1 a_{11}) & + (-y_2 a_{21}) \dots + (-y_m a_{m1}) \leq -c_1 \\
 & \vdots & \ddots \qquad \qquad \qquad \vdots \\
 & (-y_1 a_{1n}) & + (-y_2 a_{2n}) \dots + (-y_m a_{mn}) \leq -c_n \\
 & & -y_1, -y_2, \dots, -y_m \leq 0
 \end{array}$$

as we will see $\mathbf{b}^T \mathbf{y} \geq \mathbf{c}^T \mathbf{x}$, hence it is more interesting to minimize. It yields:

$$\begin{array}{ll}
 \min & \mathbf{b}^T \mathbf{y} \\
 & A^T \mathbf{y} \geq \mathbf{c} \\
 & \mathbf{y} \geq \mathbf{0}
 \end{array}$$

Example

$$\begin{array}{ll}\max & 6x_1 + 8x_2 \\ & 5x_1 + 10x_2 \leq 60 \\ & 4x_1 + 4x_2 \leq 40 \\ & x_1, x_2 \geq 0\end{array}$$

$$\left\{ \begin{array}{lll} 5\pi_1 + 4\pi_2 + 6\pi_3 \leq 0 \\ 10\pi_1 + 4\pi_2 + 8\pi_3 \leq 0 \\ 1\pi_1 + 0\pi_2 + 0\pi_3 \leq 0 \\ 0\pi_1 + 1\pi_2 + 0\pi_3 \leq 0 \\ 0\pi_1 + 0\pi_2 + 1\pi_3 = 1 \\ 60\pi_1 + 40\pi_2 \end{array} \right.$$

$$y_1 = -\pi_1 \geq 0$$

$$y_2 = -\pi_2 \geq 0$$

Duality Recipe

	Primal linear program	Dual linear program
Variables	$x_1, x_2, \dots, x_n$	$y_1, y_2, \dots, y_m$
Matrix	A	A^T
Right-hand side	$\mathbf{b}$	$\mathbf{c}$
Objective function	$\max \mathbf{c}^T \mathbf{x}$	$\min \mathbf{b}^T \mathbf{y}$
Constraints	i th constraint has $\leq$ $\geq$ $=$	$y_i \geq 0$ $y_i \leq 0$ $y_i \in \mathbb{R}$
	$x_j \geq 0$ $x_j \leq 0$ $x_j \in \mathbb{R}$	j th constraint has $\geq$ $\leq$ $=$

1. Derivation and Motivation

2. Theory

The dual of the dual is the primal:

Primal Problem:

$$\begin{aligned}\max \quad & z = c^T x \\ & Ax \leq b \\ & x \geq 0\end{aligned}$$

Let's put the dual in the standard form

Dual Problem:

$$\begin{aligned}\min \quad & b^T y \equiv -\max -b^T y \\ & -A^T y \leq -c \\ & y \geq 0\end{aligned}$$

Dual Problem:

$$\begin{aligned}\min \quad & w = b^T y \\ & A^T y \geq c \\ & y \geq 0\end{aligned}$$

Dual of Dual:

$$\begin{aligned}-\min \quad & -c^T x \\ & -Ax \geq -b \\ & x \geq 0\end{aligned}$$

Weak Duality Theorem

As we saw the dual produces upper bounds. This is true in general:

Theorem (Weak Duality Theorem)

Given:

$$(P) \max\{\mathbf{c}^T \mathbf{x} \mid A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$$

$$(D) \min\{\mathbf{b}^T \mathbf{y} \mid A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$$

for any feasible solution $\mathbf{x}$ of (P) and any feasible solution $\mathbf{y}$ of (D):

$$\mathbf{c}^T \mathbf{x} \leq \mathbf{b}^T \mathbf{y}$$

Proof:

From (D) $c_j \leq \sum_{i=1}^m y_i a_{ij} \forall j$ and from (P) $\sum_{j=1}^n a_{ij} x_j \leq b_i \forall i$

From (D) $y_i \geq 0$ and from (P) $x_j \geq 0$

$$\sum_{j=1}^n c_j x_j \leq \sum_{j=1}^n \left(\sum_{i=1}^m y_i a_{ij} \right) x_j = \sum_{i=1}^m \left(\sum_{j=1}^n a_{ij} x_j \right) y_i \leq \sum_{i=1}^m b_i y_i$$

Strong Duality Theorem

Due to Von Neumann and Dantzig 1947 and Gale, Kuhn and Tucker 1951.

Theorem (Strong Duality Theorem)

Given:

$$(P) \max\{\mathbf{c}^T \mathbf{x} \mid A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$$

$$(D) \min\{\mathbf{b}^T \mathbf{y} \mid A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$$

exactly one of the following occurs:

1. *(P) and (D) are both infeasible*
2. *(P) is unbounded and (D) is infeasible*
3. *(P) is infeasible and (D) is unbounded*
4. *(P) has feasible solution, then let an optimal be: $\mathbf{x}^* = [x_1^*, \dots, x_n^*]$
(D) has feasible solution, then let an optimal be: $\mathbf{y}^* = [y_1^*, \dots, y_m^*]$, then:*

$$\mathbf{c}^T \mathbf{x}^* = \mathbf{b}^T \mathbf{y}^*$$

Proof:

- all other combinations of 3 possibilities (Optimal, Infeasible, Unbounded) for (P) and 3 for (D) are ruled out by weak duality theorem.
- we use the simplex method. (Other proofs independent of the simplex method exist, eg, Farkas Lemma and convex polyhedral analysis)
- The last row of the final tableau will give us that for every $\mathbf{x}$:

$$\begin{aligned} z &= z^* + \sum_{k=1}^{n+m} \bar{c}_k x_k = z^* + \sum_{j=1}^n \bar{c}_j x_j + \sum_{i=1}^m \bar{c}_{n+i} x_{n+i} \\ &= z^* + \bar{\mathbf{c}}_B \mathbf{x}_B + \bar{\mathbf{c}}_N \mathbf{x}_N \end{aligned} \quad (*)$$

In addition, $z^* = \sum_{j=1}^n c_j x_j^*$ (c_j , original values) because optimal value

- We define $y_i^* = -\bar{c}_{n+i}, \quad i = 1, 2, \dots, m$

- We claim that $(y_1^*, y_2^*, \dots, y_m^*)$ is a dual feasible solution satisfying $\mathbf{c}^T \mathbf{x}^* = \mathbf{b}^T \mathbf{y}^*$.

- Let's verify the claim:

We substitute in (*): i) $z = \sum_{j=1}^n c_j x_j$; ii) $\bar{c}_{n+i} = -y_i^*$; and iii) $x_{n+i} = b_i - \sum_{j=1}^n a_{ij} x_j$ for $i = 1, 2, \dots, m$ ($n+i$ are the slack variables)

$$\begin{aligned} \sum_{j=1}^n c_j x_j &= z^* + \sum_{j=1}^n \bar{c}_j x_j - \sum_{i=1}^m y_i^* \left(b_i - \sum_{j=1}^n a_{ij} x_j \right) \\ &= \left(z^* - \sum_{i=1}^m y_i^* b_i \right) + \sum_{j=1}^n \left(\bar{c}_j + \sum_{i=1}^m a_{ij} y_i^* \right) x_j \end{aligned}$$

This must hold for every $(x_1, x_2, \dots, x_n)$ hence:

$$z^* = \sum_{i=1}^m b_i y_i^* \quad \implies y^* \text{ satisfies } c^T x^* = b^T y^*$$

$$c_j = \bar{c}_j + \sum_{i=1}^m a_{ij} y_i^*, j = 1, 2, \dots, n$$

Since $\bar{c}_k \leq 0$ for every $k = 1, 2, \dots, n + m$:

$$\bar{c}_j \leq 0 \rightsquigarrow c_j - \sum_{i=1}^m y_i^* a_{ij} \leq 0 \rightsquigarrow$$

$$\sum_{i=1}^m y_i^* a_{ij} \geq c_j \quad j = 1, 2, \dots, n$$

$$\bar{c}_{n+i} \leq 0 \rightsquigarrow y_i^* = -\bar{c}_{n+i} \geq 0,$$

$$i = 1, 2, \dots, m$$

$\Rightarrow y^*$ is also dual feasible solution

Complementary Slackness Theorem

Theorem (Complementary Slackness)

A feasible solution $\mathbf{x}^*$ for (P)

A feasible solution $\mathbf{y}^*$ for (D)

Necessary and sufficient conditions for optimality of both:

$$\left(c_j - \sum_{i=1}^m y_i^* a_{ij} \right) x_j^* = 0, \quad j = 1, \dots, n$$

If $x_j^* \neq 0$ then $\sum y_i^* a_{ij} = c_j$ (no surplus)

If $\sum y_i^* a_{ij} > c_j$ then $x_j^* = 0$

Proof:

$$z^* = \mathbf{c}^T \mathbf{x}^* \leq \mathbf{y}^* \mathbf{A} \mathbf{x}^* \leq \mathbf{b}^T \mathbf{y}^* = w^*$$

Hence from strong duality theorem:

$$\mathbf{c} \mathbf{x}^* - \mathbf{y}^* \mathbf{A} \mathbf{x}^* = 0$$

In scalars

$$\sum_{j=1}^n \underbrace{\left(c_j - \sum_{i=1}^m y_i^* a_{ij} \right)}_{\leq 0} \underbrace{x_j^*}_{\geq 0} = 0$$

Hence each term must be $= 0$

Proof in scalar form:

$$c_j x_j^* \leq \left(\sum_{i=1}^m a_{ij} y_i^* \right) x_j^* \quad j = 1, 2, \dots, n \quad \text{from feasibility in D}$$

$$\left(\sum_{j=1}^n a_{ij} x_j^* \right) y_i^* \leq b_i y_i^* \quad i = 1, 2, \dots, m \quad \text{from feasibility in P}$$

Summing in j and in i :

$$\sum_{j=1}^n c_j x_j^* \leq \sum_{j=1}^n \left(\sum_{i=1}^m a_{ij} y_i^* \right) x_j^* = \sum_{i=1}^m \left(\sum_{j=1}^n a_{ij} x_j^* \right) y_i^* \leq \sum_{i=1}^m b_i y_i^*$$

For the strong duality theorem the left hand side is equal to the right hand side and hence all inequalities become equalities.

$$\sum_{j=1}^n \underbrace{\left(c_j - \sum_{i=1}^m y_i^* a_{ij} \right)}_{\leq 0} \underbrace{x_j^*}_{\geq 0} = 0$$

Economic Interpretation of Duality Theory

$$\begin{aligned}
 \max \quad & 5x_0 + 6x_1 + 8x_2 \\
 & 6x_0 + 5x_1 + 10x_2 \leq 60 \\
 & 8x_0 + 4x_1 + 4x_2 \leq 40 \\
 & 4x_0 + 5x_1 + 6x_2 \leq 50 \\
 & x_0, x_1, x_2 \geq 0
 \end{aligned}$$

final tableau:

$$\begin{array}{ccccccc|c}
 x_0 & x_1 & x_2 & s_1 & s_2 & s_3 & -z & b \\
 \hline
 & 0 & 1 & & 0 & & & 5/2 \\
 & 1 & 0 & & 0 & & & 7 \\
 & 0 & 0 & & 1 & & & 2 \\
 \hline
 -1/5 & 0 & 0 & -1/5 & 0 & -1 & & -62
 \end{array}$$

- What is the maximum number of products that can be produced? at most m
- What are the values of the variables, the reduced costs, the dual variables, and the shadow (or marginal) prices?
- What is the value of additional manpower capacity (1st constr.)? In +1 out +1/5
- If one slack variable > 0 then overcapacity: $s_2 = 2$ then the second constraint is not tight
- How much more expensive would a non-selected product need to be? look at reduced costs:

$$c_j + \pi a_j > 0$$

Economic Interpretation of Duality Theory

Game: Suppose two economic operators:

- P owns the factory and produces goods
- D is in the market buying and selling raw material and resources
- D asks P to close and sell him/her all resources
- P considers if the offer is convenient
- D wants to spend the least possible
- y are prices that D offers for the resources
- $\sum y_i b_i$ is the amount D has to pay to have all resources of P
- $\sum y_i a_{ij} \geq c_j$ total value to make j is greater or eq. then the price per unit of product j
- P either sells all resources $\sum y_i a_{ij}$ or produces product j (c_j)
- without $\geq$ there would not be negotiation because P would be better off producing and selling
- ▶ at optimality the situation is indifferent (strong th.)
- ▶ resource 2 that was not totally utilized in the primal has been given value 0 in the dual.
(complementary slackness th.) Plausible, since we do not use all the resource, likely to place not so much value on it.
- ▶ for product 0 $\sum y_i a_{ij} > c_j$ hence not profitable producing it. (complementary slackness th.)

- Derivation:
 - Economic Interpretation
 - Bounding Approach
 - Multiplier Approach
 - Recipe
 - Lagrangian Multipliers Approach (next time)
- Theory:
 - Symmetry
 - Weak Duality Theorem
 - Strong Duality Theorem
 - Complementary Slackness Theorem