

Page 2: Full EUA Prescribing Information

Page 28: Fact Sheets for Vaccination Providers

Page 38: Fact Sheets for Recipients and Caregivers

From: (b) (6)
Sent: Saturday, December 05, 2020 6:03 PM
To: 'Harkins Tull, Elisa' <Elisa.HarkinsTull@pfizer.com>
Cc: Devlin, Carmel M <Carmel.Devlin@pfizer.com>; (b) (6) 'Boyce, Donna' <Donna.Boyce@pfizer.com>
Subject: EUA 27034 - Pfizer-BioNTech COVID-19 Vaccine - CBER comments on Full EUA PI and Fact Sheets

Dear Ms. Harkins,

Attached please find Word versions of the Full EUA Prescribing Information included in amendment 3 (dated November 25, 2020) and the EUA Fact Sheets included in your initial EUA 27034 submission (dated November 20, 2020), with CBER comments. Please submit the revised versions of these documents by Monday, December 7, 2020.

Please confirm receipt of this email and let me know if you have any questions or need additional information.

Regards,

(b) (6)

Center for Biologics Evaluation and Research
Office of Vaccines Research and Review
U.S. Food and Drug Administration

(b) (6)



THIS MESSAGE IS INTENDED ONLY FOR THE USE OF THE PARTY TO WHOM IT IS ADDRESSED AND MAY CONTAIN INFORMATION THAT IS PRIVILEGED, CONFIDENTIAL, AND PROTECTED FROM DISCLOSURE UNDER LAW. If you are not the addressee, or a person authorized to deliver the document to the addressee, you are hereby notified that any review, disclosure, dissemination, copying, or other action based on the content of this communication is not authorized. If you have received this document in error, please immediately notify the sender by e-mail or phone.

**FULL EMERGENCY USE
AUTHORIZATION (EUA) ~~FULL~~
PRESCRIBING INFORMATION**

PFIZER-BIONTECH COVID-19 VACCINE

**FULL EMERGENCY USE AUTHORIZATION PRESCRIBING
INFORMATION: CONTENTS***

- 1 AUTHORIZED USE
- 2 DOSAGE AND ADMINISTRATION
 - 2.1 Preparation for Administration
 - 2.2 Administration Information
 - 2.3 Vaccination Schedule for Individuals 16 Years of Age and Older
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
 - 5.1 Management of Acute Allergic Reactions
 - ~~5.2 Concurrent Illness at Time of Vaccination~~
 - 5.3 Altered Immunocompetence
 - 5.4 Limitation of Use
- 6 OVERALL SAFETY SUMMARY
 - 6.1 Clinical Trials Experience

- 8 ADVERSE EVENTS AND VACCINE ADMINISTRATION
ERRORS REPORTING REQUIREMENTS AND INSTRUCTIONS
- 9 OTHER REPORTING REQUIREMENTS
- 10 DRUG INTERACTIONS
- 11 USE IN SPECIFIC POPULATIONS
 - 11.1 Pregnancy
 - 11.2 Lactation
 - 11.3 Pediatric Use
 - 11.4 Geriatric Use
- 13 DESCRIPTION
- 14 CLINICAL PHARMACOLOGY
 - 14.1 Mechanism of Action
- 18 CLINICAL TRIAL RESULTS AND SUPPORTING DATA FOR
EUA
 - 18.1 Efficacy in Participants 16 Years of Age and Older
- 19 HOW SUPPLIED/STORAGE AND HANDLING
- 20 PATIENT COUNSELING INFORMATION
- 21 CONTACT INFORMATION

* Sections or subsections omitted from the full emergency use authorization
prescribing information are not listed.

Commented [A1]: Pfizer, please update the Table of
Contents to be consistent with the Full PI.

~~FULL~~ EMERGENCY USE AUTHORIZATION (EUA) ~~FULL~~-PRESCRIBING INFORMATION

1 AUTHORIZED USE

PfizerBioNTech- COVID-19 Vaccine is authorized for use under an Emergency Use Authorization (EUA) for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARSCoV2) in individuals 16 years of age and older.

2 DOSAGE AND ADMINISTRATION

For intramuscular injection only.

2.1 Preparation for Administration

- The Pfizer-BioNTech COVID-19 Vaccine Multiple Dose Vial contains a frozen suspension that does not contain preservative and must be thawed and diluted prior to administration.
- Vials may be thawed in the refrigerator [2°C to 8°C (35°F to 46°F)] or at room temperature [up to 25°C (77°F)] [*see How Supplied/Storage and Handling (19)*].
- Prior to dilution, ~~the~~ thawed suspension may contain white to off-white opaque amorphous particles.
- Dilute the vial contents using 1.8 mL of ONLY 0.9% Sodium Chloride Injection, USP to form the Pfizer-BioNTech COVID-19 Vaccine. ~~The contents of the vial must be diluted with 1.8 mL of sterile 0.9% Sodium Chloride Injection, USP to form the Pfizer BioNTech COVID-19 Vaccine.~~
- Only use 0.9% Sodium Chloride Injection, USP for as the diluent. This ~~The diluent is not packaged with the vaccine and must be sourced separately. Do not use bacteriostatic 0.9% Sodium Chloride Injection or any other diluent.~~
- ~~After dilution, the vial contains five 0.3 mL doses of Pfizer-BioNTech COVID-19 Vaccine. 5 doses of 0.3 mL.~~
- ~~0.9% Sodium Chloride Injection, USP is not packaged with the vaccine and must be sourced separately. Do not use bacteriostatic 0.9% Sodium Chloride Injection or any other diluent.~~
- After dilution, the vaccine will be an off-white suspension. Inspect vials to confirm there are no particulates and no discoloration is observed. If particulates or discoloration are observed, discard the vial.
- Strictly adhere ~~adherence~~ to aseptic techniques must be followed.

Commented [A2]: Pending CBER review of the information that will be submitted in response to CBER's December 2, 2020 CMC IR.

DILUTE BEFORE USE



- Thaw vial(s) of Pfizer-BioNTech COVID-19 Vaccine before use either by:
 - Allowing vial(s) to thaw in the refrigerator [2°C to 8°C (35°F to 46°F)]. A carton of vials may take up to 3 hours to thaw.
 - Allowing vial(s) to sit at room temperature [up to 25°C (77°F)] for 30 minutes
 - Remove ~~Removing~~ a thawed vial of Pfizer-BioNTech COVID-19 Vaccine from the refrigerator and allowing it to come to room temperature.
- ~~Before dilution allow vials to come to room temperature [up to 25°C (77°F)]. (If using a frozen vial of Pfizer-BioNTech COVID-19 Vaccine, thaw for 30 minutes at room temperature.~~
- ~~Continue preparation and administer within 2 hours.~~




~~Once at room temperature, the liquid in the vial is concentrated vaccine that requires dilution with 0.9% Sodium Chloride Injection, USP before administration (see below) to form the Pfizer-BioNTech COVID-19 Vaccine.~~

~~Once at room temperature, the liquid in the vial is concentrated vaccine that requires dilution with 0.9% Sodium Chloride Injection, USP before administration (see below).~~

~~Vials at room temperature must be diluted within 2 hours.~~

~~Vials at room temperature must be diluted within 2 hours.~~



Commented [A3]: Pfizer, please modify time as necessary if vials can be thawed at room temperature in carton.


	• <u>Before dilution i</u>nv<u>ert concentrated vaccine vial gently 10 times. to mix.</u> • <u>Do not shake.</u> • <u>Inspect contents of vial for discoloration. Vial may contain white to off-white opaque amorphous particles.</u>
	• <u>Obtain sterile 0.9% Sodium Chloride Injection, USP. Use only this as the diluent.</u> • <u>Using aseptic technique, withdraw 1.8 mL of diluent into a transfer syringe (21-gauge or narrower needle).</u> • Cleanse the <u>vaccine</u> vial stopper with a single-use antiseptic swab. • Add 1.8 mL of 0.9% Sodium Chloride Injection, USP into the Pfizer BioNTech COVID-19 Vaccine vial using a needle 21-gauge or narrower. Cleanse the vaccine vial stopper with a single use antiseptic swab. • <u>Add 1.8 mL of the 0.9% Sodium Chloride Injection, USP into the vaccine vial.</u>
	• Equalize vial pressure before removing the needle from the vial by withdrawing 1.8 mL air into the empty diluent syringe.

Commented [A4]: Pfizer, please describe how the concentrated vaccine should appear coloration-wise (outside of the particles).

Pending CBER review of the information that will be submitted in response to CBER's December 2, 2020 CMC IR.

Pfizer, please add a comment regarding when the vaccine should be discarded regarding coloration or appearance.

	- Gently invert the vial containing the diluted Pfizer-BioNTech COVID-19 Vaccine 10 times to mix. Do not shake. Diluted The vaccine should will be an off-white suspension. Discard if vaccine is discolored or contains particulate matter.
	- Record the date and time of dilution on the Pfizer-BioNTech COVID-19 Vaccine vial label. - Store between 2°C to 25°C (35°F to 77°F). - Discard any unused vaccine 6 hours after dilution.

PREPARATION OF INDIVIDUAL 0.3 mL DOSES OF DILUTED PFIZER-BIONTECH COVID-19 VACCINE	
	- Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab, and withdraw <u>0.3 mL</u> of the <u>Pfizer BioNTech COVID-19 Vaccine</u>. - Administer immediately.

2.2 Administration Information

Visually inspect each dose in the dosing syringe prior to administration. The vaccine will be an off-white suspension. During the visual inspection,

- verify the final dosing volume of 0.3 mL.
- confirm there are no particulates and that no discoloration is observed.

If the visual inspection fails, do not administer the vaccine.

Administer the Pfizer-BioNTech COVID-19 Vaccine intramuscularly in the deltoid muscle.

2.3 Vaccination Schedule for Individuals 16 Years of Age and Older

The Pfizer-BioNTech COVID-19 Vaccine is administered intramuscularly as a series of two doses (0.3 mL each) ~~at greater than or equal to three weeks~~ 21 days apart.

There are no data available on the interchangeability of the Pfizer-BioNTech COVID-19 Vaccine with other COVID-19 vaccines to complete the vaccination series. Individuals who have received one dose of Pfizer-BioNTech COVID-19 Vaccine should receive a second dose of Pfizer-BioNTech COVID-19 Vaccine to complete the vaccination series.

3 DOSAGE FORMS AND STRENGTHS

Pfizer-BioNTech COVID-19 Vaccine is a suspension for injection. After preparation, a single dose is 0.3 mL.

4 CONTRAINDICATIONS

Do not administer Pfizer-BioNTech COVID-19 Vaccine to individuals with known history of a severe allergic reaction (e.g., anaphylaxis) to any component of the Pfizer-BioNTech COVID-19 Vaccine *[see Description (13)]*.

5 WARNINGS AND PRECAUTIONS

5.1 Management of Acute Allergic Reactions

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of Pfizer-BioNTech COVID-19 Vaccine.

~~5.2 Concurrent Illness at Time of Vaccination~~

~~The administration of the Pfizer BioNTech COVID-19 Vaccine should be postponed in individuals with an acute severe febrile illness.~~

~~5.3.2~~ Altered Immunocompetence

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the Pfizer-BioNTech COVID-19 Vaccine.

~~5.4.3~~ Limitation of ~~Use~~Effectiveness

The Pfizer-BioNTech COVID-19 Vaccine may not protect all vaccine recipients.

6 OVERALL SAFETY SUMMARY

It is MANDATORY for vaccination providers to report to the complete a Vaccine Adverse Event Reporting System (VAERS) form to report and to Pfizer, Inc. all vaccine administration errors, all serious adverse events, cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and hospitalized or fatal cases of COVID-19 that result in hospitalization or death following vaccination with the Pfizer-BioNTech COVID-19 Vaccine. Please see the REQUIREMENTS AND INSTRUCTIONS FOR REPORTING ADVERSE EVENTS AND VACCINE ADMINISTRATION ERRORS ADVERSE EVENTS AND VACCINE ADMINISTRATION ERRORS REPORTING REQUIREMENTS AND INSTRUCTIONS section for details on reporting to VAERS reporting and Pfizer, Inc.

In clinical studies, the adverse reactions in participants 16 years of age and older were pain at the injection site (84.1%), fatigue (62.9%), headache (55.1%), muscle pain (38.3%), chills (31.9%), joint pain (23.6%), and fever (14.2%), and lymphadenopathy (0.3%).

Additional adverse reactions, some of which may be serious, may become apparent with more widespread use of the Pfizer-BioNTech COVID-19 Vaccine.

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of Pfizer-BioNTech COVID-19 Vaccine was evaluated in participants 16 years of age and older in two clinical studies conducted in the United States, Europe, Turkey, South Africa, and South America. Study BNT162-01 (Study 1) was a Phase 1/2, two-part, dose-escalation trial that enrolled 60 participants, 18 through 55 years of age. Study C4591001 (Study 2) is a Phase 1/2/3, multicenter, multinational, randomized, saline placebo-controlled, observer-blind, dose-finding, vaccine candidate-selection (Phase 1/2) and efficacy (Phase 2/3) study that has enrolled approximately 44,000 participants, 12 years of age or older. Of these, approximately 43,448 participants (21,720 Pfizer BioNTech COVID-19 Vaccine, 21,728 placebo) in Phase 2/3 are 16 years of age or older.

At the time of the analysis of Study 2 for the EUA, 37,586+9,067 (18,801+9,534 Pfizer-BioNTech COVID-19 Vaccine and 18,785+9,536 placebo) participants 16 years of age or older have been were followed for a median of evaluated for safety 2 months after the second dose of Pfizer-BioNTech COVID-19 Vaccine.

In Study 2, a total of 21,720 participants 16 years of age or older received at least one dose of Pfizer BioNTech COVID-19 Vaccine and a total of 21,728 participants 16 years of age or older received placebo.

The safety evaluation in Study 2 is ongoing. The safety population includes participants enrolled by October 9, 2020, and includes safety data accrued through November 14, 2020. Participants 18 years and older in the reactogenicity subset are were monitored for solicited local and systemic reaction events, and use of antipyretic medication after each vaccination in an electronic diary maintained by the participant or the participant's parent/legal guardian. Participants are being monitored for unsolicited adverse events, including serious adverse events, throughout the study [from Dose 1 through 1 month (all unsolicited adverse events) or 6 months (serious adverse events) after the last vaccination, depending on the safety parameter].

Commented [A5]: Pfizer, please confirm this is the enrolled AND randomized participants.

Demographic characteristics in Study 2 were generally similar with regard to age, gender, race, and ethnicity among participants who received Pfizer-BioNTech COVID-19 Vaccine and those who received placebo. Overall, among the participants who received Pfizer-BioNTech COVID-19 Vaccine, ~~50.645%~~ were male and ~~49.485%~~ were female, ~~83.124%~~ were White, ~~9.16%~~ were Black or African American, ~~286.04%~~ were Hispanic/Latino, 4.3% were Asian, and ~~0.57%~~ were Native American/Alaskan native.

Local and Systemic Adverse ~~Events-Reactions~~ Solicited in the Study 2

Table 1 and Table 2 present the frequency and severity of solicited local ~~reactions~~ and systemic ~~events~~~~reactions~~, respectively, within 7 days following each dose of Pfizer-BioNTech COVID-19 Vaccine and placebo in the subset of participants 186 to 55 years of age included in the EUA safety population who were monitored for reactogenicity with an electronic diary.

At this time, solicited reactogenicity data in 16 and -17 year-old participants are not available.

~~Local adverse event~~~~reactions~~ at the Pfizer BioNTech COVID-19 Vaccine site and the placebo injection site were assessed:

Table 1: Study 2 – Frequency of Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 186-55 Years of Age – Reactogenicity Subset of the Safety Population*

Local Reaction	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI^c)	N ^a	n ^b (%)	(95% CI^c)	N ^a	n ^b (%)	(95% CI^c)	N ^a	n ^b (%)	(95% CI^c)
Redness ^d												
Any	2291	104 (4.5)	(3.7, 5.5)	2298	26 (1.1)	(0.7, 1.7)	2098	123 (5.9)	(4.9, 7.0)	2103	14 (0.7)	(0.4, 1.1)
Mild	2291	70 (3.1)	(2.4, 3.8)	2298	16 (0.7)	(0.4, 1.1)	2098	73 (3.5)	(2.7, 4.4)	2103	8 (0.4)	(0.2, 0.7)
Moderate	2291	28 (1.2)	(0.8, 1.8)	2298	6 (0.3)	(0.1, 0.6)	2098	40 (1.9)	(1.4, 2.6)	2103	6 (0.3)	(0.1, 0.6)
Severe	2291	6 (0.3)	(0.1, 0.6)	2298	4 (0.2)	(0.0, 0.4)	2098	10 (0.5)	(0.2, 0.9)	2103	0	(0.0, 0.2)
Grade 4	2291	0	(0.0, 0.2)	2298	0	(0.0, 0.2)	2098	0	(0.0, 0.2)	2103	0	(0.0, 0.2)
Swelling ^d												
Any	2291	132 (5.8)	(4.8, 6.8)	2298	11 (0.5)	(0.2, 0.9)	2098	132 (6.3)	(5.3, 7.4)	2103	5 (0.2)	(0.1, 0.6)
Mild	2291	88 (3.8)	(3.1, 4.7)	2298	3 (0.1)	(0.0, 0.4)	2098	80 (3.8)	(3.0, 4.7)	2103	3 (0.1)	(0.0, 0.4)
Moderate	2291	39 (1.7)	(1.2, 2.3)	2298	5 (0.2)	(0.1, 0.5)	2098	45 (2.1)	(1.6, 2.9)	2103	2 (0.1)	(0.0, 0.3)
Severe	2291	5 (0.2)	(0.1, 0.5)	2298	3 (0.1)	(0.0, 0.4)	2098	7 (0.3)	(0.1, 0.7)	2103	0	(0.0, 0.2)
Grade 4	2291	0	(0.0, 0.2)	2298	0	(0.0, 0.2)	2098	0	(0.0, 0.2)	2103	0	(0.0, 0.2)
Pain at the injection site ^e												

Commented [A6]: Pfizer, please redo table with participants 18-55 years of age.

Commented [A7]: Pfizer, please remove the 95% CI column from each safety table, Tables 1-4, and revise footnote accordingly.

Commented [A8]: Pfizer, please delete row for Grade 4 in Tables 1-4 as values are zero.

Table 1: Study 2 – Frequency of Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 18-55 Years of Age – Reactogenicity Subset of the Safety Population*

Local Reaction	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c
Any	2291	1904 (83.1)	(81.5, 84.6)	2298	322 (14.0)	(12.6, 15.5)	2098	1632 (77.8)	(75.9, 79.6)	2103	245 (11.7)	(10.3, 13.1)
Mild	2291	1170 (51.1)	(49.0, 53.1)	2298	308 (13.4)	(12.0, 14.9)	2098	1039 (49.5)	(47.4, 51.7)	2103	225 (10.7)	(9.4, 12.1)
Moderate	2291	710 (31.0)	(29.1, 32.9)	2298	12 (0.5)	(0.3, 0.9)	2098	568 (27.1)	(25.2, 29.0)	2103	20 (1.0)	(0.6, 1.5)
Severe	2291	24 (1.0)	(0.7, 1.6)	2298	2 (0.1)	(0.0, 0.3)	2098	25 (1.2)	(0.8, 1.8)	2103	0	(0.0, 0.2)
Grade 4	2291	0 (0.0, 0.2)		2298	0 (0.0, 0.2)		2098	0 (0.0, 0.2)		2103	0 (0.0, 0.2)	
Any local reaction ^f	2291	1916 (83.6)	(82.1, 85.1)	2298	338 (14.7)	(13.3, 16.2)	2098	1638 (78.1)	(76.2, 79.8)	2103	256 (12.2)	(10.8, 13.6)

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose. Grade 4 reactions were classified by the investigator or medically qualified person.

a. N = number of participants reporting at least 1 yes or no response for the specified reaction after the specified dose.

b. n = Number of participants with the specified characteristic.

c. Exact 2-sided confidence interval (CI) based on the Clopper and Pearson method.

d. Mild: >2.0 to 5.0 cm; Moderate: >5.0 to 10.0 cm; Severe: >10.0 cm; **Grade 4: necrosis (redness and swelling categories) or exfoliative dermatitis (redness category only).**

e. Mild: does not interfere with activity; Moderate: interferes with activity; Severe: prevents daily activity; Grade 4: emergency room visit or hospitalization for severe pain at the injection site.

f. Any local reaction: any redness >2.0 cm, any swelling >2.0 cm, or any pain at the injection site.

* All Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Commented [A6]: Pfizer, please redo table with participants 18-55 years of age.

Commented [A7]: Pfizer, please remove the 95% CI column from each safety table, Tables 1-4, and revise footnote accordingly.

The mean duration of pain at the injection site after Dose 2 was 2.5 days (range 1 to 36 days), for redness 2.6 days (range 1 to 34 days), and for swelling 2.3 days (range 1 to 34 days) for participants 16 to 55 years of age in the Pfizer-BioNTech COVID-19 Vaccine group.

Table 2: Study 2 – Frequency of Solicited Systemic Events/Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 18-55 Years of Age – Safety Population*

Systemic Event	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c
Fever												
≥38.0°C	2291	85 (3.7)	(3.0, 4.6)	2298	20 (0.9)	(0.5, 1.3)	2098	331 (15.8)	(14.2, 17.4)	2103	10 (0.5)	(0.2, 0.9)
≥38.0°C to 38.4°C	2291	64 (2.8)	(2.2, 3.6)	2298	10 (0.4)	(0.2, 0.8)	2098	194 (9.2)	(8.0, 10.6)	2103	5 (0.2)	(0.1, 0.6)
>38.4°C to 38.9°C	2291	15 (0.7)	(0.4, 1.1)	2298	5 (0.2)	(0.1, 0.5)	2098	110 (5.2)	(4.3, 6.3)	2103	3 (0.1)	(0.0, 0.4)
>38.9°C to 40.0°C	2291	6 (0.3)	(0.1, 0.6)	2298	3 (0.1)	(0.0, 0.4)	2098	26 (1.2)	(0.8, 1.8)	2103	2 (0.1)	(0.0, 0.3)

Commented [A10]: Pfizer, please redo table for participants 18 to 55 years of age and older.

Also, please make table 508-compliant.

Table 2: Study 2 – Frequency of Solicited Systemic Events/Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 18-55 Years of Age – Safety Population*

Systemic Event	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c
>40.0°C	2291	0	(0.0, 0.2)	2298	2 (0.1)	(0.0, 0.3)	2098	1 (0.0)	(0.0, 0.3)	2103	0	(0.0, 0.2)
Fatigue ^d												
Any	2291	1085 (47.4)	(45.3, 49.4)	2298	767 (33.4)	(31.4, 35.3)	2098	1247 (59.4)	(57.3, 61.5)	2103	479 (22.8)	(21.0, 24.6)
Mild	2291	597 (26.1)	(24.3, 27.9)	2298	467 (20.3)	(18.7, 22.0)	2098	442 (21.1)	(19.3, 22.9)	2103	248 (11.8)	(10.4, 13.2)
Moderate	2291	455 (19.9)	(18.2, 21.6)	2298	289 (12.6)	(11.2, 14.0)	2098	708 (33.7)	(31.7, 35.8)	2103	217 (10.3)	(9.1, 11.7)
Severe	2291	33 (1.4)	(1.0, 2.0)	2298	11 (0.5)	(0.2, 0.9)	2098	97 (4.6)	(3.8, 5.6)	2103	14 (0.7)	(0.4, 1.1)
Grade 4	2291	0 (0.0, 0.2)		2298	0 (0.0, 0.2)		2098	0 (0.0, 0.2)		2103	0 (0.0, 0.2)	
Headache ^d												
Any	2291	959 (41.9)	(39.8, 43.9)	2298	775 (33.7)	(31.8, 35.7)	2098	1085 (51.7)	(49.6, 53.9)	2103	506 (24.1)	(22.2, 25.9)
Mild	2291	628 (27.4)	(25.6, 29.3)	2298	505 (22.0)	(20.3, 23.7)	2098	538 (25.6)	(23.8, 27.6)	2103	321 (15.3)	(13.8, 16.9)
Moderate	2291	308 (13.4)	(12.1, 14.9)	2298	251 (10.9)	(9.7, 12.3)	2098	480 (22.9)	(21.1, 24.7)	2103	170 (8.1)	(7.0, 9.3)
Severe	2291	23 (1.0)	(0.6, 1.5)	2298	19 (0.8)	(0.5, 1.3)	2098	67 (3.2)	(2.5, 4.0)	2103	15 (0.7)	(0.4, 1.2)
Grade 4	2291	0 (0.0, 0.2)		2298	0 (0.0, 0.2)		2098	0 (0.0, 0.2)		2103	0 (0.0, 0.2)	
Chills ^d												
Any	2291	321 (14.0)	(12.6, 15.5)	2298	146 (6.4)	(5.4, 7.4)	2098	737 (35.1)	(33.1, 37.2)	2103	79 (3.8)	(3.0, 4.7)
Mild	2291	230 (10.0)	(8.8, 11.3)	2298	111 (4.8)	(4.0, 5.8)	2098	359 (17.1)	(15.5, 18.8)	2103	65 (3.1)	(2.4, 3.9)
Moderate	2291	82 (3.6)	(2.9, 4.4)	2298	33 (1.4)	(1.0, 2.0)	2098	333 (15.9)	(14.3, 17.5)	2103	14 (0.7)	(0.4, 1.1)
Severe	2291	9 (0.4)	(0.2, 0.7)	2298	2 (0.1)	(0.0, 0.3)	2098	45 (2.1)	(1.6, 2.9)	2103	0 (0.0, 0.2)	
Grade 4	2291	0 (0.0, 0.2)		2298	0 (0.0, 0.2)		2098	0 (0.0, 0.2)		2103	0 (0.0, 0.2)	
Vomiting ^e												
Any	2291	28 (1.2)	(0.8, 1.8)	2298	28 (1.2)	(0.8, 1.8)	2098	40 (1.9)	(1.4, 2.6)	2103	25 (1.2)	(0.8, 1.7)
Mild	2291	24 (1.0)	(0.7, 1.6)	2298	22 (1.0)	(0.6, 1.4)	2098	28 (1.3)	(0.9, 1.9)	2103	16 (0.8)	(0.4, 1.2)
Moderate	2291	4 (0.2)	(0.0, 0.4)	2298	5 (0.2)	(0.1, 0.5)	2098	8 (0.4)	(0.2, 0.7)	2103	9 (0.4)	(0.2, 0.8)
Severe	2291	0 (0.0, 0.2)		2298	1 (0.0)	(0.0, 0.2)	2098	4 (0.2)	(0.1, 0.5)	2103	0 (0.0, 0.2)	
Grade 4	2291	0 (0.0, 0.2)		2298	0 (0.0, 0.2)		2098	0 (0.0, 0.2)		2103	0 (0.0, 0.2)	
Diarrhea ^f												
Any	2291	255 (11.1)	(9.9, 12.5)	2298	270 (11.7)	(10.5, 13.1)	2098	219 (10.4)	(9.2, 11.8)	2103	177 (8.4)	(7.3, 9.7)
Mild	2291	206 (9.0)	(7.9, 10.2)	2298	217 (9.4)	(8.3, 10.7)	2098	179 (8.5)	(7.4, 9.8)	2103	144 (6.8)	(5.8, 8.0)
Moderate	2291	46 (2.0)	(1.5, 2.7)	2298	52 (2.3)	(1.7, 3.0)	2098	36 (1.7)	(1.2, 2.4)	2103	32 (1.5)	(1.0, 2.1)

Commented [A10]: Pfizer, please redo table for participants 18 to 55 years of age and older.

Also, please make table 508-compliant.

Table 2: Study 2 – Frequency of Solicited Systemic Events/Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 18-55 Years of Age – Safety Population*

Systemic Event	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c
Severe	2291	3 (0.1)	(0.0, 0.4)	2298	1 (0.0)	(0.0, 0.2)	2098	4 (0.2)	(0.1, 0.5)	2103	1 (0.0)	(0.0, 0.3)
Grade 4	2291	0	(0.0, 0.2)	2298	0	(0.0, 0.2)	2098	0	(0.0, 0.2)	2103	0	(0.0, 0.2)
New or worsened muscle pain ^d												
Any	2291	487 (21.3)	(19.6, 23.0)	2298	249 (10.8)	(9.6, 12.2)	2098	783 (37.3)	(35.2, 39.4)	2103	173 (8.2)	(7.1, 9.5)
Mild	2291	256 (11.2)	(9.9, 12.5)	2298	175 (7.6)	(6.6, 8.8)	2098	326 (15.5)	(14.0, 17.2)	2103	111 (5.3)	(4.4, 6.3)
Moderate	2291	218 (9.5)	(8.3, 10.8)	2298	72 (3.1)	(2.5, 3.9)	2098	410 (19.5)	(17.9, 21.3)	2103	59 (2.8)	(2.1, 3.6)
Severe	2291	13 (0.6)	(0.3, 1.0)	2298	2 (0.1)	(0.0, 0.3)	2098	47 (2.2)	(1.7, 3.0)	2103	3 (0.1)	(0.0, 0.4)
Grade 4	2291	0	(0.0, 0.2)	2298	0	(0.0, 0.2)	2098	0	(0.0, 0.2)	2103	0	(0.0, 0.2)
New or worsened joint pain ^d												
Any	2291	251 (11.0)	(9.7, 12.3)	2298	138 (6.0)	(5.1, 7.1)	2098	459 (21.9)	(20.1, 23.7)	2103	109 (5.2)	(4.3, 6.2)
Mild	2291	147 (6.4)	(5.4, 7.5)	2298	95 (4.1)	(3.4, 5.0)	2098	205 (9.8)	(8.5, 11.1)	2103	54 (2.6)	(1.9, 3.3)
Moderate	2291	99 (4.3)	(3.5, 5.2)	2298	43 (1.9)	(1.4, 2.5)	2098	234 (11.2)	(9.8, 12.6)	2103	51 (2.4)	(1.8, 3.2)
Severe	2291	5 (0.2)	(0.1, 0.5)	2298	0	(0.0, 0.2)	2098	20 (1.0)	(0.6, 1.5)	2103	4 (0.2)	(0.1, 0.5)
Grade 4	2291	0	(0.0, 0.2)	2298	0	(0.0, 0.2)	2098	0	(0.0, 0.2)	2103	0	(0.0, 0.2)
Any systemic event ^e	2291	1538 (67.1)	(65.2, 69.1)	2298	1243 (54.1)	(52.0, 56.1)	2098	1557 (74.2)	(72.3, 76.1)	2103	803 (38.2)	(36.1, 40.3)
Use of antipyretic or pain medication ^h	2291	638 (27.8)	(26.0, 29.7)	2298	332 (14.4)	(13.0, 16.0)	2098	945 (45.0)	(42.9, 47.2)	2103	266 (12.6)	(11.3, 14.1)

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose. Grade 4 events were classified by the investigator or medically qualified person.

a. N = number of participants reporting at least 1 yes or no response for the specified event after the specified dose.

b. n = Number of participants with the specified characteristic.

c. Exact 2-sided confidence interval (CI) based on the Clopper and Pearson method.

d. Mild: does not interfere with activity; Moderate: some interference with activity; Severe: prevents daily activity; ~~Grade 4: emergency room visit or hospitalization for severe fatigue, severe headache, severe muscle pain, or severe joint pain.~~

e. Mild: 1 to 2 times in 24 hours; Moderate: >2 times in 24 hours; Severe: requires intravenous hydration; ~~Grade 4: emergency room visit or hospitalization for severe vomiting.~~

f. Mild: 2 to 3 loose stools in 24 hours; Moderate: 4 to 5 loose stools in 24 hours; Severe: 6 or more loose stools in 24 hours; ~~Grade 4: emergency room visit or hospitalization for severe diarrhea.~~

g. Any systemic event: any fever ≥38.0°C, any fatigue, any vomiting, any chills, any diarrhea, any headache, any new or worsened muscle pain, or any new or worsened joint pain.

h. Severity was not collected for use of antipyretic or pain medication.

*All randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Commented [A10]: Pfizer, please redo table for participants 18 to 55 years of age and older.

Also, please make table 508-compliant.

Table 3 and Table 4 present the frequency and severity of reported solicited local ~~reactions~~ and systemic ~~events~~~~reactions~~, respectively, within 7 days of each dose of Pfizer-BioNTech COVID-19 Vaccine and placebo for participants 56 years of age and older.

Table 3: Study 2 – Frequency of Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 56 Years of Age and Older – Safety Population*

Local Reaction	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c
Redness ^d												
Any (> 2 cm)	1802	85 (4.7)	(3.8, 5.8)	1792	19 (1.1)	(0.6, 1.7)	1660	120 (7.2)	(6.0, 8.6)	1646	12 (0.7)	(0.4, 1.3)
Mild	1802	55 (3.1)	(2.3, 4.0)	1792	12 (0.7)	(0.3, 1.2)	1660	59 (3.6)	(2.7, 4.6)	1646	8 (0.5)	(0.2, 1.0)
Moderate	1802	27 (1.5)	(1.0, 2.2)	1792	5 (0.3)	(0.1, 0.6)	1660	53 (3.2)	(2.4, 4.2)	1646	3 (0.2)	(0.0, 0.5)
Severe	1802	3 (0.2)	(0.0, 0.5)	1792	2 (0.1)	(0.0, 0.4)	1660	8 (0.5)	(0.2, 0.9)	1646	1 (0.1)	(0.0, 0.3)
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
Swelling ^d												
Any (> 2 cm)	1802	118 (6.5)	(5.4, 7.8)	1792	21 (1.2)	(0.7, 1.8)	1660	124 (7.5)	(6.3, 8.8)	1646	11 (0.7)	(0.3, 1.2)
Mild	1802	71 (3.9)	(3.1, 4.9)	1792	10 (0.6)	(0.3, 1.0)	1660	68 (4.1)	(3.2, 5.2)	1646	5 (0.3)	(0.1, 0.7)
Moderate	1802	45 (2.5)	(1.8, 3.3)	1792	11 (0.6)	(0.3, 1.1)	1660	53 (3.2)	(2.4, 4.2)	1646	5 (0.3)	(0.1, 0.7)
Severe	1802	2 (0.1)	(0.0, 0.4)	1792	0	(0.0, 0.2)	1660	3 (0.2)	(0.0, 0.5)	1646	1 (0.1)	(0.0, 0.3)
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
Pain at the injection site ^e												
Any	1802	1282 (71.1)	(69.0, 73.2)	1792	166 (9.3)	(8.0, 10.7)	1660	1098 (66.1)	(63.8, 68.4)	1646	127 (7.7)	(6.5, 9.1)
Mild	1802	1008 (55.9)	(53.6, 58.2)	1792	160 (8.9)	(7.6, 10.3)	1660	792 (47.7)	(45.3, 50.1)	1646	125 (7.6)	(6.4, 9.0)
Moderate	1802	270 (15.0)	(13.4, 16.7)	1792	6 (0.3)	(0.1, 0.7)	1660	298 (18.0)	(16.1, 19.9)	1646	2 (0.1)	(0.0, 0.4)
Severe	1802	4 (0.2)	(0.1, 0.6)	1792	0	(0.0, 0.2)	1660	8 (0.5)	(0.2, 0.9)	1646	0	(0.0, 0.2)
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
Any local reaction ^f	1802	1300 (72.1)	(70.0, 74.2)	1792	187 (10.4)	(9.1, 11.9)	1660	1110 (66.9)	(64.5, 69.1)	1646	140 (8.5)	(7.2, 10.0)

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose. Grade 4 reactions were classified by the investigator or medically qualified person.

a. N = number of participants reporting at least 1 yes or no response for the specified reaction after the specified dose.

b. n = Number of participants with the specified characteristic.

Commented [A11]: Pfizer, please make all tables 508-compliant.

Table 3: Study 2 – Frequency of Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 56 Years of Age and Older – Safety Population*

Local Reaction	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c

c. Exact 2-sided confidence interval (CI) based on the Clopper and Pearson method.

d. Mild: >2.0 to ≤5.0 cm; Moderate: >5.0 to ≤10.0 cm; Severe: >10.0 cm; ~~Grade 4: necrosis (redness and swelling categories) or exfoliative dermatitis (redness category only).~~

e. Mild: does not interfere with activity; Moderate: interferes with activity; Severe: prevents daily activity; Grade 4: emergency room visit or hospitalization for severe pain at the injection site.

f. Any local reaction: any redness >2.0 cm, any swelling >2.0 cm, or any pain at the injection site.

* All randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Commented [A11]: Pfizer, please make all tables 508-compliant.

Table 4: Study 2 – Frequency of Solicited Systemic ~~Events~~ Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 56 Years of Age and Older – Reactogenicity Subset of the Safety Population*

Systemic Event	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c
Fever												
≥38.0°C	1802	26 (1.4)	(0.9, 2.1)	1792	7 (0.4)	(0.2, 0.8)	1660	181 (10.9)	(9.4, 12.5)	1646	4 (0.2)	(0.1, 0.6)
≥38.0°C to 38.4°C	1802	23 (1.3)	(0.8, 1.9)	1792	2 (0.1)	(0.0, 0.4)	1660	131 (7.9)	(6.6, 9.3)	1646	2 (0.1)	(0.0, 0.4)
>38.4°C to 38.9°C	1802	1 (0.1)	(0.0, 0.3)	1792	3 (0.2)	(0.0, 0.5)	1660	45 (2.7)	(2.0, 3.6)	1646	1 (0.1)	(0.0, 0.3)
>38.9°C to 40.0°C	1802	1 (0.1)	(0.0, 0.3)	1792	2 (0.1)	(0.0, 0.4)	1660	5 (0.3)	(0.1, 0.7)	1646	1 (0.1)	(0.0, 0.3)
>40.0°C	1802	1 (0.1)	(0.0, 0.3)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
Fatigue ^d												
Any	1802	615 (34.1)	(31.9, 36.4)	1792	405 (22.6)	(20.7, 24.6)	1660	839 (50.5)	(48.1, 53.0)	1646	277 (16.8)	(15.1, 18.7)
Mild	1802	373 (20.7)	(18.8, 22.6)	1792	252 (14.1)	(12.5, 15.8)	1660	351 (21.1)	(19.2, 23.2)	1646	161 (9.8)	(8.4, 11.3)
Moderate	1802	240 (13.3)	(11.8, 15.0)	1792	150 (8.4)	(7.1, 9.7)	1660	442 (26.6)	(24.5, 28.8)	1646	114 (6.9)	(5.7, 8.3)
Severe	1802	2 (0.1)	(0.0, 0.4)	1792	3 (0.2)	(0.0, 0.5)	1660	46 (2.8)	(2.0, 3.7)	1646	2 (0.1)	(0.0, 0.4)
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
Headache ^d												
Any	1802	454 (25.2)	(23.2, 27.3)	1792	325 (18.1)	(16.4, 20.0)	1660	647 (39.0)	(36.6, 41.4)	1646	229 (13.9)	(12.3, 15.7)

Table 4: Study 2 – Frequency of Solicited Systemic ~~Events~~Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 56 Years of Age and Older – Reactogenicity Subset of the Safety Population*

Systemic Event	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c
Mild	1802	348 (19.3)	(17.5, 21.2)	1792	242 (13.5)	(12.0, 15.2)	1660	422 (25.4)	(23.3, 27.6)	1646	165 (10.0)	(8.6, 11.6)
Moderate	1802	104 (5.8)	(4.7, 6.9)	1792	80 (4.5)	(3.6, 5.5)	1660	216 (13.0)	(11.4, 14.7)	1646	60 (3.6)	(2.8, 4.7)
Severe	1802	2 (0.1)	(0.0, 0.4)	1792	3 (0.2)	(0.0, 0.5)	1660	9 (0.5)	(0.2, 1.0)	1646	4 (0.2)	(0.1, 0.6)
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
Chills ^d												
Any	1802	113 (6.3)	(5.2, 7.5)	1792	57 (3.2)	(2.4, 4.1)	1660	377 (22.7)	(20.7, 24.8)	1646	46 (2.8)	(2.1, 3.7)
Mild	1802	87 (4.8)	(3.9, 5.9)	1792	40 (2.2)	(1.6, 3.0)	1660	199 (12.0)	(10.5, 13.6)	1646	35 (2.1)	(1.5, 2.9)
Moderate	1802	26 (1.4)	(0.9, 2.1)	1792	16 (0.9)	(0.5, 1.4)	1660	161 (9.7)	(8.3, 11.2)	1646	11 (0.7)	(0.3, 1.2)
Severe	1802	0 (0.0, 0.2)		1792	1 (0.1)	(0.0, 0.3)	1660	17 (1.0)	(0.6, 1.6)	1646	0 (0.0, 0.2)	
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
Vomiting ^e												
Any	1802	9 (0.5)	(0.2, 0.9)	1792	9 (0.5)	(0.2, 1.0)	1660	11 (0.7)	(0.3, 1.2)	1646	5 (0.3)	(0.1, 0.7)
Mild	1802	8 (0.4)	(0.2, 0.9)	1792	9 (0.5)	(0.2, 1.0)	1660	9 (0.5)	(0.2, 1.0)	1646	5 (0.3)	(0.1, 0.7)
Moderate	1802	1 (0.1)	(0.0, 0.3)	1792	0 (0.0, 0.2)		1660	1 (0.1)	(0.0, 0.3)	1646	0 (0.0, 0.2)	
Severe	1802	0 (0.0, 0.2)		1792	0 (0.0, 0.2)		1660	1 (0.1)	(0.0, 0.3)	1646	0 (0.0, 0.2)	
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
Diarrhea ^f												
Any	1802	147 (8.2)	(6.9, 9.5)	1792	118 (6.6)	(5.5, 7.8)	1660	137 (8.3)	(7.0, 9.7)	1646	99 (6.0)	(4.9, 7.3)
Mild	1802	118 (6.5)	(5.4, 7.8)	1792	100 (5.6)	(4.6, 6.7)	1660	114 (6.9)	(5.7, 8.2)	1646	73 (4.4)	(3.5, 5.5)
Moderate	1802	26 (1.4)	(0.9, 2.1)	1792	17 (0.9)	(0.6, 1.5)	1660	21 (1.3)	(0.8, 1.9)	1646	22 (1.3)	(0.8, 2.0)
Severe	1802	3 (0.2)	(0.0, 0.5)	1792	1 (0.1)	(0.0, 0.3)	1660	2 (0.1)	(0.0, 0.4)	1646	4 (0.2)	(0.1, 0.6)
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)

Table 4: Study 2 – Frequency of Solicited Systemic ~~Events~~Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 56 Years of Age and Older – ~~Reactogenicity~~ Subset of the Safety Population*

Systemic Event	Dose 1						Dose 2					
	Pfizer-BioNTech COVID-19 Vaccine			Placebo			Pfizer-BioNTech COVID-19 Vaccine			Placebo		
	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c
New or worsened muscle pain ^d												
Any	1802	251 (13.9)	(12.4, 15.6)	1792	149 (8.3)	(7.1, 9.7)	1660	477 (28.7)	(26.6, 31.0)	1646	87 (5.3)	(4.3, 6.5)
Mild	1802	168 (9.3)	(8.0, 10.8)	1792	100 (5.6)	(4.6, 6.7)	1660	202 (12.2)	(10.6, 13.8)	1646	57 (3.5)	(2.6, 4.5)
Moderate	1802	82 (4.6)	(3.6, 5.6)	1792	46 (2.6)	(1.9, 3.4)	1660	259 (15.6)	(13.9, 17.4)	1646	29 (1.8)	(1.2, 2.5)
Severe	1802	1 (0.1)	(0.0, 0.3)	1792	3 (0.2)	(0.0, 0.5)	1660	16 (1.0)	(0.6, 1.6)	1646	1 (0.1)	(0.0, 0.3)
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
New or worsened joint pain ^d												
Any	1802	155 (8.6)	(7.3, 10.0)	1792	109 (6.1)	(5.0, 7.3)	1660	313 (18.9)	(17.0, 20.8)	1646	61 (3.7)	(2.8, 4.7)
Mild	1802	101 (5.6)	(4.6, 6.8)	1792	68 (3.8)	(3.0, 4.8)	1660	161 (9.7)	(8.3, 11.2)	1646	35 (2.1)	(1.5, 2.9)
Moderate	1802	52 (2.9)	(2.2, 3.8)	1792	40 (2.2)	(1.6, 3.0)	1660	145 (8.7)	(7.4, 10.2)	1646	25 (1.5)	(1.0, 2.2)
Severe	1802	2 (0.1)	(0.0, 0.4)	1792	1 (0.1)	(0.0, 0.3)	1660	7 (0.4)	(0.2, 0.9)	1646	1 (0.1)	(0.0, 0.3)
Grade 4	1802	0	(0.0, 0.2)	1792	0	(0.0, 0.2)	1660	0	(0.0, 0.2)	1646	0	(0.0, 0.2)
Any systemic event ^e	1802	883 (49.0)	(46.7, 51.3)	1792	679 (37.9)	(35.6, 40.2)	1660	1070 (64.5)	(62.1, 66.8)	1646	464 (28.2)	(26.0, 30.4)
Use of antipyretic or pain medication ^h	1802	358 (19.9)	(18.0, 21.8)	1792	213 (11.9)	(10.4, 13.5)	1660	625 (37.7)	(35.3, 40.0)	1646	161 (9.8)	(8.4, 11.3)

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose. Grade 4 events were classified by the investigator or medically qualified person.

a. N = number of participants reporting at least 1 yes or no response for the specified event after the specified dose.

b. n = Number of participants with the specified characteristic.

c. Exact 2-sided CI based on the Clopper and Pearson method.

d. Mild: does not interfere with activity; Moderate: some interference with activity; Severe: prevents daily activity; ~~Grade 4: emergency room visit or hospitalization for severe fatigue, severe headache, severe muscle pain, or severe joint pain.~~

e. Mild: 1 to 2 times in 24 hours; Moderate: >2 times in 24 hours; Severe: requires intravenous hydration; ~~Grade 4: emergency room visit or hospitalization for severe vomiting.~~

f. Mild: 2 to 3 loose stools in 24 hours; Moderate: 4 to 5 loose stools in 24 hours; Severe: 6 or more loose stools in 24 hours; ~~Grade 4: emergency room visit or hospitalization for severe diarrhea.~~

g. Any systemic event: any fever ≥38.0°C, any fatigue, any vomiting, any chills, any diarrhea, any headache, any new or worsened muscle pain, or any new or worsened joint pain.

h. ~~Severity was not collected for use of antipyretic or pain medication.~~

* ~~All~~ Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Unsolicited Adverse Events

Serious Adverse Events

In Study 2, ~~among in~~ participants 16 to 55 years of age ~~who had received at least 1 dose of vaccine or placebo~~ (Pfizer-BioNTech COVID-19 Vaccine = 10,841, placebo = 10,851), ~~received at least 1 dose of vaccine or placebo~~ serious adverse events from Dose 1 through 30 days after Dose 2 were reported by 0.4% of ~~Pfizer-BioNTech COVID-19 Vaccine vaccine-recipients participants~~ and by 0.3% of ~~placebo recipients participants who received at least 1 dose of Pfizer-BioNTech COVID-19 Vaccine or placebo, respectively~~. In Study 2, in participants 56 years of age and older (Pfizer-BioNTech COVID-19 Vaccine = 7960, placebo = 7934), serious adverse events were reported by 0.8% of ~~vaccine recipients participants~~ and by 0.6% of ~~placebo recipients participants~~ who received at least 1 dose of Pfizer-BioNTech COVID-19 Vaccine or placebo, respectively. There were no notable patterns or numerical imbalances between treatment groups for specific categories of serious adverse events (including neurologic, neuro-inflammatory, and thrombotic events) that would suggest a causal relationship to Pfizer-BioNTech COVID-19 Vaccine.

Non-Serious Adverse Events

Overall in Study 2, in which 10,841 participants 16 to 55 years of age received Pfizer-BioNTech COVID-19 Vaccine and 10,851 participants received placebo, non-serious adverse events from Dose 1 through 30 days after Dose 2 were reported in 29.3% of participants who received Pfizer-BioNTech COVID-19 Vaccine and 13.2% of participants in the placebo group, for participants who received at least 1 dose. Overall in Study 2, in which 7960 participants 56 years of age and older received Pfizer-BioNTech COVID-19 Vaccine, non-serious adverse events within 30 days were reported in 23.8% of participants who received Pfizer-BioNTech COVID-19 Vaccine and 11.7% of participants in the placebo group, for participants who received at least 1 dose. - The higher frequency of reported unsolicited non-serious adverse events among Pfizer BioNTech COVID-19 Vaccine recipients compared to placebo recipients was primarily attributed to local and systemic adverse events reported during the first 7 days following vaccination that are consistent with adverse reactions solicited among participants in the reactogenicity subset and presented in Tables 3 and 4. From Dose 1 through 30 days after Dose 2, reports of lymphadenopathy were imbalanced with notably more cases in the vaccine group (64) vs. the placebo group (6), which is plausibly related to vaccination. From dose 1 through 30 days after dose 2, there were three reports of Bell's palsy in the Pfizer BioNTech COVID-19 Vaccine group and none in the placebo group. This observed frequency of reported Bell's palsy is consistent with the expected background rate in the general population. attributed to adverse events consistent with reactogenicity events. There were no other notable patterns or numerical imbalances between treatment groups for specific categories of non-serious adverse events (including- other neurologic, or neuro-inflammatory, and thrombotic events) that would suggest a causal relationship to Pfizer BioNTech COVID-19 Vaccine.

8 **REQUIREMENTS AND INSTRUCTIONS FOR REPORTING ADVERSE EVENTS AND VACCINE ADMINISTRATION ERRORS** ~~REPORTING REQUIREMENTS AND INSTRUCTIONS~~

See Overall Safety Summary (Section 6) for additional information.

The vaccination provider enrolled in the federal COVID-19 Vaccination Program is responsible for MANDATORY reporting of the listed events following Pfizer-BioNTech COVID-19 Vaccine to the Vaccine Adverse Event Reporting System (VAERS) and to Pfizer Inc.:

- Vaccine administration errors whether or not associated with an adverse event
- Serious adverse events* (irrespective of attribution to vaccination)
- Cases of Multisystem Inflammatory Syndrome (MIS) in children and adults
- Cases of COVID-19 that result in hospitalization or death

*Serious adverse events are defined as:

- Death
- A life-threatening adverse event
- Inpatient hospitalization or prolongation of existing hospitalization
- A persistent or significant incapacity or substantial disruption of the ability to conduct normal life functions
- A congenital anomaly/birth defect
- An important medical event that based on appropriate medical judgement may jeopardize the individual and may require medical or surgical intervention to prevent one of the outcomes listed above

Instructions for reporting to VAERS

The vaccination provider enrolled in the federal COVID-19 Vaccination Program should complete and submit a VAERS form to FDA using one of the following methods:

- Complete and submit the report online: <https://vaers.hhs.gov/reportevent.html>, or
- If you are unable to submit this form electronically, you may fax it to VAERS at 1-877-721-0366. If you need additional help submitting a report you may call the VAERS toll-free information line at 1-800-822-7967 or send an email to info@vaers.org.

IMPORTANT: When reporting adverse events or vaccine administration errors to VAERS, please complete the entire form with detailed information. It is important that the information reported to FDA be as detailed and complete as possible. Information to include:

- Patient demographics (e.g., patient name, date of birth)
- Pertinent medical history
- Pertinent details regarding admission and course of illness
- Concomitant medications
- Timing of adverse event(s) in relationship to administration of the Pfizer-BioNTech COVID-19 Vaccine
- Pertinent laboratory and virology information
- Outcome of the event and any additional follow-up information if it is available at the time of the VAERS report. Subsequent reporting of follow-up information should be completed if additional details become available.

The following steps are highlighted to provide the necessary information for safety tracking:

1. In Box 17, provide information on Pfizer-BioNTech COVID-19 Vaccine and any other vaccines administered on the same day; and in Box 22, provide information on any other vaccines received within one month prior.

2. In Box 18, description of the event:
 - a. Write “Pfizer-BioNTech COVID-19 Vaccine EUA” as the first line.
 - b. Provide a detailed report of vaccine administration error and/or adverse event. It is important to provide detailed information regarding the patient and adverse event/medication error for ongoing safety evaluation of this unapproved vaccine. Please see information to include listed above.
3. Contact information:
 - a. In Box 13, provide the name and contact information of the prescribing healthcare provider or institutional designee who is responsible for the report.
 - b. In Box 14, provide the name and contact information of the best doctor/healthcare professional to contact about the adverse event.
 - c. In Box 15, provide the address of the facility where vaccine was given (NOT the healthcare provider’s office address).

9—OTHER REPORTING REQUIREMENTS

Instructions for reporting to Pfizer, Inc.

~~In addition, the vaccination provider must report all adverse events and the events described in Adverse Events and Vaccine Administration Errors Reporting Requirements and Instructions (Section 8) Report to Pfizer Inc. (contact information below) or by providing a copy of the VAERS form to Pfizer Inc.~~

Website	Fax number	Telephone number
www.pfizersafetyreporting.com	1-866-635-8337	1-800-438-1985

Other Reporting Instructions

~~The vaccination providers may report to VAERS and Pfizer Inc. other adverse events that are not required to be reported using the contact information above.~~

10 DRUG INTERACTIONS

~~There isare no information on data to assess the concomitant -administration of the Pfizer-BioNTech COVID-19 Vaccine with other vaccines.~~

11 USE IN SPECIFIC POPULATIONS

11.1 Pregnancy

Risk Summary

All pregnancies have a risk of birth defect, loss, or other adverse outcomes. In the US general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. Available data on Pfizer-BioNTech COVID-19 Vaccine administered to pregnant women are insufficient to inform vaccine-associated risks in pregnancy. ~~Therefore, the Pfizer-BioNTech COVID-19 Vaccine is not recommended for use during pregnancy.~~

Data

Animal Data

~~Reproduction studies in animals have not been completed.~~

11.2 Lactation

Risk Summary

Data are not available to assess the effects of Pfizer-BioNTech COVID-19 Vaccine on the breastfed infant or on milk production/excretion. ~~Therefore, the Pfizer BioNTech COVID-19 Vaccine is not recommended for use during breastfeeding.~~

~~The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Pfizer BioNTech COVID-19 Vaccine and any potential adverse effects on the breastfed child from Pfizer BioNTech COVID-19 Vaccine or from the underlying maternal condition. For preventive vaccines, the underlying maternal condition is susceptibility to disease prevented by the vaccine.~~

11.3 Pediatric Use

Emergency Use Authorization of Pfizer BioNTech COVID-19 Vaccine does not include use in individuals younger than 16 years of age~~The safety and effectiveness of Pfizer BioNTech COVID-19 Vaccine in individuals younger than 16 years of age have not been established.~~

11.4 Geriatric Use

Clinical studies of Pfizer-BioNTech COVID-19 Vaccine include participants 65 years of age and older and their data contributes to the overall assessment of safety and efficacy [see Overall Safety Summary (6.0) and Clinical Trial Results and Supporting Data for EUA (18.1)]. Of the total number of Pfizer-BioNTech COVID-19 Vaccine recipients in Study 2 (N=20,033), 21.417.1% (n=4,2943434) were 65 years of age and older and through 74 years of age and 4.3% (n=860) were 75 years of age and older. ~~[see Clinical Trial Results and Supporting Data for EUA (18.1)]~~

13 DESCRIPTION

The Pfizer-BioNTech COVID-19 Vaccine Suspension for Intramuscular Injection, is a white to off-white, ~~sterile, preservative free,~~ frozen suspension for intramuscular injection that does not contain preservative. The Pfizer-BioNTech COVID-19 Vaccine contains a nucleoside-modified messenger RNA (modRNA) encoding the viral spike glycoprotein (S) of SARS-CoV-2. The Pfizer-BioNTech COVID-19 Vaccine also includes the following ingredients: lipids ((4-hydroxybutyl)azanediyl)bis(hexane-6,1-diyl)bis(2-hexyldecanoate), 2-[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide, 1,2-distearoyl-sn-glycero-3-phosphocholine, and cholesterol), potassium chloride, monobasic potassium phosphate, sodium chloride, dibasic sodium phosphate dihydrate, and sucrose ~~[as stabilizer]~~.

~~The vial stoppers are not made with natural rubber latex. The Pfizer BioNTech COVID-19 Vaccine is packaged in a clear glass 2 mL vial with a rubber stopper (not made with natural rubber latex), aluminum overseal, and flip-off cap.~~

Commented [A12]: Pfizer, please revise this paragraph to include the amounts of each component/ingredient in 0.3 mL dose of the vaccine.

14 CLINICAL PHARMACOLOGY

14.1 Mechanism of Action

The modRNA in the Pfizer-BioNTech COVID-19 Vaccine is formulated in lipid ~~nanoparticles~~, which enable delivery of the RNA into host cells to allow expression of the SARS-CoV-2 S antigen. The vaccine elicits an immune response to the both neutralizing antibody and cellular immune responses to the S antigen, which ~~may contribute to protection~~ against COVID-19.

18 CLINICAL TRIAL RESULTS AND SUPPORTING DATA FOR EUA

18.1 Efficacy in Participants 16 Years of Age and Older

Study 2 is a multicenter, multinational, Phase 1/2/3, randomized, placebo-controlled, observer-blind, dose-finding, vaccine candidate-selection, and efficacy study in participants 12 years of age and older. Randomization was stratified by age: 12 through 15 years of age, 16 through 55 years of age, or 56 years of age and older, with a minimum of 40% of participants in the ≥56-year stratum. The study excluded participants who were immunocompromised and those who had previous clinical or microbiological diagnosis of COVID-19. Participants with preexisting stable disease, defined as disease not requiring significant change in therapy or hospitalization for worsening disease during the 6 weeks before enrollment, were included as were participants with known stable infection with human immunodeficiency virus (HIV), hepatitis C virus (HCV), or hepatitis B virus (HBV). ~~At the time of the analysis of Study 2 for the EUA, information presented is based on participants 16 years and older.~~

In the Phase 2/3 portion approximately 44,000 participants 12 years of age and older were randomized equally and received 2 doses of Pfizer-BioNTech COVID-19 Vaccine or placebo separated by 21 days. Participants are planned to be followed for up to 24 months, for assessments of safety and efficacy against COVID-19.

The population for the analysis of the primary efficacy endpoint included, 36,621 participants 12 years of age and older (18,242 in the Pfizer-BioNTech COVID-19 Vaccine group and 18,379 in the placebo group) who did not have evidence of prior infection with SARS-CoV-2 through 7 days after the second dose. Table 5 presents the specific demographic characteristics in the studied population.

Commented [A13]: Pfizer, please fix this to only include participants 16 years of age and older.

Please also adjust the number of participants and the table below as necessary.

For the table, please remove all data for 12-15 year olds.

Please remove the Total column.

Table 5: Demographics (population for the primary efficacy endpoint)*

	Pfizer-BioNTech COVID-19 Vaccine (N=18,242) n (%)	Placebo (N=18,379) n (%)	Total (N=36,621) n (%)
Sex			
Male	9318 (51.1)	9225 (50.2)	18,543 (50.6)
Female	8924 (48.9)	9154 (49.8)	18,078 (49.4)
Age (years)			
Mean (SD)	50.6 (15.70)	50.4 (15.81)	50.5 (15.76)
Median	52.0	52.0	52.0
Min, max	(12, 89)	(12, 91)	(12, 91)
Age group			
≥12 through 15 years	46 (0.3)	42 (0.2)	88 (0.2)
≥16 through 64 years	14,216 (77.9)	14,299 (77.8)	28,515 (77.9)
≥65 through 74 years	3176 (17.4)	3226 (17.6)	6402 (17.5)
≥75 years	804 (4.4)	812 (4.4)	1616 (4.4)

Table 5: Demographics (population for the primary efficacy endpoint)*

	Pfizer-BioNTech COVID-19 Vaccine (N=18,242) n (%)	Placebo (N=18,379) n (%)	Total (N=36,621) n (%)
Race			
White	15,110 (82.8)	15,301 (83.3)	30,411 (83.0)
Black or African American	1617 (8.9)	1617 (8.8)	3234 (8.8)
American Indian or Alaska native	118 (0.6)	106 (0.6)	224 (0.6)
Asian	815 (4.5)	810 (4.4)	1625 (4.4)
Native Hawaiian or other Pacific Islander	48 (0.3)	29 (0.2)	77 (0.2)
Other†	534 (2.9)	516 (2.8)	1050 (2.9)
Ethnicity			
Hispanic or Latino	4886 (26.8)	4857 (26.4)	9743 (26.6)
Not Hispanic or Latino	13,253 (72.7)	13,412 (73.0)	26,665 (72.8)
Not reported	103 (0.6)	110 (0.6)	213 (0.6)
Comorbidities¹			
Yes	8432 (46.2)	8450 (46.0)	16,882 (46.1)
No	9810 (53.8)	9929 (54.0)	19,739 (53.9)

* All eligible randomized participants who receive all vaccination(s) as randomized within the predefined window, have no other important protocol deviations as determined by the clinician, and have no evidence of SARS-CoV-2 infection prior to 7 days after Dose 2.

† Includes multiracial and not reported.

¹ Number of subjects who have 1 or more comorbidities that increase the risk of severe COVID-19 disease: e.g., asthma, BMI ≥ 30 kg/m², chronic pulmonary disease, diabetes mellitus, hypertension.

- Chronic lung disease (e.g., emphysema and chronic bronchitis, idiopathic pulmonary fibrosis, and cystic fibrosis) or moderate to severe asthma
- Significant cardiac disease (e.g., heart failure, coronary artery disease, congenital heart disease, cardiomyopathies, and pulmonary hypertension)
- Severe obesity (body mass index ≥ 40 kg/m²)
- Diabetes (Type 1, Type 2 or gestational)
- Liver disease
- Human Immunodeficiency Virus (HIV) infection.

Efficacy Against COVID-19

The population included in the primary efficacy analysis included those 16 years of age and older who had been enrolled from X through Y date and followed for the development of Covid-19 through November 14, 2020. At the time of the analysis of the primary efficacy endpoint, participants 16 years of age and older had been followed for the development of symptomatic COVID-19 for a median of 2 months. Participants were followed for the development of symptomatic COVID-19 for 2214 person-years for Pfizer-BioNTech COVID-19 Vaccine and 2222 person-years in the placebo group. There were 8 confirmed COVID-19 cases identified in Pfizer-BioNTech COVID-19 Vaccine and 162 in placebo groups, respectively.

In this analysis, compared to placebo, efficacy of Pfizer-BioNTech COVID-19 Vaccine in participants from first COVID-19 occurrence from 7 days after Dose 2 without evidence of prior infection with SARS-CoV-2 was 95.0% (95% credible interval of 90.3% to 97.6%) in participants 16 years of age and older. In a separate analysis, compared to placebo, efficacy of Pfizer-BioNTech COVID-19 Vaccine in participants from first

Commented [A14]: Pfizer, please note that this revision is made for consistency with the shell tables sent on November 17, 2020.

Commented [A15]: Pfizer, please add the dates to this sentence.

Commented [A16]: Pfizer, please present the primary analyses in tabular form and include rows presenting subgroup analyses for subgroups 16-64 and > 65, while deleting the other subgroup analyses.

Please revise the text accordingly.

COVID-19 occurrence from 7 days after Dose 2 compared to participants with or without evidence of prior infection with SARS-CoV-2 was 94.6% (95% credible interval of 89.9% to 97.3%) in participants 16 years of age and older.

Commented [A17]: To be revised as per comment above.

Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19* (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhea; or vomiting).

*Case definition: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhea; or vomiting.

The subgroup analyses of vaccine efficacy including important demographic characteristics is presented in Table 6.

Table 6: Subgroup Analyses of Vaccine Efficacy - Participants Without Evidence of Infection Prior to 7 Days After Dose 2 - Evaluable Efficacy Population

Subgroup	Pfizer-BioNTech COVID-19 Vaccine N ^a =18,198 Cases n ^{1b} Surveillance Time ^c (n ^{2d})	Placebo N ^a =18,325 Cases n ^{1b} Surveillance Time ^c (n ^{2d})	Vaccine Efficacy % (95% CI) ^e
Age (years)			
16 to 64	TBD	TBD	TBD
≥65	1 0.508 (3848)	19 0.511 (3880)	94.7 (66.7, 99.9)
≥75	0 0.102 (774)	5 0.106 (785)	100.0 (-13.1, 100.0)
Sex			
Female	5 1.090 (8536)	81 1.114 (8749)	93.7 (84.7, 98.0)
Male	3 1.124 (8875)	81 1.108 (8762)	96.4 (88.9, 99.3)
Ethnicity			
Hispanic or Latino	3 0.605 (4764)	53 0.600 (4746)	94.4 (82.7, 98.9)
Not Hispanic or Latino	5 1.596 (12548)	109 1.608 (12661)	95.4 (88.9, 98.5)
Race			
Black or African American	0 0.165 (1502)	7 0.164 (1486)	100.0 (31.2, 100.0)
White	7 1.889 (14504)	146 1.903 (14670)	95.2 (89.8, 98.1)
All others ^f	1 0.160 (1405)	9 0.155 (1355)	89.3 (22.6, 99.8)

Table 6: Subgroup Analyses of Vaccine Efficacy - Participants Without Evidence of Infection Prior to 7 Days After Dose 2 - Evaluable Efficacy Population

Subgroup	Pfizer-BioNTech COVID-19 Vaccine N ^a =18,198 Cases n1 ^b Surveillance Time ^c (n2 ^d)	Placebo N ^a =18,325 Cases n1 ^b Surveillance Time ^c (n2 ^d)	Vaccine Efficacy % (95% CI) ^e

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Confidence interval (CI) for vaccine efficacy is derived based on the Clopper and Pearson method adjusted to the surveillance time.

f. All others = American Indian or Alaska Native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories.

The subgroup analyses of vaccine efficacy by risk status in participants is presented in Table 7.

Table 7: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Risk Status – Participants Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%) (95% CI) ^e	
	Pfizer-BioNTech COVID-19 Vaccine (N ^a =18198) n1 ^b Surveillance Time ^c (n2 ^d)	Placebo (N ^a =18325) n1 ^b Surveillance Time ^c (n2 ^d)				
First COVID-19 occurrence from 7 days after Dose 2						
At risk ^f						
Yes	4	1.025 (8030)	86	1.025 (8029)	95.3	(87.7, 98.8)
No	4	1.189 (9381)	76	1.197 (9482)	94.7	(85.9, 98.6)
Age group (years) and at risk						
16-64 and not at risk	4	0.962 (7671)	69	0.964 (7701)	94.2	(84.4, 98.5)
16-64 and at risk	3	0.744 (5878)	74	0.746 (5917)	95.9	(87.6, 99.2)
≥65 and not at risk	0	0.227 (1701)	7	0.233 (1771)	100.0	(29.0, 100.0)
≥65 and at risk	1	0.281 (2147)	12	0.279 (2109)	91.7	(44.2, 99.8)
Obese ^g						
Yes	3	0.763 (6000)	67	0.782 (6103)	95.4	(86.0, 99.1)
No	5	1.451 (11406)	95	1.439 (11404)	94.8	(87.4, 98.3)
Age group (years) and obese						
16-64 and not obese	4	1.107 (8811)	83	1.101 (8825)	95.2	(87.3, 98.7)
16-64 and obese	3	0.598 (4734)	60	0.609 (4789)	94.9	(84.4, 99.0)
≥65 and not obese	1	0.343 (2582)	12	0.338 (2567)	91.8	(44.5, 99.8)
≥65 and obese	0	0.165 (1265)	7	0.173 (1313)	100.0	(27.1, 100.0)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Table 7: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Risk Status – Participants Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

	Vaccine Group (as Randomized)					
	Pfizer-BioNTech COVID-19 Vaccine (N ^a =18198)		Placebo (N ^a =18325)			
Efficacy Endpoint Subgroup	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)	VE (%)	(95% CI ^e)

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.

f. At risk is defined as having at least one of the Charlson Comorbidity Index (CMI) category or obesity (BMI ≥30 kg/m²).

g. Obese is defined as BMI ≥30 kg/m².

19 HOW SUPPLIED/STORAGE AND HANDLING

Pfizer-BioNTech COVID-19 Vaccine Suspension for Intramuscular Injection, Multiple Dose Vials are supplied in a carton containing 25 multiple dose vials (NDC 59267-1000-3) or 195 multiple dose vials (NDC 59267-1000-2). ~~Not all pack sizes may be available.~~

Frozen Vials Prior to Use

Cartons of Pfizer-BioNTech COVID-19 Vaccine ~~Suspension for Intramuscular Injection~~-Multiple Dose Vials (after dilution each vial contains 5 doses of 0.3 mL) arrive in thermal containers with dry ice. Once received, remove the vial cartons immediately from the thermal container and store in ~~the~~ an ultra-low temperature freezer between -80°C to -60°C (-112°F to -76°F). Vials must be kept frozen between -80°C to -60°C (-112°F to -76°F) and protected from light, in the original cartons, until ready to use.

If an ultra-low temperature freezer is not available, the thermal container in which the Pfizer-BioNTech COVID-19 Vaccine arrives may be used as temporary storage when consistently re-filled to the top of the container with dry ice. Refer to the re-icing guidelines packed in the original thermal container for instructions regarding the use of the thermal container for temporary storage. The thermal container maintains a temperature range of -90°C to -60°C (-130°F to -76°F). Storage within this temperature range is not considered an excursion from the recommended storage condition.

Thawed Vials Prior to Dilution

Prior to dilution, multiple dose vials of Pfizer-BioNTech COVID-19 Vaccine may be thawed and stored in the refrigerator [2°C to 8°C (35°F to 46°F)]. ~~for up to 5 days (120 hours).~~ A carton of 25 vials or 195 vials may take up to 2 or 3 hours, respectively, to thaw in the refrigerator, ~~respectively~~, whereas a fewer number of vials will thaw in less time. ~~Vials may be stored in the refrigerator for up to 5 days (120 hours).~~ Frozen vials may also be

Commented [A18]: Pfizer, please insert a time frame, if needed, for how long it can be safely stored in the thermal container.
Are there any temperature monitors in the boxes?

thawed at room temperature [up to 25°C (-77°F)] for 30 minutes. During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.

Commented [A19]: Pfizer, this reflects the time in the Dosage and Administration section. Please modify as necessary.

Prior to dilution, the multiple dose vial may be stored at room temperature for no more than 2 hours. ~~During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.~~ Thawed vials can be handled in room light conditions.

Do not refreeze thawed vials ~~[see Dosage and Administration (2.1)].~~

Vials After Dilution

After dilution, multiple dose vials of the Pfizer-BioNTech COVID-19 Vaccine must be stored between 2°C to 25°C (35°F to 77°F) and used within 6 hours from the time of dilution. Any vaccine remaining in vials must be discarded after 6 hours. During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light. After dilution, the vaccine vials can be handled in room light conditions. Do not refreeze. ~~If the vaccine is frozen, it must be discarded.~~

20 PATIENT COUNSELING INFORMATION

Advise the recipient or caregiver to read the Fact Sheet for Recipients and Caregivers.

21 CONTACT INFORMATION

For general questions, visit the website or call the telephone number provided below.

Website	Telephone number
www.cvdvaccine.com 	1-877-829-2619 (1-877-VAX-CO19)

This Full EUA Prescribing Information may have been updated. For the most recent Full EUA Prescribing Information, please see www.cvdvaccine.com.



Manufactured by
Pfizer Inc., New York, NY 10017

BIONTECH

Manufactured for
BioNTech Manufacturing GmbH
An der Goldgrube 12
55131 Mainz, Germany

LAB-1457-0.4

Revised: November 2020

FACT SHEET FOR HEALTHCARE PROVIDERS ADMINISTERING VACCINE (VACCINATION PROVIDERS)

EMERGENCY USE AUTHORIZATION (EUA) OF THE PFIZER-BIONTECH COVID-19 VACCINE TO PREVENT CORONAVIRUS DISEASE 2019 (COVID-19)

The U.S. Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) to permit the emergency use of the unapproved product, **Pfizer-BioNTech COVID-19 Vaccine**, for active immunization to prevent COVID-19 in individuals 16 years of age and older.

Commented [A20]: The age range for the EUA will be pending our review.

SUMMARY OF INSTRUCTIONS FOR COVID-19 VACCINATION PROVIDERS

Vaccination providers enrolled in the federal COVID-19 Vaccination Program must report all vaccine administration errors, all serious adverse events, cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and cases of COVID-19 that result in hospitalization or death following administration of Pfizer-BioNTech COVID-19 Vaccine. See "MANDATORY REQUIREMENTS FOR THE PFIZER-BIONTECH COVID-19 VACCINE ADMINISTRATION UNDER EMERGENCY USE AUTHORIZATION" for reporting requirements.

- The Pfizer-BioNTech COVID-19 Vaccine is a suspension for intramuscular injection.
- After preparation, the Pfizer-BioNTech COVID-19 Vaccine is administered as a series of two doses (0.3 mL each) 21 days apart.

See the Full EUA ~~Full~~ Prescribing Information for instructions for preparation and administration.

For information on clinical trials that are testing the use of the Pfizer-BioNTech COVID-19 Vaccine for active immunization against COVID-19, please see www.clinicaltrials.gov.

This Fact Sheet may have been updated. For the most recent Fact Sheet, please see www.cvdvaccine.com.

DESCRIPTION OF COVID-19

Coronavirus disease 2019 (COVID-19) is an infectious disease caused by the novel coronavirus, SARS-CoV-2, that appeared in late 2019. It is predominantly a respiratory illness that can affect other organs. People with COVID-19 have reported a wide range of symptoms, ranging from mild symptoms to severe illness. Symptoms may appear 2-14 days after exposure to the virus. Symptoms may include: fever or chills; cough; shortness of breath; fatigue; muscle or body aches; headache; new loss of taste or smell; sore throat; congestion or runny nose; nausea or vomiting; diarrhea.

Commented [A21]: Pfizer, please insert page numbers (pagination) in the document.

DOSAGE AND ADMINISTRATION

Additionally, please include the date of the Fact Sheet version on each page.

Storage and Handling

Frozen Vials Prior to Use

Cartons of Pfizer-BioNTech COVID-19 Vaccine Suspension for Intramuscular Injection Multiple Dose Vials (after dilution each vial contains 5 doses of 0.3 mL) arrive in thermal containers with dry ice. Once received, remove the vial cartons immediately from the thermal container and store in the freezer between -80°C to -60°C (-112°F to -76°F). Vials must be kept frozen between -80°C to -60°C (-112°F to -76°F) and protected from light, in the original cartons, until ready to use.

If an ultra-low temperature freezer is not available, the thermal container in which the Pfizer-BioNTech COVID-19 Vaccine arrives may be used as temporary storage when consistently re-filled to the top of the container with dry ice. Refer to the re-icing guidelines packed in the original thermal container for instructions regarding the use of the thermal container for temporary storage. The thermal container maintains a temperature range of -90°C to -60°C (-130°F to -76°F). Storage within this temperature range is not considered an excursion from the recommended storage condition.

Thawed Vials Prior to Dilution

Prior to dilution, multiple dose vials of Pfizer-BioNTech COVID-19 Vaccine may be thawed and stored in the refrigerator [2°C to 8°C (35°F to 46°F)]. A carton of 25 vials or 195 vials may take up to 2 or 3 hours, respectively, to thaw in the refrigerator, whereas a fewer number of vials will thaw in less time. Vials may be stored in the refrigerator for up to 5 days (120 hours). Frozen vials may also be thawed at room temperature [up to 25°C (77°F)]. Prior to dilution, the multiple dose vial may be stored at room temperature for no more than 2 hours. During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light. Thawed vials can be handled in room light conditions. Do not refreeze thawed vials (see *Dose Preparation*).

Vials after Dilution

After dilution, multiple dose vials of the Pfizer-BioNTech COVID-19 Vaccine must be stored between 2°C to 25°C (35°F to 77°F) and used within 6 hours from the time of dilution. Any vaccine remaining in vials must be discarded after 6 hours. During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light. After dilution, the vaccine vials can be handled in room light conditions. Do not freeze. If the vaccine is frozen, it must be discarded.

Dosing and Schedule

The Pfizer-BioNTech COVID-19 Vaccine is administered intramuscularly as a series of two doses (0.3 mL each) 21 days apart.

There are no data available on the interchangeability of the Pfizer-BioNTech COVID-19 Vaccine with other COVID-19 vaccines to complete the vaccination series. Individuals who have received one dose of Pfizer-BioNTech COVID-19 Vaccine should receive a second dose of Pfizer-BioNTech COVID-19 Vaccine to complete the vaccination series.

Dose Preparation

- The Pfizer-BioNTech COVID-19 Vaccine Multiple Dose Vial contains a frozen suspension that does not contain preservative and must be thawed and diluted prior to administration.
- Vials may be thawed in the refrigerator or at room temperature [up to 25°C (77°F)].
- Prior to dilution, the thawed suspension may contain white to off-white opaque amorphous particles.
- The contents of the vial must be diluted with 1.8 mL of sterile 0.9% Sodium Chloride Injection, USP to form the Pfizer-BioNTech COVID-19 Vaccine.
- After dilution, the vial contains 5 doses of 0.3 mL.
- 0.9% Sodium Chloride Injection, USP is not packaged with the vaccine and must be sourced separately. Do not use bacteriostatic 0.9% Sodium Chloride Injection or any other diluent.
- After dilution, the vaccine will be an off-white suspension. Inspect vials to confirm there are no particulates and no discoloration is observed.
- Strict adherence to aseptic techniques must be followed.








No more than
2 hours at room
temperature
(up to 25 °C / 77 °F)


- Remove a thawed vial of Pfizer-BioNTech COVID-19 Vaccine from the refrigerator and allow it to come to room temperature.
- If using a frozen vial of Pfizer-BioNTech COVID-19 Vaccine, thaw for 30 minutes at room temperature.

Vials at room temperature must be diluted within 2 hours.

Commented [A22]: Pfizer, please make the directions in each panel consistent with the Full EUA Prescribing Information.

 Gently x 10	• Invert gently 10 times to mix. • <u>Do not shake.</u>
 1.8 mL of 0.9% sodium chloride injection	• Obtain sterile 0.9% Sodium Chloride Injection, USP. • Cleanse the vial stopper with a single-use antiseptic swab. • Add 1.8 mL of 0.9% Sodium Chloride Injection, USP into the Pfizer-BioNTech COVID-19 Vaccine vial using a needle 21-gauge or narrower.
 Pull back plunger to 1.8 mL to remove air from vial	• Equalize vial pressure before removing the needle from the vial by withdrawing 1.8 mL air into the empty diluent syringe.
 Gently x 10	• Gently invert the vial containing the Pfizer-BioNTech COVID-19 Vaccine 10 times to mix. • <u>Do not shake.</u>

	- Record the date and time of dilution on the Pfizer-BioNTech COVID-19 Vaccine vial label. - Store between 2°C to 25°C (35°F to 77°F). - Discard any unused vaccine 6 hours after dilution.
---	---

PREPARATION OF INDIVIDUAL 0.3 mL DOSES OF PFIZER-BIONTECH COVID-19 VACCINE	
	- Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab, and withdraw <u>0.3 mL</u> of the Pfizer-BioNTech COVID-19 Vaccine. - Administer immediately.

Administration

Visually inspect each dose in the dosing syringe prior to administration. The vaccine will be an off-white suspension. During the visual inspection,

- verify the final dosing volume of 0.3 mL.
- confirm there are no particulates and that no discoloration is observed.

If the visual inspection fails, do not administer the vaccine.

Administer the Pfizer-BioNTech COVID-19 Vaccine intramuscularly in the deltoid muscle.

Contraindications

Do not administer Pfizer-BioNTech COVID-19 Vaccine to individuals with known history of a severe allergic reaction (e.g., anaphylaxis) to any component of the Pfizer-BioNTech COVID-19 Vaccine (see [Full EUA Full Prescribing Information](#)).

Warnings

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of Pfizer-BioNTech COVID-19 Vaccine.

~~The administration of the Pfizer BioNTech COVID-19 Vaccine should be postponed in individuals with an acute severe febrile illness.~~

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the Pfizer-BioNTech COVID-19 Vaccine.

Pfizer-BioNTech COVID-19 Vaccine may not protect all vaccine recipients.

Pregnancy and Lactation

~~There is no experience with the use of the Pfizer BioNTech COVID-19 Vaccine during pregnancy or breastfeeding. Therefore, the Pfizer BioNTech COVID-19 Vaccine is not recommended for use during pregnancy or breastfeeding.~~

Adverse Reactions

Adverse reactions that have been reported for the Pfizer-BioNTech COVID-19 Vaccine are injection site pain, fatigue, headache, muscle pain, chills, joint pain, fever, and lymphadenopathy (see [Full](#) ~~EUA~~ [Full](#) Prescribing Information).

Additional adverse reactions, some of which may be serious, may become apparent with more widespread use of the Pfizer-BioNTech COVID-19 Vaccine.

Use with Other Vaccines

There is no information on the co-administration of the Pfizer-BioNTech COVID-19 Vaccine with other vaccines.

INFORMATION TO PROVIDE TO VACCINE RECIPIENTS/CAREGIVERS

As the vaccination provider, you must communicate to the recipient or their caregiver, information consistent with the "Fact Sheet for Recipients and Caregivers" (and provide a copy or direct the individual to the website www.cvdvaccine.com to obtain the Fact Sheet) prior to the individual receiving Pfizer-BioNTech COVID-19 Vaccine, including:

- FDA has authorized the emergency use of the Pfizer-BioNTech COVID-19 Vaccine, which is not an FDA-approved vaccine.
- The recipient or their caregiver has the option to accept or refuse Pfizer-BioNTech COVID-19 Vaccine.

- The significant known and potential risks and benefits of Pfizer-BioNTech COVID-19 Vaccine, and the extent to which such risks and benefits are unknown.
- Information about available alternative vaccines and the risks and benefits of those alternatives.

For information on clinical trials that are testing the use of the Pfizer-BioNTech COVID-19 Vaccine to prevent COVID-19, please see www.clinicaltrials.gov.

Provide a vaccination card to the recipient or their caregiver with the date when the recipient needs to return for the second dose of Pfizer-BioNTech COVID-19 Vaccine.

MANDATORY REQUIREMENTS FOR PFIZER-BIONTECH COVID-19 VACCINE ADMINISTRATION UNDER EMERGENCY USE AUTHORIZATION

In order to mitigate the risks of using this unapproved product under EUA and to optimize the potential benefit of Pfizer-BioNTech COVID-19 Vaccine, the following items are required. Use of unapproved Pfizer-BioNTech COVID-19 Vaccine for active immunization to prevent COVID-19 under this EUA is limited to the following (all requirements **must** be met):

1. Pfizer-BioNTech COVID-19 Vaccine is authorized for use in individuals 16 years of age and older.
2. The vaccination provider must communicate to the individual receiving the Pfizer-BioNTech COVID-19 Vaccine or their caregiver, information consistent with the "Fact Sheet for Recipients and Caregivers" prior to the individual receiving Pfizer-BioNTech COVID-19 Vaccine.
3. The vaccination provider is responsible for mandatory reporting of the following to the Vaccine Adverse Event Reporting System (VAERS) and to Pfizer Inc.:
 - vaccine administration errors whether or not associated with an adverse event,
 - serious adverse events* (irrespective of attribution to vaccination),
 - cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and
 - cases of COVID-19 that result in hospitalization or death.

Complete and submit reports to VAERS online at <https://vaers.hhs.gov/reportevent.html> or by calling 1-800-822-7967. The reports should include the words "Pfizer-BioNTech COVID-19 Vaccine EUA" in the description section of the report.

Report to Pfizer Inc. using the contact information below or by providing a copy of the VAERS form to Pfizer Inc.

<u>Website</u>	<u>Fax number</u>	<u>Telephone number</u>
www.pfizersafetyreporting.com	1-866-635-8337	1-800-438-1985

4. The vaccination provider is responsible for responding to FDA requests for information about vaccine administration errors, adverse events, cases of MIS in adults and children and cases of COVID-19 that result in hospitalization or death following administration of Pfizer-BioNTech COVID-19 Vaccine to recipients.

Commented [A23]: Pending review.

* Serious adverse events are defined as:

- Death;
- A life-threatening adverse event;
- Inpatient hospitalization or prolongation of existing hospitalization;
- A persistent or significant incapacity or substantial disruption of the ability to conduct normal life functions;
- A congenital anomaly/birth defect;
- An important medical event that based on appropriate medical judgement may jeopardize the individual and may require medical or surgical intervention to prevent one of the outcomes listed above.

REPORTING OF OTHER REPORTING REQUIREMENTS ADVERSE EVENTS TO VAERS AND PFIZER, INC.

Vaccination providers may report to VAERS and Pfizer Inc. other adverse events that are not required to be reported using the contact information above.

~~In addition, you must report all adverse events, vaccine administration errors whether or not associated with an adverse event, serious adverse events* (irrespective of attribution to vaccination), cases of Multisystem Inflammatory Syndrome (MIS), and cases of COVID-19 that result in hospitalization or death to Pfizer Inc. at the contact information below or by providing a copy of the VAERS form by faxing to Pfizer Inc.~~

Website	Fax number	Telephone number
www.pfizersafetyreporting.com	1-866-635-8337	1-800-438-1985

ADDITIONAL INFORMATION

For general questions, visit the website or call the telephone number provided below.

To access the most recent Pfizer-BioNTech COVID-19 Vaccine ~~Full EUA~~ Full Prescribing Information and Fact Sheets, please scan the QR code provided below.

Global website	Telephone number
www.cvdvaccine.com 	1-877-829-2619 (1-877-VAX-CO19)

AVAILABLE ALTERNATIVES

There is no approved alternative vaccine to prevent COVID-19. There may be clinical trials or availability under EUA of other COVID-19 vaccines.

AUTHORITY FOR ISSUANCE OF THE EUA

The Secretary of Health and Human Services (HHS) has declared a public health emergency that justifies the emergency use of drugs and biological products during the Pfizer-BioNTech COVID-19 pandemic Vaccine to prevent COVID-19. In response, FDA has issued an EUA for the unapproved product, Pfizer-BioNTech COVID-19 Vaccine, for active immunization against COVID-19 in individuals 16 years of age and older.

~~As a vaccination provider, you must comply with the mandatory requirements of the EUA (see above).~~

FDA issued this EUA, based on Pfizer-BioNTech's request and submitted data.

Although limited scientific information is available, based on the totality of the scientific evidence available to date, it is reasonable to believe that the Pfizer-BioNTech COVID-19 Vaccine may be effective for the prevention of COVID-19 in individuals as specified in the Full EUA Full Prescribing Information.

Commented [A24]: Pfizer, please append the Full EUA Prescribing Information to this Fact Sheet to make one document.

This EUA for the Pfizer-BioNTech COVID-19 Vaccine will end when the Secretary of HHS determines that the circumstances justifying the EUA no longer exist or when there is a change in the approval status of the product such that an EUA is no longer needed.

For additional information about Emergency Use Authorization visit FDA at:
<https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization>.

The Countermeasures Injury Compensation Program

The Countermeasures Injury Compensation Program (CICP) is a federal program that has been created to help pay for related costs of medical care and other specific expenses to compensate people injured after use of certain medical countermeasures. Medical countermeasures are specific vaccines, medications, devices, or other items used to prevent, diagnose, or treat the public during a public health emergency or a security threat. For more information about CICP regarding the Pfizer-BioNTech COVID-19 Vaccine used to prevent COVID-19, visit www.hrsa.gov/cicp, email cicp@hrsa.gov, or call: 1-855-266-2427.



Manufactured by
Pfizer Inc., New York, NY 10017

BIONTECH

Manufactured for
BioNTech Manufacturing GmbH
An der Goldgrube 12
55131 Mainz, Germany

LAB-1450-0.4

Revised: November 2020

END SHORT VERSION FACT SHEET

Long Version (Full EUA Prescribing Information) Begins On Next Page

FACT SHEET FOR RECIPIENTS AND CAREGIVERS

EMERGENCY USE AUTHORIZATION (EUA) OF THE PFIZER-BIONTECH COVID-19 VACCINE TO PREVENT CORONAVIRUS DISEASE 2019 (COVID-19) IN INDIVIDUALS 16 YEARS OF AGE AND OLDER

Commented [A25]: The age range for the EUA will be pending our review.

You are being offered the Pfizer-BioNTech COVID-19 Vaccine to prevent Coronavirus Disease 2019 (COVID-19) caused by SARS-CoV-2. This Fact Sheet contains information to help you understand the risks and benefits of the Pfizer-BioNTech COVID-19 Vaccine, which you may receive because there is currently a pandemic of COVID-19.

The Pfizer-BioNTech COVID-19 Vaccine is a vaccine and may prevent you from getting COVID-19. There is no U.S. Food and Drug Administration (FDA) approved vaccine to prevent COVID-19.

Read this Fact Sheet for information about the Pfizer-BioNTech COVID-19 Vaccine. Talk to the vaccination provider if you have questions. It is your choice to receive the Pfizer-BioNTech COVID-19 Vaccine.

The Pfizer-BioNTech COVID-19 Vaccine is administered as a 2-dose series, 21 days apart, into the muscle of the upper arm.

The Pfizer-BioNTech COVID-19 Vaccine may not protect everyone.

This Fact Sheet may have been updated. For the most recent Fact Sheet, please see www.cvdvaccine.com.

WHAT YOU NEED TO KNOW BEFORE YOU GET THIS VACCINE?

WHAT IS COVID-19?

COVID-19 disease is caused by a coronavirus called SARS-CoV-2. This type of coronavirus has not been seen before. You can get COVID-19 through contact with another person who has the virus. It is predominantly a respiratory illness that can affect other organs. People with COVID-19 have had a wide range of symptoms reported, ranging from mild symptoms to severe illness. Symptoms may appear 2-14 days after exposure to the virus. Symptoms may include: fever or chills; cough; shortness of breath; fatigue; muscle or body aches; headache; new loss of taste or smell; sore throat; congestion or runny nose; nausea or vomiting; diarrhea.

WHAT IS THE PFIZER-BIONTECH COVID-19 VACCINE?

The Pfizer-BioNTech COVID-19 Vaccine is an investigational (not FDA-approved) vaccine that may prevent COVID-19. There is no FDA-approved vaccine to prevent COVID-19.

Commented [A26]: Pfizer, please insert page numbers (pagination) in the document.

The FDA has authorized the emergency use of the Pfizer-BioNTech COVID-19 Vaccine to prevent COVID-19 in individuals 16 years of age and older under an Emergency Use Authorization (EUA).

For more information on EUA, see the “What is an Emergency Use Authorization (EUA)?” section at the end of this Fact Sheet.

WHAT SHOULD YOU MENTION TO YOUR VACCINATION PROVIDER BEFORE YOU GET THE PFIZER-BIONTECH COVID-19 VACCINE?

Tell the vaccination provider about all of your medical conditions, including if you:

- have any allergies
- have a fever
- have a bleeding disorder or are on a blood thinner
- are immunocompromised or are on a medicine that affects your immune system
- are pregnant or plan to become pregnant
- are breastfeeding
- have received another COVID-19 vaccine

WHO SHOULD GET THE PFIZER-BIONTECH COVID-19 VACCINE?

FDA has authorized the emergency use of the Pfizer-BioNTech COVID-19 Vaccine in individuals 16 years of age and older.

WHO SHOULD NOT GET THE PFIZER-BIONTECH COVID-19 VACCINE?

You should not get the Pfizer-BioNTech COVID-19 Vaccine if you:

- had a severe allergic reaction after a previous dose of this vaccine
- had a severe allergic reaction to any ingredient of this vaccine (see **WHAT ARE THE INGREDIENTS IN THE PFIZER-BIONTECH COVID-19 VACCINE?** at the end of this Fact Sheet)

HOW IS THE PFIZER-BIONTECH COVID-19 VACCINE GIVEN?

The Pfizer-BioNTech COVID-19 Vaccine will be given to you as an injection into the muscle of your upper arm.

The Pfizer-BioNTech COVID-19 Vaccine vaccination series is 2 doses given 21 days apart.

If you receive one dose of the Pfizer-BioNTech COVID-19 Vaccine, you should receive a second dose of the same vaccine 21 days later to complete the vaccination series.

HAS THE PFIZER-BIONTECH COVID-19 VACCINE BEEN USED SAFELY BEFORE?

The Pfizer-BioNTech COVID-19 Vaccine is an investigational COVID-19 vaccine. In clinical trials, approximately 20,000 individuals 16 years of age and older have received at least 1 dose of the Pfizer-BioNTech COVID-19 Vaccine.

Commented [A27]: Pending review...

WHAT ARE THE BENEFITS OF THE PFIZER-BIONTECH COVID-19 VACCINE?

In an ongoing clinical trial, the Pfizer-BioNTech COVID-19 Vaccine has been shown to prevent COVID-19 following 2 doses given 21 days apart. The duration of protection against COVID-19 is currently unknown.

WHAT ARE THE RISKS OF THE PFIZER-BIONTECH COVID-19 VACCINE?

Side effects that have been reported with the Pfizer-BioNTech COVID-19 Vaccine are:

- injection site pain
- tiredness
- headache
- muscle pain
- chills
- joint pain
- fever
- swollen lymph nodes (lymphadenopathy)

These may not be all the possible side effects of the Pfizer-BioNTech COVID-19 Vaccine. Serious and unexpected side effects may occur. The Pfizer-BioNTech COVID-19 Vaccine is still being studied in clinical trials.

WHAT IF I DECIDE NOT TO GET THE PFIZER-BIONTECH COVID-19 VACCINE?

It is your choice to receive or not receive the Pfizer-BioNTech COVID-19 Vaccine. Should you decide not to receive it, it will not change your standard medical care.

ARE OTHER CHOICES AVAILABLE FOR PREVENTING COVID-19 BESIDES PFIZER-BIONTECH COVID-19 VACCINE?

Currently, there is no FDA-approved alternative vaccine available for prevention of COVID-19. FDA may allow the emergency use of other vaccines to protect against COVID-19.

CAN I RECEIVE THE PFIZER-BIONTECH COVID-19 VACCINE WITH OTHER VACCINES?

There is no information on the use of the Pfizer-BioNTech COVID-19 Vaccine with other vaccines.

WHAT IF I AM PREGNANT OR BREASTFEEDING?

~~There is no experience giving the Pfizer-BioNTech COVID-19 Vaccine to pregnant or breastfeeding women.~~ If you are pregnant or breastfeeding, discuss your options with your healthcare provider.

Commented [A28]: Pfizer, please delete.

WILL THE PFIZER-BIONTECH COVID-19 VACCINE GIVE ME COVID-19?

No. The Pfizer-BioNTech COVID-19 Vaccine does not contain SARS-CoV-2 and cannot give you COVID-19.

KEEP YOUR VACCINATION CARD

When you get your first dose, you will get a vaccination card to show you when to return for your second dose of Pfizer-BioNTech COVID-19 Vaccine. Remember to bring your card when you return.

SERIOUS SIDE EFFECTS

There is a remote chance that the Pfizer-BioNTech COVID-19 Vaccine could cause a severe allergic reaction. A severe allergic reaction would usually occur within a few minutes to one hour after getting a dose of the Pfizer-BioNTech COVID-19 Vaccine. Signs of a severe allergic reaction can include:

- Difficulty breathing

- Swelling of your face and throat
- A fast heartbeat
- A bad rash all over your body
- Dizziness and weakness

WHAT SHOULD I DO ABOUT SIDE EFFECTS?

If you experience a severe allergic reaction, call 9-1-1, or go to the nearest hospital.

Call the vaccination provider or your healthcare provider if you have any side effects that bother you or do not go away.

Report vaccine side effects to **FDA/CDC Vaccine Adverse Event Reporting System (VAERS)**. The VAERS toll-free number is 1-800-822-7967 or report online to <https://vaers.hhs.gov/reportevent.html>. Please include "Pfizer BioNTech COVID-19 Vaccine EUA" in the first line of box #18 of the report form.

In addition, you can report side effects to Pfizer Inc. at the contact information provided below.

Website	Fax number	Telephone number
www.pfizersafetyreporting.com	1-866-635-8337	1-800-438-1985

ADDITIONAL INFORMATION

If you have questions, visit the website or call the telephone number provided below.

To access the most recent Fact Sheets, please scan the QR code provided below.

Global website	Telephone number
www.cvdvaccine.com 	1-877-829-2619 (1-877-VAX-CO19)

WHAT ARE THE INGREDIENTS IN THE PFIZER-BIONTECH COVID-19 VACCINE?

The Pfizer-BioNTech COVID-19 Vaccine includes the following ingredients: mRNA, lipids ((4-hydroxybutyl)azanediyl)bis(hexane-6,1-diyl)bis(2-hexyldecanoate), 2-[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide, 1,2-Distearoyl-sn-glycero-3-phosphocholine, and cholesterol), potassium chloride, monobasic potassium phosphate, sodium chloride, dibasic sodium phosphate dihydrate, and sucrose.

HOW CAN I LEARN MORE?

- Ask the vaccination provider.
- Visit CDC at <https://www.cdc.gov/coronavirus/2019-ncov/index.html>.
- Visit FDA at <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization>.
- Contact your local or state public health department.

WHERE WILL MY VACCINATION INFORMATION BE RECORDED?

The vaccination provider may include your vaccination information in your state/local jurisdiction's Immunization Information System (IIS) or other designated system. This will ensure that you receive the same vaccine when you return for the second dose. For more information about IISs visit: <https://www.cdc.gov/vaccines/programs/iis/about.html>.

WHAT IS THE COUNTERMEASURES INJURY COMPENSATION PROGRAM?

The Countermeasures Injury Compensation Program is a federal program that may help pay for costs of medical care and other specific expenses of certain people who have been seriously injured by certain medicines or vaccines, including this vaccine. Generally, a claim must be submitted to the CICI within one (1) year from the date of receiving the vaccine. To learn more about this program, visit www.hrsa.gov/cicp/ or call 1-855-266-2427. The Countermeasures Injury Compensation Program is a federal program that may help pay for costs of medical care and other specific expenses of certain people who have been seriously injured by certain medicines or vaccines. To learn more about this program, visit www.hrsa.gov/cicp/ or call 1-855-266-2427.

WHAT IS AN EMERGENCY USE AUTHORIZATION (EUA)?

The United States FDA has made the Pfizer-BioNTech COVID-19 Vaccine available under an emergency access mechanism called an EUA. The EUA is supported by a Secretary of Health and Human Service (HHS) declaration that circumstances exist to justify the emergency use of drugs and biological products during the COVID-19 pandemic.

The Pfizer-BioNTech COVID-19 Vaccine has not undergone the same type of review as an FDA-approved or cleared product. FDA may issue an EUA when certain criteria are met, which includes that there are no adequate, approved, available alternatives. In addition, the FDA decision is based on the totality of scientific evidence available showing that the product may be effective to prevent COVID-19 during the COVID-19 pandemic and that the known and potential benefits of the product outweigh the known and potential risks of the product. All of these criteria must be met to allow for the product to be used in the treatment of patients during the COVID-19 pandemic.

The EUA for the Pfizer-BioNTech COVID-19 Vaccine is in effect for the duration of the COVID-19 EUA declaration justifying emergency use of these products, unless terminated or revoked (after which the products may no longer be used).



Manufactured by
Pfizer Inc., New York, NY 10017

BIONTECH

Manufactured for
BioNTech Manufacturing GmbH
An der Goldgrube 12
55131 Mainz, Germany

LAB-1451-0.4

Revised: November 2020

Commented [A29]: Pfizer, please include the 2-D bar code requested by CDC in the lower right corner of the last page of this Fact Sheet. The box with the barcode will include the following language:
Scan to capture that this Fact Sheet was provided to the vaccine recipient for the electronic medical record/immunization information system.