

Pfizer-BioNTech COVID-19 Vaccine

EUA 27034

**Response to CBER Comments Received on
24 November 2020, and Subsequent Teleconference on 25 November 2020,
Regarding Datasets in the
Request for Emergency Use Authorization Submission**

November 2020

TABLE OF CONTENTS

1. INTRODUCTION	3
2. CBER COMMENTS FROM 24 NOVEMBER 2020, AND SUBSEQUENT DISCUSSION WITH PFIZER VIA TELECONFERENCE ON 25 NOVEMBER 2020.....	3
2.1. CBER Comment 1.....	3
2.2. CBER Comment 2.....	3
2.3. CBER Comment 3.....	4
2.4. CBER Comment 4.....	5
2.5. CBER Comment 5.....	5
2.6. CBER Comment 6.....	6
2.7. CBER Comment 7.....	7
2.8. CBER Comment 8.....	8
2.9. CBER Comment 9.....	8
2.10. CBER Comment 10.....	8
2.11. CBER Comment 11.....	10
2.12. CBER Comment 12.....	12

1. INTRODUCTION

Reference is made to Request for Emergency Use Authorization (EUA 27034) submitted 20 November 2020 by Pfizer on behalf of BioNTech. Further reference is made to CBER's 24 November 2020 discussion points regarding datasets communicated by email from (b) (6) (CBER), to Elisa Harkins (Pfizer, Inc.), and subsequent discussion between CBER and Pfizer during a teleconference held 25 November 2020.

CBER comments are presented in ***bold italics*** and Pfizer-BioNTech's responses follow in plain text.

Per the discussion on 25 November 2020, at CBER's request, we are providing annotated, mock tables to facilitate review. See [Appendix 1](#).

2. CBER COMMENTS FROM 24 NOVEMBER 2020, AND SUBSEQUENT DISCUSSION WITH PFIZER VIA TELECONFERENCE ON 25 NOVEMBER 2020

2.1. CBER Comment 1

We are not able to identify all of the subjects included in your Safety population (N=37,586) and in your All Enrolled population (N=43,448).

Response

The core safety population in the EUA submission included 37,706 subjects 16 years of age or older enrolled by 09 October 2020 in the Phase 2/3 portion of the study who received at least one dose of vaccination. Note that HIV-positive participants were included in the safety population and shown as part of the study demographics and disposition, but did not contribute to the safety analyses. Excluding HIV-positive subjects, the total number of subjects in the core safety analysis was 37,586. Please consider ADSL.SAFFL='Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL="Y" and ADSL.HIVFL ne 'Y' and Not (ADSL.SAFFL='Y' and (ADSL.VAX101=' ' and ADSL.VAX102=' ')) to select the subjects for the core safety analyses.

The all enrolled safety population included 43,448 subjects 16 years of age or older in Phase 2/3 who received at least one dose of vaccination. This number included HIV positive subjects (ADSL.HIVFL='Y'). Excluding HIV positive subjects, the total number of subjects in the all enrolled safety analyses was 43,252.

2.2. CBER Comment 2

On page 74 of the document titled "Emergency Use Authorization (EUA) Request for Pfizer-BioNTech COVID-19 Vaccine" (Section m1.19 of EUA 27034.0) you have indicated that there are 37,706 participants who have a median follow-up of 2 months after Dose 2 and 37,586 participants (presented in Table 19 on page 75) who had a median of at least 2

months of follow-up after Dose 2 (including those analyzed in Phase 2). We have not been able to reproduce these values from your datasets.

Response

As clarified above, the core safety population in the EUA submission included 37,706 Phase 2/3 subjects 16 years of age or older, but a total of 37,586 subjects (excluding 120 HIV positive subjects) contributed to the safety analyses. Please use the combination of Safety Flag (ADSL.SAFFL='Y') and Phase 3 30k flag (ADSL.DS30KFL='Y') to select the core safety population and use ADSL.HIVFL ne 'Y' to exclude HIV positive subjects for safety analyses.

2.3. CBER Comment 3

In addition to recreating the study populations in comments 1 and 2, we would like to understand the reasons for these differences and your rationale for excluding various subjects/populations from your analyses.

Response

The eSub data submitted with the EUA included all data by the cutoff date of 14 November 2020 from all Phase 1/2/3 subjects enrolled in the study. The main safety analyses for Phase 2/3 in the EUA excluded the following subjects/subsets:

- Phase 1 subjects (ADSL.PHASEN=1): AEs and SAEs for Phase 1 subjects BNT162b2 30 µg group were summarized separately. Phase 1 subjects BNT162b2 30 µg group were also included in the listings.
- HIV positive subjects (ADSL.HIVFL='Y'): According to the Protocol, HIV-positive subjects in Phase 3 will not be included in analyses of the overall study objectives, with the exception of the specific exploratory objective for this group. HIV-positive subjects were included in the safety population and shown as part of the study demographics and disposition but not included in the safety analyses.
- Subjects 12-15 years of age (ADSL.AGEGR1N = 1): The intended EUA indication and primary focus of the analyses was for subjects 16 years of age or older. Reactogenicity data for 12-15 years of age were summarized separately as supportive information.
- Subjects with special data issues noted in ADRG:
 - Multi-enrolled subjects (ADSL.MULENRFL = 'Y'). During the conduct of this study, 1 subject was identified to be randomized across 2 sites in this trial. Because the significant misconduct of this subject compromised the integrity of the study data, results from this subject were excluded entirely from all efficacy and safety analyses, including disposition and demographic tabulations. Separated listings were generated for this subject.

- b. Three subjects (ADSL.SAFFL='Y' and (ADSL.VAX101=' ' and ADSL.VAX102=' ')) with unknown actual vaccination information due to medication error. Subjects were vaccinated as per CRF, but due to lack of matching actual vaccination data, these were not assigned to any actual dosing group. These subjects were excluded from all tables/figures, but included in all listings.

2.4. CBER Comment 4

Please confirm total number of subjects who missed their second vaccination and indicate whether the full safety data from these subjects were submitted, and in which dataset(s).

Response

There are 2449 subjects who did not have vaccination 2 in the EUA database. These included subjects who were recently enrolled and had not reached vaccination 2 yet and those who withdrew from treatment before vaccination 2. At the time of data extraction, the study was still ongoing and data cleaning is still underway. Updated data will be included for future reporting events and at study completion.

The full safety data for all the enrolled subjects, including subjects who did not receive the second vaccination, were submitted in ADAE, ADCEVD and ADFACEVD.

2.5. CBER Comment 5

Suspected cases of COVID entered in CE, such as the case of subject C4591001 1231 12312982, which were not confirmed and whose relatedness to vaccine cannot be ruled out, should be reclassified as SAEs in ADAE. Please discuss the number of subjects that would meet the criteria to be reclassified.

Response

Following internal review of potential COVID-19 illness events and their sequelae, any SAE determined NOT to meet endpoint criteria is reported back to the investigator site of incidence for reporting as an SAE.

The example subject does have an SAE record in the AE domain for suspected COVID-19 which is marked as not related, as assessed by the investigator.

There are >3000 subjects with potential COVID-19 illness reported (ie, with protocol-specified symptoms that could represent COVID-19 and should therefore trigger collection of a swab for SARS-CoV-2 NAAT) but were not confirmed to have COVID-19, as per the primary endpoint case flag/CDECASE.

In accordance with protocol section 8.3.7, potential COVID-19 illnesses and their sequelae that are consistent with the clinical endpoint definition were not to be recorded as AEs. These data were to be captured as efficacy assessment data only on the relevant pages of the CRF.

Potential COVID-19 illnesses and their sequelae were not to be reported according to the standard process for expedited reporting of SAEs, even though the event may meet the definition of an SAE, except if:

- the event is, in the investigator's opinion, of greater intensity, frequency, or duration than expected for the individual participant, or
- the investigator considers that there is a reasonable possibility that the event was related to study intervention.

2.6. CBER Comment 6

Using your flags (safety population, randomized population and enrolled population), we get the following numbers:

<i>Enrolled:</i>	43850
<i>Are Eligible for the study at Randomization:</i>	43780
<i>Randomized:</i>	43848
<i>Randomized + Eligible for Study at Randomization</i>	43780
<i>Safety Pop:</i>	43748
<i>Phase 3 30k Subjects</i>	37801
<i>Safety Pop + Eligible for study at Rand:</i>	43714
<i>Phase 3 30k Subjects + Safety Pop flag:</i>	37711
<i>Phase 3 30k Subjects + Safety + Eligible for study at Randomization:</i>	37682

We have tried using many other variables in combination and we are not able to match your reported numbers.

Response

Population	CBER	Pfizer Reporting	Difference	Explanation
Enrolled	43850	Confirmed	0	43850 subjects in ADSL
Randomized	43848	43548	300	We are excluding 195 Phase 1 subjects (ADSL.PHASEN = 1), 100 subjects from 12-15 Years age group (ADSL.AGEGR1N=1), 2 Subject(s) with enrollment at multiple sites (ADSL. MULENRFL= 'Y') and 3 subjects with unknown actual vaccination information (ADSL.SAFFL ='Y' and (ADSL. VAX101=""= ' ' and VAX102=""))) from Phase 2/3 disposition summary.
Eligible for study at randomization	43780	Confirmed	0	we have 68 subjects who are randomized and not meeting eligibility criteria (ADSL.RANDFL = 'Y' AND ADSL.INCL1FL = 'N'). 2 subjects did not get correct randomized number assigned and are

Population	CBER	Pfizer Reporting	Difference	Explanation
				not part of randomization population (ADSL.ARMCD = 'NOTASSGN').
Safety Population	43748	43448	300	We are excluding 195 Phase 1 subjects, 100 subjects from 12-15 Years age group, 2 Subject(s) with enrollment at multiple sites and 3 subjects with unknown actual vaccination information from Phase 2/3 safety population summary.
Safety Pop + Eligible for study at Rand	43714	Confirmed	0	we have 34 subjects who are part of safety pop but not meeting eligibility criteria (ADSL.SAFFL = 'Y' AND INCL1FL = 'N')
Phase 3 30k Subjects	37801	37796	5	37796 is the list of subjects with randomization <=9 OCT. We are excluding 2 subject(s) with enrollment at multiple sites and 3 subjects with unknown actual vaccination information.
Phase 3 30k Subject + Safety Pop flag	37711	37706	5	We are excluding 2 subject(s) with enrollment at multiple sites and 3 subjects with unknown actual vaccination information.
Phase 3 30k Subjects + Safety + Eligible for study at Randomization	37682	Confirmed	0	We have 29 subjects who are part of ~30k population, and part of safety population and not meeting eligibility criteria (ADSL.SAFFL = 'Y AND DS30KFL = "Y" AND INCL1FL = 'N')

2.7. CBER Comment 7

Regarding the HIV+ population:

Using Enrolled Population there are 196 HIV+ subjects (ADSL.HIVFL = 'Y')

Using randomized flag there are 196 HIV+ subjects

Using Safety population flag alone there are 196 HIV+ subjects

However, using combination of Safety Flag (ADSL.SAFFL = 'Y') and Phase 3 30k flag (ADSL.DS30KFL = 'Y'), there are only 120 HIV+ subjects.

Response

These two numbers match ours. There are 76 HIV subjects enrolled after 09Oct2020 who are and not included in the 30k Phase 3 subjects.

2.8. CBER Comment 8

In your final analysis datasets, there are numerous records in the CE and AE datasets missing start dates.

Response

For CE, as noted in the Response to FDA Information Request (BB-IND 19736) sent on 12NOV2020:

As start and stop dates were not collected for each potential COVID-19 symptom individually, CESTDTC and CEENDTC will not be populated for each symptom, but the date first symptom started and date last symptom resolved will be mapped to CESTDTC and CEENDTC in the “COVID-19 like illness” and “COVID-19 confirmed” records.

Hence CESTDTC and CEENDTC are missing for the individual symptoms (where CESCAT = “SIGNS AND SYMPTOMS OF DISEASE”), however, the date first symptom started can be found in the FACE dataset (FATESTCD = “FSYMDATE”) along with date last symptom resolved (FATESTCD = “LSYMDATE”), or indication that symptoms are ongoing (FATESTCD = “SYMONGO”).

Similarly, start and stop dates were not collected in the CRF for potential COVID-19 illness (records where CECAT = “EFFICACY” and CEOCCUR is null).

In the AE domain, there is only one event of tinnitus for subject C4591001 1056 10561120 found with missing timing information. Per query response, this AE was reported by the subject to a Help Desk and the subject subsequently withdrew consent. Only the AE term is known; no other details were provided.

2.9. CBER Comment 9

In your final analysis datasets, there are 9 records in the DS dataset missing Start Date/Time of Disposition Event (DSSTDTC) and Study Day of Start of Disposition Event (DSSTDY).

Response

At the time of data extraction, the study was still ongoing and data cleaning is still underway. Updated data will be included for future reporting events and at study completion.

2.10. CBER Comment 10

In your final analysis datasets there are 1,713 AE records that don't have AESEV or AETOXGR populated.

Response

For reactogenicity events (AECAT = "REACTOGENICITY"), severity was not collected after the diary period hence the severity is not reported in the AE domain (1364 records). This was communicated to the Agency in (b) (4) Response to the Information Request. The maximum severity reported during the diary period is reported in CESEV in the CE domain.

Additionally, toxicity grade was not collected for medication errors (326 records) (AECAT = "MEDICATION ERROR") and exposure during pregnancy (21 records) (AEDECOD = "Exposure during pregnancy" or "Maternal exposure during pregnancy").

The remaining two records with missing toxicity grade are due to incomplete data or unknown values at time of data cut (subjects 10561120/tinnitus and 12481121/vomiting).

We have 320 subjects with ADAE.AECAT = "MEDICATION ERROR". The breakdown for the preferred term is as follows.

AEDECOD	Frequency
Expired product administered	6
Incorrect dosage administered	80
Incorrect dose administered	18
Product administration error	8
Product dispensing error	1
Product storage error	141
Product temperature excursion issue	3
Wrong product administered	63

None of these medication errors subjects had an SAE.

For the same 320 subjects as per ADDS, breakdown for the disposition term are as shown below:

Based on ADDS.DSPHASE = "VACCINATION", 37 subjects discontinued from the vaccination phase.

DSDECOD	Frequency
COMPLETED	279
MEDICATION ERROR WITHOUT ASSOCIATED ADVERSE EVENT	5
NO LONGER MEETS ELIGIBILITY CRITERIA	1
WITHDRAWAL BY SUBJECT	31

Based on ADDS.DSPHASE = "FOLLOW-UP", 2 subjects discontinued from the study.

DSDECOD	Frequency
MEDICATION ERROR WITHOUT ASSOCIATED ADVERSE EVENT	1
WITHDRAWAL BY SUBJECT	1

2.11. CBER Comment 11

In your interim analysis datasets there is 1 record where the death date in the Death Detail domain is different from in the Disposition domain and in your final analysis datasets there are 3 records where the death date in the Death Detail domain is different from in the Disposition domain as indicated in the tables below.

DDDTC date is after RFPENDTC, death date in Death Detail domain is different from in Disposition domain.

Response

DDDTC (Date/Time of Collection) in the DD domain represents the Date of Collection/Notification of Death.

C4591001: DEATH DETAILS CODED (DEATH DTL) [DD002]	
Death Details [DD01002]	
1. * Date of Collection / Notification of Death: [Date of Collection / Notification of Death]	[DDDTC_002] (MM/DD/YYYY) Req / Req / Req (2020-2025)

Per SDTM IG 3.2 Section 6.3 Findings for Death Details (DD) (b) (6) Notes for DDDTC: Date/time of collection of the diagnosis or other death assessment data in ISO 8601 format. This is not necessarily the date of death.

Date of death is captured in the Disposition data (DS.DSSTDTC where DSDECOD = "DEATH") and additionally reported in DM.DTHDTC.

- Study c4591001-ia-efficacy - 1 records (33.33%)

USUBJID	DDTEST	DDORRES	DDDTTC	RFPENDTTC	Last Disposition date	DSDECOD
C4591001 1162 11621327	PRIMARY CAUSE OF	atherosclerotic disease	9/24/2020	9/13/2020	9/13/2020	DEATH

- Study c4591001-safety-fa-eff - 3 records (50%)

USUBJID	DDTEST	DDORRES	DDDTTC	RFPENDTTC	Last Disposition date	DSDECOD
C4591001 1066 10661350	PRIMARY CAUSE OF DEATH	myocardial infarction	11/10/2020	11/3/2020	11/3/2020	DEATH
C4591001 1081 10811194	PRIMARY CAUSE OF DEATH	Cause is currently unknown. Detail will be updated once the death certificate has been received	11/11/2020	11/1/2020	11/1/2020	DEATH
C4591001 1162 11621327	PRIMARY CAUSE OF DEATH	atherosclerotic disease	9/24/2020	9/13/2020	9/13/2020	DEATH

2.12. CBER Comment 12

In your interim analysis datasets there are 97 records missing MedDRA coding information and in your final analysis datasets there are 89 records missing MedDRA coding information.

Response

The study was ongoing at the time of data cutoff. Therefore, some AE terms were not yet coded in MedDRA. These terms were set to 'Uncoded' in AE dataset for transparency. These terms will continue to be cleaned and coded for future reporting events.

Appendix 1.

C4591001 EUA Safety Analysis – Key Safety Table Annotated Mocks

Table	File Name (.html, -sas.txt)	Title	Input Datasets	Population Subset used
1	adds-s002-p3- rand	Disposition of All Randomized Subjects – ~38000 Subjects for Phase 2/3 Analysis	ADDS ADSL	ADSL.RANDFL='Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and not(ADSL.SAFFL='Y' and NOT (ADSL.VAX101="" and ADSL.VAX102=""))
2	adce-s010-lr- age-p3-saf	Local Reactions, by Maximum Severity, Within 7 Days After Each Dose, by Age Group – Reactogenicity Subset for Phase 2/3 Analysis – Safety Population	ADFACEVD ADSL	ADAE.SAFFL='Y' and ADAE.HIVFL='N' and ADFACEVD.FAOBJ in ('PAIN AT INJECTION SITE' 'SWELLING' 'REDNESS') and ADFACEVD.KNOWVFL='Y' and ADSL.TRTA ne "" and ADSL.PHASEN NE 1 and ADSL.ARMCD ne "" and ADSL.AGEGR1N^=1 and ADSL.MULENRFL^='Y' and ADSL.DS30KFL='Y' and NOT (ADSL.VAX101="" and ADSL.VAX102="")
3	adce-s020-se- age-p3-saf	Systemic Events, by Maximum Severity, Within 7 Days After Each Dose, by Age Group – Reactogenicity Subset for Phase 2/3 Analysis – Safety Population	ADFACEVD ADSL	ADSL.SAFFL='Y' and ADSL.TRTA ne "" and ADFACEVD.FAOBJ not in ('PAIN AT INJECTION SITE' 'SWELLING' 'REDNESS') and index(uppercase(ADFACEVD.FAOBJ),'HOSPI')=0 and ADSL.KNOWVFL='Y' and ADSL.PHASEN NE 1 and ADSL.ARMCD ne "" and ADSL.AGEGR1N^=1 and ADSL.MULENRFL^='Y' and ADSL.DS30KFL='Y' and ADSL.HIVFL='N' and NOT (ADSL.VAX101="" and ADSL.VAX102="")
4	adae-s091-all- pd2-p3-saf	Number (%) of Subjects Reporting at Least 1 Adverse Event From Dose 1 to 1 Month After Dose 2 – ~38000 Subjects for Phase 2/3 Analysis – Safety Population	ADAE ADSL	ADSL.SAFFL='Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and ADSL.HIVFL ne 'Y' and NOT (ADSL.VAX101="" AND ADSL.VAX102="")
5	adae-s091-all- 2mpd2-p23-saf	Number (%) of Subjects Reporting at Least 1 Adverse Event From Dose 1 to Data Cutoff Date (14NOV2020) – Subjects With 2 Months Follow-Up Time After Dose 2 for Phase 2/3 Analysis – Safety Population	ADAE ADSL	ADSL.SAFFL='Y' and ADSL.FU2MPD2 >=56 and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.HIVFL ne 'Y' and NOT (ADSL.VAX101="" AND ADSL.VAX102="")
6	adae-vax-tier2- p3-saf	Tier 2 Adverse Events Reported From Dose 1 to 1 Month After Dose 2, by System Organ Class and Preferred Term – ~38000 Subjects for Phase 2/3 Analysis – Safety Population	ADAE ADSL	ADSL.SAFFL='Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and ADSL.HIVFL ne 'Y' and NOT (ADSL.VAX101="" AND ADSL.VAX102="")

Table	File Name (.html, -sas.txt)	Title	Input Datasets	Population Subset used
7	adae-s130-1md2-soc-p3-saf	Number (%) of Subjects Reporting at Least 1 Adverse Event From Dose 1 to 1 Month After Dose 2, by System Organ Class and Preferred Term – ~38000 Subjects for Phase 2/3 Analysis – Safety Population	ADAE ADSL	ADSL.SAFFL='Y' and ADSL.HIVFL ne 'Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and NOT (ADSL.VAX101=' AND ADSL.VAX102=')
8	adae-s130-1md2-ser-p3-saf	Number (%) of Subjects Reporting at Least 1 Serious Adverse Event From Dose 1 to 1 Month After Dose 2, by System Organ Class and Preferred Term – ~38000 Subjects for Phase 2/3 Analysis – Safety Population	ADAE ADSL	ADSL.SAFFL='Y' and ADSL.HIVFL ne 'Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and NOT (ADSL.VAX101=' AND ADSL.VAX102=')
9	adae-s130-ser-vax2-p3-saf	Number (%) of Subjects Reporting at Least 1 Serious Adverse Event From Dose 1 to Data Cutoff Date (14NOV2020), by System Organ Class and Preferred Term – ~38000 Subjects for Phase 2/3 Analysis – Safety Population	ADAE ADSL	ADSL.SAFFL='Y' and ADSL.HIVFL ne 'Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and NOT (ADSL.VAX101=' AND ADSL.VAX102=')

Table 1

Disposition of All Randomized Subjects – ~38000 Subjects for Phase 2/3 Analysis

ADSL.TRT01P

ADSL.RANDFL='Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and not(SAFFL ='Y' and (VAX101=''' and VAX102 ='''))

		Vaccine Group (as Randomized)		
		BNT162b2 (30 µg)	Placebo	Total
		(N^a=xx)	(N^a=xx)	(N^a=xx)
		n^b (%)	n^b (%)	n^b (%)
Randomized	ADSL.VAX101DT eq . and ADSL.VAX102DT eq .	xx (xxx.x)	xx (xxx.x)	xx (xxx.x)
Not vaccinated		xx (xxx.x)	xx (xxx.x)	xx (xxx.x)
Vaccinated	ADSL.VAX101DT ne .	xx (xxx.x)	xx (xxx.x)	xx (xxx.x)
Dose 1	ADSL.VAX102DT ne .			xx (xxx.x)
Dose 2				xx (xxx.x)
Completed 1-month post-Dose 2 visit (vaccination period)	ADDS.DSPHASEN=26 and ADDS.DSDECODN=2			xx (xxx.x)
Discontinued from vaccination period but continue in the study	ADDS.DSPHASEN=26 and ADDS.EOTDCDT ne . and ADDS.EOSDCDT eq . and ADDS.DSDECODN not in (. 2) and (ADSL.VAX101DT ne . or ADSL.VAX102DT ne .)			
Discontinued after Dose 1 and before Dose 2	ADSL.VAX101DT ne . and (ADSL.VAX102DT eq . or ADDS.ASTDT < ADSL.VAX102DT)			
Discontinued after Dose 2 and before 1-month post-Dose 2 visit	ADSL.VAX101DT ne . and ADSL.VAX102DT ne . and (ADSL.VAX102DT <= ASTDT and (M1PD2DT eq . or ASTDT<M1PD2DT)			
Reason for discontinuation from vaccination period	ADDS.DSDECOD			
Adverse event				
Withdrawal by subject				
Physician decision		xx (xxx.x)	xx (xxx.x)	xx (xxx.x)
Death		xx (xxx.x)	xx (xxx.x)	xx (xxx.x)
Study terminated by sponsor		xx (xxx.x)	xx (xxx.x)	xx (xxx.x)
Pregnancy				
Other				
Withdrawal from the study	ADDS.DSPHASEN=31 and ADDS.EOSDCDT ne . and ADDS.DSDECOD not in (. 2) and (ADSL.VAX101DT ne . or ADSL.VAX102DT ne .)			
Withdrawal after Dose 1 and before Dose 2	ADSL.VAX101DT ne . and (ADSL.VAX102DT eq . or ADDS.ASTDT < ADSL.VAX102DT)			
Withdrawal after Dose 2 and before 1-month post-Dose 2 visit	ADSL.VAX101DT ne . and ADSL.VAX102DT ne . and (ADSL.VAX102DT <= ADDS.ASTDT and (ADDS.M1PD2DT eq . or ADDS.ASTDT<ADDS.M1PD2DT)			
Withdrawal after 1-month post-Dose 2 visit				
Reason for withdrawal from the study				
Adverse event				
Withdrawal by subject				
Physician decision		xx (xxx.x)	xx (xxx.x)	xx (xxx.x)
Death	ADSL.VAX101DT ne . and ADSL.VAX102DT ne . and ADDS.M1PD2DT ne . and ADSL.M1PD2DT le ADDS.ASTDT			

PFIZER CONFIDENTIAL

Pregnancy	xx (xxx.x)	xx (xxx.x)	xx (xxx.x)
Other	xx (xxx.x)	xx (xxx.x)	xx (xxx.x)

Note: HIV-positive subjects are included in this summary but not included in the analyses of the overall study objectives.

Note: Subject C4591001 xxx was randomized but did not sign informed consent and is not included in any analysis population.

Note: Because of a dosing error, Subject[s] C4591001 xxxx xxxxx [and C4591001 xxxx xxxxxx] received an additional dose of BNT162b2 (30 µg) at an unscheduled visit after receiving one dose of BNT162b2 (30 µg) and one dose of placebo.

a. N = number of randomized subjects in the specified group, or the total sample. This value is the denominator for the percentage calculations.

b. n = Number of subjects with the specified characteristic.

PFIZER CONFIDENTIAL SDTM Creation: DDMMYYYY (HH:MM) Source Data: abcdefgh Table Generation: DDMMYYYY (HH:MM)

(Cutoff date: ddMmmYYYY, Snapshot Date: ddMmmYYYY) Output File: (b) (6)/C4591001/abcd_XNNN

Table 2. **Local Reactions, by Maximum Severity, Within 7 Days After Each Dose, by Age Group – Reactogenicity**
Subset for Phase 2/3 Analysis – Safety Population

AGEGRIN		ADFACEVD.ATPTREF			Vaccine Group (as Administered)			ADSL.TRT01A	
Age Group	Dose	Local Reaction	N ^a	n ^b (%)	(95% CI ^c)	N ^a	n ^b (%)	(95% CI ^c)	
1	Any	Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	ADSL.SAFFL='Y' and ADSL.HIVFL='N' and ADFACEVD.FAOBJ in ('PAIN AT INJECTION SITE' 'SWELLING' 'REDNESS') and ADFACEVD.KNOWVFL='Y' and ADSL.TRTA ne '' and ADSL.PHASEN ne 1 and ADSL.ARMCD ne '' and ADSL.AGEGR1N^=1 and ADSL.MULENRFL^='Y' and ADSL.DS30KFL='Y' and NOT (ADSL.VAX101=''' and ADSL.VAX102='''
		Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Severe	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Grade 4	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Swelling ^d							
		Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Severe	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Grade 4	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Pain at the injection site ^e							
		Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Severe	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Grade 4	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
		Any local reaction ^f	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)	
	2 <Repeat for Dose 2>								
	Any dose<Repeat for any dose>								
	<Repeat for >55 Years>								

ADFACEVD.FATESTCD='MAXSEV'
and AVALC ne'' and EVENTFL='Y'

ADSL.SAFFL='Y' and ADSL.HIVFL='N' and ADFACEVD.FAOBJ in
('PAIN AT INJECTION SITE' 'SWELLING' 'REDNESS') and
ADFACEVD.KNOWVFL='Y' and ADSL.TRTA ne ' and ADSL.PHASEN
ne 1 and ADSL.ARMCD ne '' and ADSL.AGEGR1N^=1 and
ADSL.MULENRFL^='Y' and ADSL.DS30KFL='Y' and NOT
(ADSL.VAX101='' and ADSL.VAX102='')

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose.

Note: Grade 4 reactions were classified by the investigator or medically qualified person.

- a. N = number of subjects reporting at least 1 yes or no response for the specified reaction after the specified dose.
- b. n = Number of subjects with the specified characteristic.
- c. Exact 2-sided CI based on the Clopper and Pearson method.
- d. Mild: >2.0 to 5.0 cm; moderate: >5.0 to 10.0 cm; severe: >10.0 cm; Grade 4: necrosis (redness and swelling categories) or exfoliative dermatitis (redness category only).
- e. Mild: does not interfere with activity; moderate: interferes with activity; severe: prevents daily activity; Grade 4: emergency room visit or hospitalization for severe pain at the injection site.
- f. Any local reaction: any redness >2.0 cm, any swelling >2.0 cm, or any pain at the injection site.

PFIZER CONFIDENTIAL SDTM Creation: DDMMYYYY (HH:MM) Source Data: abcdefgh Table Generation: DDMMYYYY (HH:MM)

(Cutoff date: ddMmmYYYY, Snapshot Date: ddMmmYYYY) Output File: (b) (6) /C4591001/abcd_XNNN

Table 3.

Systemic Events by Maximum Severity, Within 7 Days After Each Dose, by Age Group – Reactogenicity Subset for Phase 2/3 Analysis – Safety Population

ADSL.AGEGR1N

ADFACEVD.FATESTCD="MAXSEV" and AVALC ne "" and
EVENTFL="Y"

ADSL.TRT01A

ADFACEVD.ATPTREF

ADSL.SAFFL="Y" and ADSL.TRTA ne "" and ADFACEVD.FAOBJ not
in ("PAIN AT INJECTION SITE" "SWELLING" "REDNESS") and
index(uppercase(ADFACEVD.FAOBJ),"HOSPI")=0 and
ADSL.KNOWVFL="Y" and ADSL.PHASEN ne 1 and ADSL.ARMCD ne
" and ADSL.AGEGR1N^=1 and ADSL.MULENRFL^="Y" and
ADSL.DS30KFL="Y" and ADSL.HIVFL="N" and NOT (ADSL.VAX101=""
and ADSL.VAX102="")

ADFACEVD.FAOBJ in FAOBJ not in ("PAIN
AT INJECTION SITE" "SWELLING"
"REDNESS") and
index(uppercase(ADFACEVD.FAOBJ),"HOSPI")=
0

Age Group	Dose	Systemic Event	Vaccine Group (as Administered)			Placebo		
			N ^a	n ^b (%)	(95% CI) ^c	N ^a	n ^b (%)	(95% CI) ^c
16-55 Years	1	Fever						
		≥38.0°C	NN	nn (xx.x)		NN	nn (xx.x)	
	2	≥38.0°C to 38.4°C	NN	nn (xx.x)		NN	nn (xx.x)	
		≥38.0°C to 38.9°C	NN	nn (xx.x)		NN	nn (xx.x)	
	3	>38.9°C	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
		>40.0°C	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
	4	Fatigue ^d						
		Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
		Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
		Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
	5	Headache ^d						
		Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
		Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
		Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
	6	Chills ^d						
		Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
		Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
		Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
	7	Vomiting ^e						
		Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
		Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
		Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)

Severe	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Grade 4	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Diarrhea ^f						
Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Severe	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Grade 4	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
New or worsened muscle pain ^d						
Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Severe	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Grade 4	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
New or worsened joint pain ^d						
Any	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Mild	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Moderate	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Severe	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Grade 4	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Any systemic event ^g	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
Use of antipyretic or pain medication ^h	NN	nn (xx.x)	(xx.x, xx.x)	NN	nn (xx.x)	(xx.x, xx.x)
2 <Repeat for Dose 2>						
Any dose <Repeat for any dose>						
<Repeat for >55 Years>						

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose. Grade 4 events were classified by the investigator or medically qualified person.

- a. N = number of subjects reporting at least 1 yes or no response for the specified event after the specified dose.
- b. n = Number of subjects with the specified characteristic.
- c. Exact 2-sided CI based on the Clopper and Pearson method.
- d. Mild: does not interfere with activity; moderate: some interference with activity; severe: prevents daily activity; Grade 4: emergency room visit or hospitalization for severe fatigue, severe headache, severe muscle pain, or severe joint pain.
- e. Mild: 1 to 2 times in 24 hours; moderate: >2 times in 24 hours; severe: requires intravenous hydration; Grade 4: emergency room visit or hospitalization for severe vomiting.
- f. Mild: 2 to 3 loose stools in 24 hours; moderate: 4 to 5 loose stools in 24 hours; severe: 6 or more loose stools in 24 hours; Grade 4: emergency room visit or hospitalization for severe diarrhea.
- g. Any systemic event: any fever $\geq 38.0^{\circ}\text{C}$, any fatigue, any vomiting, any chills, any diarrhea, any headache, any new or worsened muscle pain, or any new or worsened joint pain.
- h. Severity was not collected for use of antipyretic or pain medication.

PFIZER CONFIDENTIAL SDTM Creation: DDMMYYYY (HH:MM) Source Data: abcdefgh Table Generation: DDMMYYYY (HH:MM)
(Cutoff date: ddMmmYYYY, Snapshot Date: ddMmmYYYY) Output File: (b) (6)/C4591001/abcd_XNNN

Table 4:

AECAT='ADVERSE EVENT' and ADSL.SAFFL='Y' and ADAE.VPHASEN in (1,2,3) and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and ADSL.V01DT >= ADAE.ASTDT

ADSL.SAFFL="Y" and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne "Y" and ADSL.DS30KFL="Y" and ADSL.HIVFL ne 'Y' and NOT ((ADSL.VAX101=' ' AND ADSL.VAX102=' '))

**Number (%) of Subjects Reporting at Least 1 Adverse Event From Dose 1 to 1 Month After Dose 2 –
~38000 Subjects for Phase 2/3 Analysis – Safety Population**

		Vaccine Group (as Administered)	
		BNT162b2 (30 µg) (N^a=xx)	Placebo (N^a=xx)
Adverse Event	ADSL.TRT01A	n^b (%)	n^b (%)
Any event	Upcase(ADAE.AREL)="RELATED"	xx (xx.x)	xx (xx.x)
Related ^c		xx (xx.x)	xx (xx.x)
Severe	ADAE.ATOXGRN=3	xx (xx.x)	xx (xx.x)
Life-threatening	ADAE.ATOXGRN=4	xx (xx.x)	xx (xx.x)
Any serious adverse event		xx (xx.x)	xx (xx.x)
Related ^c	ADAE.AESER="Y"	xx (xx.x)	xx (xx.x)
Severe		xx (xx.x)	xx (xx.x)
Life-threatening		xx (xx.x)	xx (xx.x)
Any adverse event leading to withdrawal	index (upcase(ADAE.AEACN),'DRUG WITHDRAWN') > 0 or ADAE.AESUBJDC='Y'	xx (xx.x)	xx (xx.x)
Related ^c		xx (xx.x)	xx (xx.x)
Severe		xx (xx.x)	xx (xx.x)
Life-threatening	ADAE.AESDTH='Y' or ADAE.AEOUT='FATAL'	xx (xx.x)	xx (xx.x)
Death		xx (xx.x)	xx (xx.x)

- a. N = number of subjects in the specific vaccine group or for the percentage calculations.
b. n = Number of subjects reporting at least 1 occurrence of the specified event category. For "any event", n = the number of subjects reporting at least 1 occurrence of any event.
c. PFIZER CONFIDENTIAL SDTM Creation: DDMMYYYY (HH:MM) Source Data: abcdefgh Table Generation: DDMMYYYY (HH:MM) (Cutoff date: ddMmmYYYY, Snapshot Date: ddMmmYYYY) Output File: (b) (6) /C4591001/abcd_XNNN

Table 5

ADAE.AECAT='ADVERSE EVENT' and
ADSL.SAFFL='Y' and ADAE.VPHASEN >=1 and
ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1
and ADSL.MULENRFL ne 'Y' and ADSL.HIVFL
ne 'Y'

ADSL.SAFFL='Y' and ADSL.FU2MPD2 >=56 and ADSL.PHASEN ne 1 and
ADSL.AGEGR1N ne 1 and MULENRFL ne 'Y' and HIVFL ne 'Y' and NOT
((VAX101=' AND VAX102='))

Number (%) of Subjects Reporting at Least 1 Adverse Event From Dose 1 to Data Cutoff Date (14NOV2020) –
Subjects With 2 Months Follow-Up Time After Dose 2 for Phase 2/3 Analysis – Safety Population

ADSL.TRT01A

Adverse Event	Vaccine Group (as Administered)		
	BNT162b2 (30 µg) (N ^a =xx) n ^b (%)	Placebo (N ^a =xx) n ^b (%)	Total (N ^a =xx) n ^b (%)
Any event	Upcase(ADAE.AREL)="RELATED" (xx.x)	xx (xx.x)	xx (xx.x)
Related ^c	xx (xx.x)	xx (xx.x)	xx (xx.x)
Severe	ADAE.ATOXGRN=3 xx (xx.x)	xx (xx.x)	xx (xx.x)
Life-threatening	ADAE.ATOXGRN=4 xx (xx.x)	xx (xx.x)	xx (xx.x)
Any serious adverse event	xx (xx.x)	xx (xx.x)	xx (xx.x)
Related ^c	ADAE.AESER="Y" xx (xx.x)	xx (xx.x)	xx (xx.x)
Severe	xx (xx.x)	xx (xx.x)	xx (xx.x)
Life-threatening			xx (xx.x)
Any adverse event leading to withdrawal	index (upcase(ADAE.AEACN),'DRUG WITHDRAWN') > 0 or ADAE.AESUBJDC='Y' xx (xx.x)		xx (xx.x)
Related ^c			xx (xx.x)
Severe	xx (xx.x)	xx (xx.x)	xx (xx.x)
Life-threatening		xx (xx.x)	xx (xx.x)
Death	ADAE.AESDTH='Y' or ADAE.AEOUT='FATAL' xx (xx.x)	xx (xx.x)	xx (xx.x)

a. N = number of subjects in the specific vaccine group.

b. n = Number of subjects reporting at least 1 occurrence of the specified event category. For "any event", n = the number of subjects reporting at least 1 occurrence of any event.

c. Assessed by the investigator as related to investigational product.

PFIZER CONFIDENTIAL SDTM Creation: DDMMYYYY (HH:MM) Source Data: abcdefgh Table Generation: DDMMYYYY (HH:MM)

(Cutoff date: ddMmmYYYY, Snapshot Date: ddMmmYYYY) Output File: ((b) (6)/C4591001/abcd_XNNN

Table 6

ADAE.AECAT = 'ADVERSE EVENT' and
ADAE.VPHASEN in (1,2) and TIER2FL='Y'

ADSL.SAFFL = 'Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1
and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and ADSL.HIVFL
ne 'Y' and NOT (ADSL.VAX101=' AND ADSL.VAX102=')

Tier 2 Adverse Events Reported From Dose 1 to 1 Month After Dose 2, by System Organ Class and Preferred Term – ~38000 Subjects

Phase 2/3 Analysis – Safety Population							
Vaccine Group (as Administered)							
BNT162b2 (30 µg) (N ^a =xx)			Placebo (N ^a =xx)			Difference	
System Organ Class	n ^b (%)	(95% CI) ^c	n ^b (%)	(95% CI) ^c	% ^d	(95% CI) ^e	
Preferred Term							
<System organ class>							
<Preferred term>	xx (xx.x)	(xx.x, xx.x)					
		(xx.x, xx.x)					
		(xx.x, xx.x)					
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)	xx.x	(xx.x, xx.x)	

TIER2FL derived as below:
create table per as select a.*,b.bign,a.n/b.bign*100 as per,'Y' as
TIER2FL from smaln a left join bign b on a.ACTARM=b.ACTARM and
a.PHASENX=b.PHASENX having per>=1;

Method Used:
proc binomial data=XXXX out=XXXX1 max_time=120 alpha=0.95 ;
by _event;
RISKDIFF / AS ONE STD;
popl_trt_id;
outcome_exist;
run;

Note: MedDRA (v23.1) coding dictionary applied.

Note: Tier 2 events are "common" adverse events with an incidence rate $\geq 1.0\%$ in any vaccine group (preferred term level). No Tier 1 events were identified at this stage for this program.

Note: The 95% confidence interval quantifies the precision of the risk difference estimate. Confidence intervals are not adjusted for multiplicity. They should only be used to identify potentially important adverse events.

- N = number of subjects in the specified group. This value is the denominator for the percentage calculations.
- n = Number of subjects reporting at least 1 occurrence of the specified adverse event.
- Exact 2-sided CI based on the Clopper and Pearson method.
- Difference in proportions, expressed as a percentage (BNT162b2 [30 µg] – placebo) for subjects ≥ 16 years.
- 2-Sided CI, based on the Miettinen and Nurminen method for the difference in proportions, expressed as a percentage.

PFIZER CONFIDENTIAL SDTM Creation: DDMMYY (HH:MM) Source Data: ADSL Table Generation: DDMMYY (HH:MM)

(Cutoff date: ddMmmYYYY, Snapshot Date: ddMmmYYYY) Output File: (b) (6) /C4591001/abcd_XNNN

TABLE 7

Number (%) of Subjects Reporting at Least 1 Adverse Event From Dose 1 to 1 Month After Dose 2, by System Class and Preferred Term – ~38000 Subjects for Phase 2/3 Analysis – Safety Population				
System Organ Class Preferred Term	Vaccine Group (as Administered)			Placebo
	BNT162b2 (30 µg) (N ^a =xx)			(N ^a =xx)
	n ^b (%)	(95% CI) ^c	n ^b (%)	(95% CI) ^c
<Any event>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<System organ class>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<System organ class>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)

ADAE.AECAT = 'ADVERSE EVENT' and
ADSL.SAFFL='Y' and ADAE.VPHASEN in (1,2,3) and
ADSL.PHASEN ne 1 ADSL.AGEGR1N ne 1 and
ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and
ADSL.V01DT >= ADAE.ASTDT

ADSL.SAFFL='Y' and ADSL.HIVFL ne 'Y' and
ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1
and ADSL.MULENRFL ne 'Y' and
ADSL.DS30KFL='Y') and NOT
((ADSL.VAX101=' AND ADSL.VAX102='))

ADAE.AEBODSYS

ADAE.AEDECOD

ADSL.TRT01A

Note: MedDRA (v23.1) coding dictionary applied.

a. N = number of subjects in the specified group. This value is the denominator for the percentage calculations.

b. n = Number of subjects reporting at least 1 occurrence of the specified event. For "any event", n = number of subjects reporting at least 1 occurrence of any event.

c. Exact 2-sided CI based on the Clopper and Pearson method.

PFIZER CONFIDENTIAL SDTM Creation: DDMMYYYY (HH:MM) Source Data: ADSL Table Generation: DDMMYYYY (HH:MM)

(Cutoff date: ddMmmYYYY, Snapshot Date: ddMmmYYYY) Output File: (b) (6) C4591001/abcd_XNNN

PFIZER CONFIDENTIAL

Page 25

FDA-CBER-2021-5683-1788614

090177e1959f4bcfApproved On: 26-Nov-2020 01:08 (GMT)

TABLE 8

ADAE.AECAT = 'ADVERSE EVENT' and ADAE.AESER='Y' and ADAE.SAFFL='Y' and ADAE.VPHASEN in (1,2,3) and ADAE.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADAE.MULENRFL ne 'Y' and ADAE.DS30KFL='Y' and ADSL.V01DT >= ADAE.ASTDT
ADAE.AECAT = 'ADVERSE EVENT' and ADAE.AESER='Y' and ADSL.SAFFL='Y' and ADAE.VPHASEN in (1,2,3) and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and ADSL.V01DT >= ADAE.ASTDT

ADSL.SAFFL='Y' and ADSL.HIVFL ne 'Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y' and NOT ((ADSL.VAX101=' AND ADSL.VAX102='))

Number (%) of Subjects Reporting at Least 1 Serious Adverse Event From Dose 1 to 1 Month After Dose 2, by System Organ Class and Preferred Term – ~38000 Subjects for Phase 2/3 Analysis – Safety Population

System Organ Class Preferred Term	Vaccine Group (as Administered)			
	BNT162b2 (30 µg) (N ^a =xx)		Placebo (N ^a =xx)	
	n ^b (%)	(95% CI) ^c	n ^b (%)	(95% CI) ^c
<Any event>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<System organ class>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<System organ class>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)

Note: MedDRA (v23.1) coding dictionary applied.

a. N = number of subjects in the specified group. This value is the denominator for the percentage calculations.

b. n = Number of subjects reporting at least 1 occurrence of the specified event. For "any event", n = number of subjects reporting at least 1 occurrence of any event.

c. Exact 2-sided CI based on the Clopper and Pearson method.

PFIZER CONFIDENTIAL SDTM Creation: DDMMYYYY (HH:MM) Source Data: ADSL Table Generation: DDMMYYYY (HH:MM)

(Cutoff date: ddMmmYYYY, Snapshot Date: ddMmmYYYY) Output File: (b) (6) C4591001/abcd_XNNN

TABLE 9

ADAE.AECAT = 'ADVERSE EVENT' and ADAE.AESER='Y' and ADSL.SAFFL='Y' and ADAE.VPHASEN in (1,2,3,99) and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y'

ADSL.SAFFL='Y' and ADSL.HIVFL ne 'Y' and ADSL.PHASEN ne 1 and ADSL.AGEGR1N ne 1 and ADSL.MULENRFL ne 'Y' and ADSL.DS30KFL='Y') and NOT ((ADSL.VAX101=' AND ADSL.VAX102='))

Number (%) of Subjects Reporting at Least 1 Serious Adverse Event From Dose 1 to Data Cutoff Date (14NOV2020), by System Organ Class and Preferred Term -- 38000 Subjects for Phase 2/3 Analysis
Safety Population

ADAE.AEBODSYS
ADAE.AEDECOD

ADSL.TRT01A

System Organ Class Preferred Term	Vaccine Group (as Administered)			
	BNT162b2 (30 µg) (N ^a =xx)		Placebo (N ^a =xx)	
	n ^b (%)	(95% CI ^c)	n ^b (%)	(95% CI ^c)
<Any event>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<System organ class>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<System organ class>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)
<Preferred term>	xx (xx.x)	(xx.x, xx.x)	xx (xx.x)	(xx.x, xx.x)

Note: MedDRA (v23.1) coding dictionary applied.

a. N = number of subjects in the specified group. This value is the denominator for the percentage calculations.

b. n = Number of subjects reporting at least 1 occurrence of the specified event. For "any event", n = number of subjects reporting at least 1 occurrence of any event.

c. Exact 2-sided CI based on the Clopper and Pearson method.

PFIZER CONFIDENTIAL SDTM Creation: DDMMYYYY (HH:MM) Source Data: ADSL Table Generation: DDMMYYYY (HH:MM)
(Cutoff date: ddMmmYYYY, Snapshot Date: ddMmmYYYY) Output File: (b) (6) C4591001/abcd_XNNN

Document Approval Record

Document Name:	COVID-19 2020 Response to 24Nov20 CBER Comments - Datasets
Document Title:	COVID-19 2020 Response to 24Nov20 CBER Comments - Datasets

Signed By:	Date(GMT)	Signing Capacity
(b) (4), (b) (6)	26-Nov-2020 00:29:09	Business Line Approver
Perez, John	26-Nov-2020 01:08:00	Final Approval